

Afghanistan



Country report card

This report card contains the latest data available on the Global Obesity Observatory on overweight and obesity . Where available, data on common and relevant obesity drivers and comorbidities are also presented.




Contents	Page
Obesity prevalence	3
Overweight/obesity by education	6
Overweight/obesity by age	9
Overweight/obesity by region	11
Overweight/obesity by socio-economic group	13
Double burden of underweight & overweight	16
Insufficient physical activity	19
Average daily frequency of carbonated soft drink consumption	25
Estimated per capita fruit intake	26
Prevalence of less than daily fruit consumption	27
Prevalence of less than daily vegetable consumption	28
Average weekly frequency of fast food consumption	29
Estimated per-capita processed meat intake	30
Estimated per capita whole grains intake	31
Mental health - depression disorders	32
Mental health - anxiety disorders	38
% Infants exclusively breastfed 0-5 months	43
Oesophageal cancer	44
Breast cancer	46
Colorectal cancer	47
Pancreatic cancer	49
Gallbladder cancer	51
Kidney cancer	53
Cancer of the uterus	55
Raised blood pressure	56
Raised cholesterol	59
Raised fasting blood glucose	62
Diabetes prevalence	64
Ovarian Cancer	65
Leukemia	66
Leukemia	68
Liver and intrahepatic bile duct Cancer	70
Multiple Myeloma	72
Non Hodgkin Lymphoma	74
Thyroid Cancer	76



Obesity prevalence

Adults, 2018






Survey type:	Measured
Age:	18-69
Sample size:	3666
Area covered:	National
References:	Afghanistan STEPS Survey 2018 https://extranet.who.int/ncdsmicrodata/index.php/catalog/782 (Last Accessed 03.11.2020)
	Unless otherwise noted overweight refers to a RMI between 25kg and 29 9kg/m² obesity refers to a RMI greater than 30kg/m²

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².



Children, 2014






Survey type:	Self-reported
Age:	13-17
Sample size:	2579
Area covered:	National
References:	Global School-based Student Health Survey (GSHS), available at https://www.who.int/ncds/surveillance/gshs/2014-GSHS-Afghanistan-fact-sheet.pdf?ua=1 (last accessed 25.11.20)
Notes:	WHO cutoffs.
Cutoffs:	WHO



0-5 years, 2018

Overweight or obesity



References: Other: Afghanistan Health Survey 2018

Notes:

UNICEF/WHO/World Bank Joint Child Malnutrition Estimates Expanded Database: Overweight (Survey Estimates), May 2023, New York.

For more information about the methodology, please consult https://data.unicef.org/resources/jme-2023-country-consultations/
Percentage of children under 5 years of age falling above 2 standard deviations (moderate and severe) from the median weight-forheight of the reference population.


Definitions: =>+2SD



Overweight/obesity by education

Adults, 2013

Obesity




Survey type:	Measured
Age:	25+
Sample size:	1200
Area covered:	Jalalabad City
References:	Saeed KMI. Prevalence and associated risk factors for Obesity in Jalabad city - Afghanistan. Alexandria Journal of Medicine (2015) 51, 347–352

 $Unless \ otherwise \ noted, overweight \ refers \ to \ a \ BMI \ between \ 25kg \ and \ 29.9kg/m^2, obesity \ refers \ to \ a \ BMI \ greater \ than \ 30kg/m^2.$



Girls, 2013




=>+2SD



0-5 years, 2018

Definitions:



Measured




Overweight/obesity by age

Women, 2013

Survey type:





Sample size:	11124
Area covered:	National
References:	National Nutrition Survey Afghanistan 2013. Survey Report. Available from:

https://reliefweb.int/sites/reliefweb.int/files/resources/Report%20NNS%20Afghanistan%202013%20(July%2026-14).pdf. [Last


Accessed 16th January 2019].

Notes: Demographic Health Survey data includes ever married women aged 15-49 years only and may include males aged 15-59.

 $Unless \ otherwise \ noted, overweight \ refers \ to \ a \ BMI \ between \ 25kg \ and \ 29.9kg/m^2, obesity \ refers \ to \ a \ BMI \ greater \ than \ 30kg/m^2.$



Girls, 2013






Overweight/obesity by region

Adults, 2015

Obesity



Sample size: 1172

Area covered: Regional - Kabul

References:


 $Profile of risk factors for noncommunicable diseases in major cities of Afghanistan: WHO STEPwise approach. \\ \underline{https://applications.emro.who.int/emhj/v26/04/10203397-2020-2604-388-399.pdf} \ (last accessed 19.05.2022) \\ \underline{https://applications.emro.who.int/emhj/v26/04$

 $Unless \ otherwise \ noted, overweight \ refers \ to \ a \ BMI \ between \ 25kg \ and \ 29.9kg/m^2, obesity \ refers \ to \ a \ BMI \ greater \ than \ 30kg/m^2.$



0-5 years, 2018

Overweight or obesity



Sample size: 19692

References:

Other: Afghanistan Health Survey 2018

Notes:

UNICEF/WHO/World Bank Joint Child Malnutrition Estimates Expanded Database: Overweight (Survey Estimates), May 2023, New York.


For more information about the methodology, please consult https://data.unicef.org/resources/jme-2023-country-consultations/
Percentage of children under 5 years of age falling above 2 standard deviations (moderate and severe) from the median weight-forheight of the reference population.

Definitions: =>+2SD



Overweight/obesity by socio-economic group


Women, 2013





Girls, 2013





Survey type:	Measured
Age:	15-19
Sample size:	5808

Area covered: National

References: National Nutrition Survey Afghanistan 2013. Survey Report. Available from: https://reliefweb.int/sites/reliefweb.int/files/resources/Report%20NNS%20Afghanistan%202013%20(July%2026-14).pdf. [Last

https://reliefweb.int/sites/reliefweb.int/files/resources/Report%20NNS%20Afghanistan%202013%20[July%2026-14].pdf. [Last Accessed 16th January 2019].


Notes: WHO cut-offs used.

Cutoffs: WHO



0-5 years, 2018

Overweight or obesity



Sample size: 19692

References:

Other: Afghanistan Health Survey 2018

Notes:


UNICEF/WHO/World Bank Joint Child Malnutrition Estimates Expanded Database: Overweight (Survey Estimates), May 2023, New York.

For more information about the methodology, please consult https://data.unicef.org/resources/jme-2023-country-consultations/
Percentage of children under 5 years of age falling above 2 standard deviations (moderate and severe) from the median weight-forheight of the reference population.

Definitions: =>+2SD




Double burden of underweight & overweight





Children, 2022



Age:			5-19

References: NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population representative studies with 222 million children, adolescents, and adults. Lancet 2024; published online Feb 29. https://doi.org/10.1016/S0140-6736(23)02750-2.


Notes:	Age standardised estimates

Definitions: Combined prevalence of BMI < -2SD and BMI > 2SD (double burden of thinness and obesity)

Cutoffs: BMI < -2SD and BMI > 2SD



0-5 years, 2018-2018



Age: 0-5

References:


UNICEF data available at https://data.unicef.org/topic/nutrition/malnutrition/ (last accessed 12.03.25)

Notes:

Children under 5 falling below -2 standard deviations from the median height for age and falling at or above +2 standard deviations from the median weight-for-height of the reference population




Insufficient physical activity





Men, 2022



References:

WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at <a href="https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-aged-type-details/GHO/prevalence-


18-years-(age-standardized-estimate)-(-)

Definitions:

Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.



Women, 2022



References:

WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at <a href="https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-type-details/GHO/prevalence-of-insufficient-physical-activity-aged-type-details/GHO/prevalence-


18-years-(age-standardized-estimate)-(-)

Definitions:

Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.




Children, 2016



Survey type:	Self-reported
Age:	11-17
References:	Global Health Observatory data repository, World Health Organisation, https://apps.who.int/gho/data/node.main.A893ADO?lang=en (last accessed 16.03.21)
Notes:	% of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.
Definitions:	% Adolescents insufficiently active (age standardised estimate)




Boys, 2016



Survey type:	Self-reported
Age:	11-17
References:	Global Health Observatory data repository, World Health Organisation, https://apps.who.int/gho/data/node.main.A893ADO?lang=en (last accessed 16.03.21)
Notes:	% of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.
Definitions:	% Adolescents insufficiently active (age standardised estimate)



Girls, 2016




Survey type:	Self-reported
Age:	11-17
References:	Global Health Observatory data repository, World Health Organisation, https://apps.who.int/gho/data/node.main.A893ADO?lang=en (last accessed 16.03.21)
Notes:	% of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.
Definitions:	% Adolescents insufficiently active (age standardised estimate)



Average daily frequency of carbonated soft drink consumption

Children, 2009-2015



Survey type: Measured


Age: 12-17

References:

Beal et al. (2019). Global Patterns of Adolescent Fruit, Vegetable, Carbonated Soft Drink, and Fast-food consumption: A meta-analysis of global school-based student health surveys. Food and Nutrition Bulletin. https://doi.org/10.1177/0379572119848287 sourced from Food Systems Dashboard https://www.foodsystemsdashboard.org/food-system




Estimated per capita fruit intake





Prevalence of less than daily fruit consumption

Children, 2008-2015



Age: 12-17

References:

Global School-based Student Health Surveys. Beal et al (2019). Global Patterns of Adolescent Fruit, Vegetable, Carbonated Soft Drink, and Fast-food consumption: A meta-analysis of global school-based student health surveys. Food and Nutrition Bulletin. https://doi.org/10.1177/0379572119848287. Sourced from Food Systems Dashboard http://www.foodsystemsdashboard.org/food-

system


Definitions:

 $\label{prevalence} Prevalence\ of\ less-than-daily\ fruit\ consumption\ (\%\ less-than-daily\ fruit\ consumption)$



Prevalence of less than daily vegetable consumption

Children, 2008-2015



Survey type: Measured

Age: 12-17

References:

Beal et al. (2019). Global Patterns of Adolescent Fruit, Vegetable, Carbonated Soft Drink, and Fast-food consumption: A meta-analysis of global school-based student health surveys. Food and Nutrition Bulletin. https://doi.org/10.1177/0379572119848287 sourced from Food Systems Dashboard https://www.foodsystemsdashboard.org/food-system


Definitions:

Prevalence of less-than-daily vegetable consumption (% less-than-daily vegetable consumption)



Average weekly frequency of fast food consumption

Children, 2009-2015




References:

Beal et al. (2019). Global Patterns of Adolescent Fruit, Vegetable, Carbonated Soft Drink, and Fast-food consumption: A meta-analysis of global school-based student health surveys. Food and Nutrition Bulletin. https://doi.org/10.1177/0379572119848287 sourced from Food Systems Dashboard https://www.foodsystemsdashboard.org/food-systems




Estimated per-capita processed meat intake






Estimated per capita whole grains intake






Mental health - depression disorders






Men, 2021






Women, 2021





Children, 2021



Area covered: National

References:


Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from http://vizhub.healthdata.org/gbd-compare. (Last accessed 23.04.25)

Definitions:

Number living with depressive disorder per 100,000 population (Under 20 years of age) $\,$



Boys, 2021



Area covered: National

References:


Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from http://vizhub.healthdata.org/gbd-compare. (Last accessed 23.04.25)

Definitions:

Number living with depressive disorder per 100,000 population (Under 20 years of age) $\,$



Girls, 2021



Area covered: National

References:

Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from http://vizhub.healthdata.org/gbd-compare. (Last accessed 23.04.25)

Definitions:

Number living with depressive disorder per 100,000 population (Under 20 years of age) $\,$



Mental health - anxiety disorders

Adults, 2021




Number living with anxiety per 100,000 population




Men, 2021

Definitions:








References: Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from http://vizhub.healthdata.org/gbd-compare. (Last accessed 23.04.25).

Definitions:Number living with anxiety per 100,000 population




Children, 2021



References:

Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from http://vizhub.healthdata.org/gbd-compare. (Last accessed 23.04.25)

Boys, 2021




References:

Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from http://vizhub.healthdata.org/gbd-compare. (Last accessed 23.04.25)



Girls, 2021




References:

Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from http://vizhub.healthdata.org/gbd-compare. (Last accessed 23.04.25)



% Infants exclusively breastfed 0-5 months

0-5 years, 2010-2023



References:

Afghanistan Multiple Indicator Cluster Survey 2022-23, Summary Findings Report

Notes:

Full details are available. Original citation United Nations Children's Fund, Division of Data, Analysis, Planning and Monitoring (2023).

Global UNICEF Global Databases: Infant and Young Child Feeding: Exclusive breastfeeding, New York, October 2023.



Oesophageal cancer

Men, 2022










Breast cancer

Women, 2022





Area covered: National

References:


Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]

Definitions: Incidence per 100,000




Colorectal cancer

Men, 2022










Pancreatic cancer

Men, 2022










Gallbladder cancer

Men, 2022











Kidney cancer

Men, 2022










Cancer of the uterus


Women, 2022





Raised blood pressure

Adults, 2015



References:


Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A875?lang=en

Definitions:

Age Standardised estimated % Raised blood pressure 2015 (SBP>=140 OR DBP>=90).



Men, 2015




References:

Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A875?lang=en

Definitions:

Age Standardised estimated % Raised blood pressure 2015 (SBP>=140 OR DBP>=90).





References:

Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A875?lang=en


Definitions:

Age Standardised estimated % Raised blood pressure 2015 (SBP>=140 OR DBP>=90).



Raised cholesterol

Adults, 2008



References:


Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885

Definitions:

% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).



Men, 2008




References:

 ${\sf Global\ Health\ Observatory\ data\ repository,\ World\ Health\ Organisation,\ } \underline{{\sf http://apps.who.int/gho/data/node.main.} A885}$

Definitions:

% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).





References:

 ${\sf Global\ Health\ Observatory\ data\ repository,\ World\ Health\ Organisation,\ \underline{http://apps.who.int/gho/data/node.main.} \underline{A885}}$


Definitions:

% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).



Raised fasting blood glucose

Men, 2014




References:

Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A869?lang=en

Definitions:

Age Standardised % raised fasting blood glucose (>= 7.0 mmoVL or on medication).





References:

Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A869?lang=en


Definitions:

Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication).



Diabetes prevalence

Adults, 2024




References:

Reproduced with kind permission International Diabetes Federation. IDF Diabetes Atlas, 11th edn. Brussels, Belgium:International Diabetes Federation, 2025. http://www.diabetesatlas.org



Ovarian Cancer

Women, 2022



Age: 20+

References:

Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]

Definitions: Incidence per 100,000



Leukemia

Men, 2022										
4										
3										
2										
1										
0———										
⁻¹ 1	0	1	2	3	4	5	6			
Age:							20+			
Area covered:							National			
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]									
Definitions:						Incidence pe	er 100,000			
	Unless ot	therwise noted, overw	veight refers to a BMI	between 25kg and 29	9.9kg/m², obesity refers	to a BMI greater thar	n 30kg/m².			




Women, 20	022										
4											
3											
2											
1											
0———											
⁻¹ 1	0	1	2	3	4	5	6				
Age:							20+				
Area covered:							National				
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]										
Definitions:						Incidence pe	er 100,000				
	Unless oth	erwise noted, overw	reight refers to a BMI l	petween 25kg and 29	.9kg/m², obesity refers	s to a BMI greater than	n 30kg/m².				



Leukemia

Men, 2022



Area covered: National


References:

Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]

Definitions: Incidence per 100,000

 $Unless \ otherwise \ noted, overweight \ refers \ to \ a \ BMI \ between \ 25kg \ and \ 29.9kg/m^2, obesity \ refers \ to \ a \ BMI \ greater \ than \ 30kg/m^2.$





Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]


Definitions: Incidence per 100,000

 $Unless \ otherwise \ noted, overweight \ refers \ to \ a \ BMI \ between \ 25kg \ and \ 29.9kg/m^2, obesity \ refers \ to \ a \ BMI \ greater \ than \ 30kg/m^2.$




Liver and intrahepatic bile duct Cancer

Men, 2022










Multiple Myeloma

Men, 2022



References:

Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today,


accessed [16.07.24]

Definitions:

Indicence per 100,000

 $Unless \ otherwise \ noted, overweight \ refers \ to \ a \ BMI \ between \ 25kg \ and \ 29.9kg/m^2, obesity \ refers \ to \ a \ BMI \ greater \ than \ 30kg/m^2.$





Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer References:

Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]


Definitions: Indicence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².



Non Hodgkin Lymphoma

Men, 2022



Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today,


accessed [16.07.24]

Definitions:

Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².





Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]


Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².



Thyroid Cancer


Men, 2022



Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]

Definitions: Incidence per 100,000





PDF created on June 17, 2025