




Australia



Country report card - adults


This report card contains the latest data available on the Global Obesity Observatory on overweight and obesity for adults. Where available, data on common and relevant obesity drivers and comorbidities are also presented.

View the latest version of this report on the Global Obesity Observatory at <https://data.worldobesity.org/country/australia-10/>

Contents	Page
Obesity prevalence	3
Trend: % Adults living with obesity in Australia 2007-2022	4
Trend: % Adults living with overweight or obesity in Australia 2007-2022	6
Trend: % Adults living with obesity in selected countries in the Asia/Oceania Region 1975-2019, selected countries	8
Trend: % Adults living with obesity in selected countries worldwide 1976-2018, selected countries	13
Overweight/obesity by education	18
Overweight/obesity by age	19
Overweight/obesity by region	20
Overweight/obesity by socio-economic group	21
Overweight/obesity by ethnicity	22
Double burden of underweight & overweight	24
Insufficient physical activity	25
Estimated per capita fruit intake	28
Estimated per-capita processed meat intake	29
Estimated per capita whole grains intake	30
Mental health - depression disorders	31
Mental health - anxiety disorders	34
Percent of population who cannot afford a healthy diet	37
Oesophageal cancer	38
Breast cancer	40
Colorectal cancer	41
Pancreatic cancer	43
Gallbladder cancer	45
Kidney cancer	47
Cancer of the uterus	49
Raised blood pressure	50
Raised cholesterol	53
Raised fasting blood glucose	56
Diabetes prevalence	58
Ovarian Cancer	59
Leukemia	60
Leukemia	62
Liver and intrahepatic bile duct Cancer	64
Multiple Myeloma	66
Non Hodgkin Lymphoma	68
Thyroid Cancer	70

Obesity prevalence

Adults, 2022-2023



Survey type: Measured

Age: 18+

Sample size: ~12846

Area covered: National


References: Australian National Health Survey 2022-2023. <https://www.abs.gov.au/statistics/health/health-conditions-and-risks/waist-circumference-and-bmi/2022#body-mass-index-bmi-> (Accessed 03.01.2024)

Notes: Provision of height, weight and waist measurements were voluntary. Self-reported health status, height, and weight was collected for all participants. In 2022, 41.8% of adult respondents did not have their height and/or weight measured. For these people, height and weight were imputed using a range of information including their self-reported height and weight

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

% Adults living with obesity in Australia 2007-2022

Men



Survey type:

Measured


References:

- 2007: Australian Bureau of Statistics (ABS). National Health Survey: Summary of results. Canberra, Australia, ABS, 2009. WHO Global InfoBase reference:102910
- 2011: Australian Institute of Health and Welfare 2015. Cardiovascular disease, diabetes and chronic kidney disease—Australian facts: Risk factors. Cardiovascular, diabetes and chronic kidney disease series no. 4. Cat. no. CDK 4. Canberra: AIHW. ABS 2013a. Australian Health Survey: biomedical results for chronic diseases, 2011–12. ABS. cat. no. 4364.0.55.005. Canberra: ABS. Australian Health Survey 2011–12. <http://www.aihw.gov.au/publication-detail/?id=60129550538>
- 2014: Australian National Health Survey, 2014–15 First Results. Australian Bureau of Statistics. <http://www.abs.gov.au/ausstats/abs@.nsf/Latestproducts/4364.0.55.001Appendix22014-15?opendocument&tabname=Notes&prodno=4364.0.55.001&issue=2014-15&num=&view=> (last accessed 27th September 2017)
- 2017: Australian National Health Survey 2017–18 (provisional results). <http://abs.gov.au/ausstats/abs@.nsf/Lookup/by%20Subject/4364.0.55.001~2017-18~Main%20Features~About%20the%20National%20Health%20Survey~5> (accessed 12.12.18)
- 2022: Australian National Health Survey 2022–2023. <https://www.abs.gov.au/statistics/health/health-conditions-and-risks/waist-circumference-and-bmi/2022#body-mass-index-bmi> (Accessed 03.01.2024)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women



Survey type: Measured

References: 2007: Australian Bureau of Statistics (ABS). National Health Survey: Summary of results. Canberra, Australia, ABS, 2009. WHO Global InfoBase reference:102910

2011: Australian Institute of Health and Welfare 2015. Cardiovascular disease, diabetes and chronic kidney disease—Australian facts: Risk factors. Cardiovascular, diabetes and chronic kidney disease series no. 4. Cat. no. CDK 4. Canberra: AIHW. ABS 2013a. Australian Health Survey: biomedical results for chronic diseases, 2011–12. ABS. cat. no. 4364.0.55.005. Canberra: ABS. Australian Health Survey 2011–12. <http://www.aihw.gov.au/publication-detail/?id=60129550538>

2014: Australian National Health Survey, 2014–15 First Results. Australian Bureau of Statistics. <http://www.abs.gov.au/ausstats/abs@.nsf/Latestproducts/4364.0.55.001Appendix22014-15?opendocument&tabname=Notes&prodno=4364.0.55.001&issue=2014-15&num=&view=> (last accessed 27th September 2017)

2017: Australian National Health Survey 2017–18 (provisional results). <http://abs.gov.au/ausstats/abs@.nsf/Lookup/by%20Subject/4364.0.55.001~2017-18~Main%20Features~About%20the%20National%20Health%20Survey~5> (accessed 12.12.18)


2022: Australian National Health Survey 2022–2023. <https://www.abs.gov.au/statistics/health/health-conditions-and-risks/waist-circumference-and-bmi/2022#body-mass-index-bmi-> (Accessed 03.01.2024)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with overweight or obesity in Australia 2007-2022

Men



Survey type:

Measured


References:

- 2007: Australian Bureau of Statistics (ABS). National Health Survey: Summary of results. Canberra, Australia, ABS, 2009. WHO Global InfoBase reference:102910
- 2011: Australian Institute of Health and Welfare 2015. Cardiovascular disease, diabetes and chronic kidney disease—Australian facts: Risk factors. Cardiovascular, diabetes and chronic kidney disease series no. 4. Cat. no. CDK 4. Canberra: AIHW. ABS 2013a. Australian Health Survey: biomedical results for chronic diseases, 2011–12. ABS. cat. no. 4364.0.55.005. Canberra: ABS. Australian Health Survey 2011–12. <http://www.aihw.gov.au/publication-detail/?id=60129550538>
- 2014: Australian National Health Survey, 2014–15 First Results. Australian Bureau of Statistics. <http://www.abs.gov.au/ausstats/abs@.nsf/Latestproducts/4364.0.55.001Appendix22014-15?opendocument&tabname=Notes&prodno=4364.0.55.001&issue=2014-15&num=&view=> (last accessed 27th September 2017)
- 2017: Australian National Health Survey 2017–18 (provisional results). <http://abs.gov.au/ausstats/abs@.nsf/Lookup/by%20Subject/4364.0.55.001~2017-18~Main%20Features~About%20the%20National%20Health%20Survey~5> (accessed 12.12.18)
- 2022: Australian National Health Survey 2022–2023. <https://www.abs.gov.au/statistics/health/health-conditions-and-risks/waist-circumference-and-bmi/2022#body-mass-index-bmi-> (Accessed 03.01.2024)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women



Survey type: Measured

References:

2007: Australian Bureau of Statistics (ABS). National Health Survey: Summary of results. Canberra, Australia, ABS, 2009. WHO Global InfoBase reference:102910

2011: Australian Institute of Health and Welfare 2015. Cardiovascular disease, diabetes and chronic kidney disease—Australian facts: Risk factors. Cardiovascular, diabetes and chronic kidney disease series no. 4. Cat. no. CDK 4. Canberra: AIHW. ABS 2013a. Australian Health Survey: biomedical results for chronic diseases, 2011–12. ABS. cat. no. 4364.0.55.005. Canberra: ABS. Australian Health Survey 2011–12. <http://www.aihw.gov.au/publication-detail/?id=60129550538>

2014: Australian National Health Survey, 2014–15 First Results. Australian Bureau of Statistics. <http://www.abs.gov.au/ausstats/abs@.nsf/Latestproducts/4364.0.55.001Appendix22014-15?opendocument&tabname=Notes&prodno=4364.0.55.001&issue=2014-15&num=&view=> (last accessed 27th September 2017)

2017: Australian National Health Survey 2017–18 (provisional results). <http://abs.gov.au/ausstats/abs@.nsf/Lookup/by%20Subject/4364.0.55.001~2017-18~Main%20Features~About%20the%20National%20Health%20Survey~5> (accessed 12.12.18)


2022: Australian National Health Survey 2022–2023. <https://www.abs.gov.au/statistics/health/health-conditions-and-risks/waist-circumference-and-bmi/2022#body-mass-index-bmi-> (Accessed 03.01.2024)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

**% Adults living with obesity in selected countries in the Asia/Oceania
Region 1975-2019, selected countries**

Men



References:

1976, 1981, 1986: Yoshiike N, Seino F, Tajima S, Arai Y, Kawano M, Furuhashi T, Inoue S. Twenty-year changes in the prevalence of overweight in Japanese adults: The National Nutrition Survey 1976-95. *Obesity Reviews* 2002;3:183-190

1993: Solon FS. Nutrition related chronic diseases in the Philippines. Makati city, Philippines: Nutrition Center of the Philippines Report Series, vol 2, No.1, cited in Reference 53

1995, 1996: Martorell R, Khan LK, Hughes ML, Grummer Strawn LM. Obesity in women from developing countries. *EJCN* (2000) 54:247-252

1998: Colaguir S, Colagiuri R, Na'ati S, Muimuiheata S, Hussain Z and Palau T. (2002). The prevalence of diabetes in the Kingdom of Tonga. *Diabetes care*, 25: 1378 - 1383.

2000: Asia Pacific Cohort Studies Collaboration. The burden of overweight and obesity in the Asia-Pacific region. *Obesity Reviews* 2007;8:191-196.

2001: SCN (2004). 5th Report on the World Nutrition Situation. Nutrition for Improved Development Outcomes. Appendix 11

2002: Report of the 2002 China National Nutrition and Health Survey. 2004. (In Chinese). Chinese Ministry of Public Health (CMPH).

2003: <http://www.fnri.dost.gov.ph/files/fnri%20files/nns/factsandfigures2003/anthropometric.pdf> (last accessed June 14th 2011)

2004: Tonga STEPS Survey 2004

2005, 2013: Chang HC, Yang HC, Chang HY, et al. Morbid obesity in Taiwan: Prevalence, trends, associated social demographics, and lifestyle factors. *PLoS One*. 2017;12(2):e0169577. Published 2017 Feb 2. doi:10.1371/journal.pone.0169577

2006: Ministry of Health and Population - MOHP/Nepal, New ERA/Nepal, and Macro International. 2007. Nepal Demographic and Health Survey 2006. Kathmandu, Nepal: MOHP/Nepal, New ERA/Nepal, and Macro International.

2007: National Institute of Population Research and Training (NIPORT), Mitra and Associates, and Macro International. 2008. Bangladesh Demographic and Health Survey 2007: Key Findings. Calverton, Maryland, USA: NIPORT, Mitra and Associates, and Macro International.

2009: Yan, S., Li, J., Li, S., Zhang, B., Du, S., Gordon-Larsen, P., Adair, L. and Popkin, B. (2012), The expanding burden of cardiometabolic risk in China: the China Health and Nutrition Survey. *Obesity Reviews*. doi: 10.1111/j.1467-789X.2012.01016.x

2011: National Institute of Population Research and Training - NIPORT/Bangladesh, Mitra and Associates/Bangladesh, and ICF International. 2013. Bangladesh Demographic and Health Survey 2011. Dhaka, Bangladesh: NIPORT, Mitra and Associates, and ICF International.

2012: STEPS Survey Tonga 2012 available at https://www.who.int/ncds/surveillance/steps/2012_Tonga_STEPSReport.pdf (last accessed 04.05.2020)


2014: Australian National Health Survey, 2014-15 First Results. Australian Bureau of Statistics. <http://www.abs.gov.au/ausstats/abs@.nsf/Latestproducts/4364.0.55.001Appendix22014-15?opendocument&tabname=Notes&prodno=4364.0.55.001&issue=2014-15&num=&view=> (last accessed 27th September 2017)

2015: Institute for Public Health (IPH) 2015. National Health and Morbidity Survey 2015 (NHMS 2015). Vol. II: Non-Communicable Diseases, Risk Factors & Other Health Problems; 2015.

2016: Ministry of Health - MOH/Nepal, New ERA/Nepal, and ICF. 2017. Nepal Demographic and Health Survey 2016. Kathmandu, Nepal: MOH/Nepal, New ERA/Nepal, and ICF.

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women




References:

- 1976, 1981, 1986: Yoshiike N, Seino F, Tajima S, Arai Y, Kawano M, Furuhashi T, Inoue S. Twenty-year changes in the prevalence of overweight in Japanese adults: The National Nutrition Survey 1976-95. *Obesity Reviews* 2002;3:183-190
- 1993: Solon FS. Nutrition related chronic diseases in the Philippines. Makati city, Philippines: Nutrition Center of the Philippines Report Series, vol 2, No.1, cited in Reference 53
- 1995, 1996: Martorell R, Khan LK, Hughes ML, Grummer Strawn LM. Obesity in women from developing countries. *EJCN* (2000) 54:247-252
- 1998: Colaguir S, Colagiuri R, Na'ati S, Muimuiheata S, Hussain Z and Palau T. (2002). The prevalence of diabetes in the Kingdom of Tonga. *Diabetes care*, 25: 1378 - 1383.
- 2000: Asia Pacific Cohort Studies Collaboration. The burden of overweight and obesity in the Asia-Pacific region. *Obesity Reviews* 2007;8:191-196.
- 2001: SCN (2004). 5th Report on the World Nutrition Situation. Nutrition for Improved Development Outcomes. Appendix 11
- 2002: Report of the 2002 China National Nutrition and Health Survey. 2004. (In Chinese). Chinese Ministry of Public Health (CMPH).
- 2003: <http://www.fnri.dost.gov.ph/files/fnri%20files/nns/factsandfigures2003/anthropometric.pdf> (last accessed June 14th 2011)
- 2004: Tonga STEPS Survey 2004
- 2005, 2013: Chang HC, Yang HC, Chang HY, et al. Morbid obesity in Taiwan: Prevalence, trends, associated social demographics, and lifestyle factors. *PLoS One*. 2017;12(2):e0169577. Published 2017 Feb 2. doi:10.1371/journal.pone.0169577
- 2006: Ministry of Health and Population - MOHP/Nepal, New ERA/Nepal, and Macro International. 2007. Nepal Demographic and Health Survey 2006. Kathmandu, Nepal: MOHP/Nepal, New ERA/Nepal, and Macro International.
- 2007: National Institute of Population Research and Training (NIPORT), Mitra and Associates, and Macro International. 2008. Bangladesh Demographic and Health Survey 2007: Key Findings. Calverton, Maryland, USA: NIPORT, Mitra and Associates, and Macro International.
- 2009: Yan, S., Li, J., Li, S., Zhang, B., Du, S., Gordon-Larsen, P., Adair, L. and Popkin, B. (2012), The expanding burden of cardiometabolic risk in China: the China Health and Nutrition Survey. *Obesity Reviews*. doi: 10.1111/j.1467-789X.2012.01016.x
- 2011: National Institute of Population Research and Training - NIPORT/Bangladesh, Mitra and Associates/Bangladesh, and ICF International. 2013. Bangladesh Demographic and Health Survey 2011. Dhaka, Bangladesh: NIPORT, Mitra and Associates, and ICF International.
- 2012: STEPS Survey Tonga 2012 available at https://www.who.int/ncds/surveillance/steps/2012_Tonga_STEPSReport.pdf (last accessed 04.05.2020)
- 2014: Australian National Health Survey, 2014-15 First Results. Australian Bureau of Statistics. <http://www.abs.gov.au/ausstats/abs@.nsf/Latestproducts/4364.0.55.001Appendix22014-15?opendocument&tabname=Notes&prodno=4364.0.55.001&issue=2014-15&num=&view=> (last accessed 27th September 2017)
- 2015: Institute for Public Health (IPH) 2015. National Health and Morbidity Survey 2015 (NHMS 2015). Vol. II: Non-Communicable Diseases, Risk Factors & Other Health Problems; 2015.
- 2016: Ministry of Health - MOH/Nepal, New ERA/Nepal, and ICF. 2017. Nepal Demographic and Health Survey 2016. Kathmandu, Nepal: MOH/Nepal, New ERA/Nepal, and ICF.

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

**% Adults living with obesity in selected countries worldwide 1976-2018,
selected countries**


Men



- References:**
- 1960, 1971, 1973, 1976, 1988, 1991: Flegal KM, Carroll MD, Kuczmarski RJ, Johnson CL. Overweight and obesity in the United States: prevalence and trends, 1960-1994. *International Journal of Obesity* (1998);22:39-47
 - 1975: Monteiro CA, Conde WL, Popking BM. Is obesity replacing or adding to undernutrition? Evidence from different social classes in Brazil. 2002. *Public Health Nutrition*:51(1A), 105-112
 - 1981, 1986: Yoshiike N, Seino F, Tajima S, Arai Y, Kawano M, Furuhashi T, Inoue S. Twenty-year changes in the prevalence of overweight in Japanese adults: The National Nutrition Survey 1976-95. *Obesity Reviews* 2002;3:183-190
 - 1982, 1993: Rodriguez-Ojea A, Jimenez S, Berdasco A, Esquivel M. The nutrition transition in Cuba in the nineties:an overview. *Public health Nutrition* 2002;5(1A), 129-133
 - 1985: Berg C, Rosengren A, Aires N, appas G, Toren K, Thelle D, Lissner L. Trends in overweight and obesity from 1985 to 2002 in Goteborg, West Sweden. *IJO* 2005 Aug;29(8):916-24
 - 1990: Berg C, Rosengren A, Aires N, appas G, Toren K, Thelle D, Lissner L. Trends in overweight and obesity from 1985 to 2002 in Goteborg, West Sweden. *IJO* 2005 online published ahead of print.
 - 1992: Uauy R, Albal C, Kain J. Obesity Trends in Latin America: Transiting from Under-to Overweight. *Journal of Nutrition* 2001;131:S893-S899
 - 1995: Health Survey for England 1995.
 - 1996: Health Survey for England 1996.
 - 1997: Filozof C, Gonzales C, Sereday M, Mazza C, Braguinsky J. Obesity prevalence and trends in Latin American countries. *Obesity Reviews*, 2001;2:99-196
 - 1998: Scottish Health Survey 1998
 - 1999: Health Survey for England 1999.
 - 2000: Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of Overweight and Obesity in the United States, 1999-2004. *JAMA* 2006;295(13):1549-1555
 - 2001: Health Survey for England 2001.
 - 2002: Monteiro CA, Conde WL and Popkin BA. (2007). Income-specific trends in obesity in Brazil: 1975 - 2003. *American Journal of Public Health*, 97 (10): 1808 - 1812.
 - 2003: 2003 ENS Report. Final results on the National Health Survey. <http://epi.minsal.cl/epi/html/invest/ENS/informeFinalENS.pdf>.
 - 2004: Health Survey for England 2004.
 - 2005: Health Survey for England 2005.
 - 2006: Health Survey for England 2006.
 - 2007: Peltonen M, Harald K, Männistö S, Saarikoski L, Lund L, Sundvall J, Juolevi A, Laatikainen T, Aldén-Nieminen H, Luoto R, Jousilahti P, Salomaa V, Taimi M, Vartiainen E. Kansallinen FINRISKI 2007 -terveyystutkimus, Tutkimuksen toteutus ja tulokset: Taulukkoliite. Kansanterveyslaitos. Yliopistopaino, Helsinki 2008.
 - 2008: Health Survey for England 2008.
 - 2009: NHANES Survey - Published in Flegal KM, Carroll MD, Kit BK, Ogden CL. Prevalence of Obesity and Trends in the Distribution of

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women




References:

- 1960, 1971, 1973, 1976, 1988, 1991: Flegal KM, Carroll MD, Kuczmarski RJ, Johnson CL. Overweight and obesity in the United States: prevalence and trends, 1960-1994. *International Journal of Obesity* (1998);22:39-47
- 1975: Monteiro CA, Conde WL, Popking BM. Is obesity replacing or adding to undernutrition? Evidence from different social classes in Brazil. 2002. *Public Health Nutrition*:51(1A), 105-112
- 1981, 1986: Yoshiike N, Seino F, Tajima S, Arai Y, Kawano M, Furuhashi T, Inoue S. Twenty-year changes in the prevalence of overweight in Japanese adults: The National Nutrition Survey 1976-95. *Obesity Reviews* 2002;3:183-190
- 1982, 1993: Rodriguez-Ojeda A, Jimenez S, Berdasco A, Esquivel M. The nutrition transition in Cuba in the nineties: an overview. *Public health Nutrition* 2002;5(1A), 129-133
- 1985: Berg C, Rosengren A, Aires N, Appas G, Toren K, Thelle D, Lissner L. Trends in overweight and obesity from 1985 to 2002 in Goteborg, West Sweden. *IJO* 2005 Aug;29(8):916-24
- 1990: Berg C, Rosengren A, Aires N, Appas G, Toren K, Thelle D, Lissner L. Trends in overweight and obesity from 1985 to 2002 in Goteborg, West Sweden. *IJO* 2005 online published ahead of print.
- 1992: Uauy R, Albal C, Kain J. Obesity Trends in Latin America: Transiting from Under- to Overweight. *Journal of Nutrition* 2001;131:S893-S899
- 1995: Health Survey for England 1995.
- 1996: Health Survey for England 1996.
- 1997: Filozof C, Gonzales C, Sereday M, Mazza C, Braguinsky J. Obesity prevalence and trends in Latin American countries. *Obesity Reviews*, 2001;2:99-196
- 1998: Scottish Health Survey 1998
- 1999: Health Survey for England 1999.
- 2000: Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of Overweight and Obesity in the United States, 1999-2004. *JAMA* 2006;295(13):1549-1555
- 2001: Health Survey for England 2001.
- 2002: Monteiro CA, Conde WL and Popkin BA. (2007). Income-specific trends in obesity in Brazil: 1975 - 2003. *American Journal of Public Health*, 97 (10): 1808 - 1812.
- 2003: 2003 ENS Report. Final results on the National Health Survey. <http://epi.minsal.cl/epi/html/invest/ENS/informeFinalENS.pdf>.
- 2004: Health Survey for England 2004.
- 2005: Health Survey for England 2005.
- 2006: Health Survey for England 2006.
- 2007: Peltonen M, Harald K, Mannistö S, Saarikoski L, Lund L, Sundvall J, Juolevi A, Laatikainen T, Aldén-Nieminen H, Luoto R, Jousilahti P, Salomaa V, Taimi M, Vartiainen E. Kansallinen FINRISKI 2007 -terveys tutkimus, Tutkimuksen toteutus ja tulokset: Taulukkoliite. Kansanterveyslaitos. Yliopistopaino, Helsinki 2008.
- 2008: Health Survey for England 2008.
- 2009: NHANES Survey - Published in Flegal KM, Carroll MD, Kit BK, Ogden CL. Prevalence of Obesity and Trends in the Distribution of

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Overweight/obesity by education

Adults, 2022-2023



Survey type: Measured

Age: 18+

Sample size: ~12846

Area covered: National


References: Australian National Health Survey 2022-2023. <https://www.abs.gov.au/statistics/health/health-conditions-and-risks/waist-circumference-and-bmi/2022#body-mass-index-bmi-> (Accessed 03.01.2024)

Notes: Provision of height, weight and waist measurements were voluntary. Self-reported health status, height, and weight was collected for all participants. In 2022, 41.8% of adult respondents did not have their height and/or weight measured. For these people, height and weight were imputed using a range of information including their self-reported height and weight

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Overweight/obesity by age

Adults, 2022-2023



Survey type: Measured

Sample size: ~12846

Area covered: National


References: Australian National Health Survey 2022-2023. <https://www.abs.gov.au/statistics/health/health-conditions-and-risks/waist-circumference-and-bmi/2022#body-mass-index-bmi-> (Accessed 03.01.2024)

Notes: Provision of height, weight and waist measurements were voluntary. Self-reported health status, height, and weight was collected for all participants. In 2022, 41.8% of adult respondents did not have their height and/or weight measured. For these people, height and weight were imputed using a range of information including their self-reported height and weight

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Overweight/obesity by region

Adults, 2022-2023



Survey type: Measured

Age: 18+

Sample size: ~12846

Area covered: National


References: Australian National Health Survey 2022-2023. <https://www.abs.gov.au/statistics/health/health-conditions-and-risks/waist-circumference-and-bmi/2022#body-mass-index-bmi-> (Accessed 03.01.2024)

Notes: Provision of height, weight and waist measurements were voluntary. Self-reported health status, height, and weight was collected for all participants. In 2022, 41.8% of adult respondents did not have their height and/or weight measured. For these people, height and weight were imputed using a range of information including their self-reported height and weight

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Overweight/obesity by socio-economic group

Adults, 2022-2023



Survey type: Measured

Age: 18+

Sample size: ~12846

Area covered: National

References: Australian National Health Survey 2022-2023. <https://www.abs.gov.au/statistics/health/health-conditions-and-risks/waist-circumference-and-bmi/2022#body-mass-index-bmi-> (Accessed 03.01.2024)

Notes: Provision of height, weight and waist measurements were voluntary. Self-reported health status, height, and weight was collected for all participants. In 2022, 41.8% of adult respondents did not have their height and/or weight measured. For these people, height and weight were imputed using a range of information including their self-reported height and weight


Definitions: A lower Index of Disadvantage quintile (e.g. the first quintile) indicates relatively greater disadvantage and a lack of advantage in general. A higher Index of Disadvantage (e.g. the fifth quintile) indicates a relative lack of disadvantage and greater advantage in general.

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Overweight/obesity by ethnicity

Ethnic groups are as defined by publication of origin and are not as defined by WOF. In some instances ethnicity is conflated with nationality and/or race.


Men, 2011



Survey type:	Self-reported
Age:	18+
Sample size:	16,044
References:	Menigoz, Karen, et al. "Ethnic Differences in Overweight and Obesity and the Influence of Acculturation on Immigrant Bodyweight: Evidence from a National Sample of Australian Adults." BMC Public Health, vol. 16, no. 1, 5 Sept. 2016, www.ncbi.nlm.nih.gov/pmc/articles/PMC5011908/ , 10.1186/s12889-016-3608-6. Accessed 30 Sept. 2021.
Definitions:	Country of birth

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2011



Survey type: Self-reported

Age: 18+

Sample size: 16,044


References: Menigoz, Karen, et al. "Ethnic Differences in Overweight and Obesity and the Influence of Acculturation on Immigrant Bodyweight: Evidence from a National Sample of Australian Adults." BMC Public Health, vol. 16, no. 1, 5 Sept. 2016, www.ncbi.nlm.nih.gov/pmc/articles/PMC5011908/, 10.1186/s12889-016-3608-6. Accessed 30 Sept. 2021.

Definitions: Country of birth

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Double burden of underweight & overweight


Adults, 2022



Survey type:	Measured
Age:	20+
References:	NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population representative studies with 222 million children, adolescents, and adults. Lancet 2024; published online Feb 29. https://doi.org/10.1016/S0140-6736(23)02750-2 .
Notes:	Age Standardised estimates
Definitions:	Combined prevalence of BMI<18.5 kg/m ² and BMI≥30 kg/m ² (double burden of underweight and obesity)

Insufficient physical activity

Adults, 2022



Survey type: Self-reported


Age: 18+

Area covered: National

References: WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-\(age-standardized-estimate\)-\(-\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-))

Definitions: Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.

Men, 2022



Survey type: Self-reported


Age: 18+

Area covered: National

References: WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-\(age-standardized-estimate\)-\(-\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-))

Definitions: Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.

Women, 2022



Survey type: Self-reported

Age: 18+


Area covered: National

References: WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-\(age-standardized-estimate\)-\(-\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-))

Definitions: Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.

Estimated per capita fruit intake

Adults, 2017



Survey type: Measured


Age: 25+

References: Global Burden of Disease, the Institute for Health Metrics and Evaluation <http://ghdx.healthdata.org/>

Definitions: Estimated per-capita fruit intake (g/day)

Estimated per-capita processed meat intake

Adults, 2017



Survey type: Measured


Age: 25+

References: Global Burden of Disease, the Institute for Health Metrics and Evaluation <http://ghdx.healthdata.org/>

Definitions: Estimated per-capita processed meat intake (g per day)

Estimated per capita whole grains intake

Adults, 2017



Survey type: Measured


Age: 25+

References: Global Burden of Disease, the Institute for Health Metrics and Evaluation <http://ghdx.healthdata.org/>

Definitions: Estimated per-capita whole grains intake (g/day)

Mental health - depression disorders

Adults, 2021




Age: 20+

Area covered: National

References: Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from <http://vizhub.healthdata.org/gbd-compare>. (Last accessed 23.04.25)

Definitions: Number living with depression per 100,000 population (adults 20+ years)

Men, 2021



Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from http://vizhub.healthdata.org/gbd-compare . (Last accessed 23.04.25)
Definitions:	Number living with depression per 100,000 population (adults 20+ years)

Women, 2021



Age: 20+


Area covered: National

References: Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from <http://vizhub.healthdata.org/gbd-compare>. (Last accessed 23.04.25)

Definitions: Number living with depression per 100,000 population (adults 20+ years)

Mental health - anxiety disorders

Adults, 2021




Age: 20+

References: Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from <http://vizhub.healthdata.org/gbd-compare>. (Last accessed 23.04.25).

Definitions: Number living with anxiety per 100,000 population

Men, 2021




Age: 20+

References: Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from <http://vizhub.healthdata.org/gbd-compare>. (Last accessed 23.04.25).

Definitions: Number living with anxiety per 100,000 population

Women, 2021




Age: 20+

References: Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from <http://vizhub.healthdata.org/gbd-compare>. (Last accessed 23.04.25).

Definitions: Number living with anxiety per 100,000 population

Percent of population who cannot afford a healthy diet

Adults, 2022



Area covered:


National

References:

The Food Systems Dashboard. The Global Alliance for Improved Nutrition (GAIN), The Columbia Climate School, and Cornell University College of Agriculture and Life Sciences. 2024. Geneva, Switzerland. <https://www.foodsystemsdashboard.org>. DOI: <https://doi.org/10.36072/db>.


Oesophageal cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Indicence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Indicence per 100,000

Breast cancer

Women, 2022




Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Colorectal cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Incidence per 100,000


Pancreatic cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Incidence per 100,000

Gallbladder cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Indicence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Indicence per 100,000

Kidney cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Incidence per 100,000

Cancer of the uterus

Women, 2022




Age: 20+

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Raised blood pressure


Adults, 2015



References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A875?lang=en>

Definitions: Age Standardised estimated % Raised blood pressure 2015 (SBP \geq 140 OR DBP \geq 90).


Men, 2015



References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A875?lang=en>

Definitions: Age Standardised estimated % Raised blood pressure 2015 (SBP \geq 140 OR DBP \geq 90).

Women, 2015




References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A875?lang=en>

Definitions: Age Standardised estimated % Raised blood pressure 2015 (SBP \geq 140 OR DBP \geq 90).

Raised cholesterol


Adults, 2008



References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A885>

Definitions: % Raised total cholesterol (≥ 5.0 mmol/L) (age-standardized estimate).


Men, 2008



References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A885>

Definitions: % Raised total cholesterol (≥ 5.0 mmol/L) (age-standardized estimate).

Women, 2008




References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A885>

Definitions: % Raised total cholesterol (≥ 5.0 mmol/L) (age-standardized estimate).

Raised fasting blood glucose


Men, 2014



References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A869?lang=en>

Definitions: Age Standardised % raised fasting blood glucose (≥ 7.0 mmol/L or on medication).

Women, 2014




References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A869?lang=en>

Definitions: Age Standardised % raised fasting blood glucose (≥ 7.0 mmol/L or on medication).

Diabetes prevalence

Adults, 2024




References:

Reproduced with kind permission International Diabetes Federation. IDF Diabetes Atlas, 11th edn. Brussels, Belgium:International Diabetes Federation, 2025. <http://www.diabetesatlas.org>

Ovarian Cancer

Women, 2022




Age: 20+

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Leukemia

Men, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022



Age: 20+

Area covered: National


References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Leukemia

Men, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022



Age: 20+

Area covered: National


References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Liver and intrahepatic bile duct Cancer

Men, 2022




Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Women, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Multiple Myeloma

Men, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022



Age: 20+

Area covered: National


References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Non Hodgkin Lymphoma

Men, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022



Age: 20+

Area covered: National


References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Thyroid Cancer

Men, 2022




Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Women, 2022



Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

PDF created on June 17, 2025