

Report card


Brazil



Contents	Page
Obesity prevalence	2
Trend: % Adults living with obesity, 1975-2019	5
Trend: % Adults living with overweight or obesity, 1975-2019	8
Trend: % Adults living with obesity, 2006-2019	11
Trend: % Adults living with overweight or obesity, 2006-2019	14
Trend: % Adults living with obesity, selected countries, 1960-2022	17
Trend: % Adults living with obesity, selected countries, 1960-2023	22
Overweight/obesity by education	27
Overweight/obesity by age	32
Overweight/obesity by region	34
Overweight/obesity by socio-economic group	39
Insufficient physical activity	42
Estimated per capita fruit intake	48
Estimated per-capita processed meat intake	49
Estimated per capita whole grains intake	50
Mental health - depression disorders	51
Mental health - anxiety disorders	52
% Infants exclusively breastfed 0-5 months	53
Oesophageal cancer	54
Breast cancer	56
Colorectal cancer	57
Pancreatic cancer	59
Gallbladder cancer	61
Kidney cancer	63
Cancer of the uterus	65
Raised blood pressure	66
Raised cholesterol	69
Raised fasting blood glucose	72
Diabetes prevalence	74
Contextual factors	75

Obesity prevalence

Adults, 2019




Survey type:	Self-reported
Age:	18+
Sample size:	52443
Area covered:	National

References: Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil

Notes: Data from Capitals of 26 Brazilian States and the Federal District

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Children, 2019



Survey type:	Measured
Age:	15-17
Area covered:	National
References:	Pesquisa nacional de saúde : 2019 : ciclos de vida : Brasil / IBGE, Coordenação de Trabalho e Rendimento. - Rio de Janeiro : IBGE, 2021. Available at https://www.ibge.gov.br/en/statistics/social/health/16840-national-survey-of-health.html?=&t=resultados (last accessed 04.04.23)
Cutoffs:	WHO

Infants, 2019


█ Overweight or obesity



Age:	0-5
Sample size:	10780287
References:	NNS: Estudo Nacional de Alimentação e Nutrição Infantil (ENANI), Brasil 2019
Notes:	UNICEF/WHO/World Bank Joint Child Malnutrition Estimates Expanded Database: Overweight (Survey Estimates), May 2023, New York. For more information about the methodology, please consult https://data.unicef.org/resources/jme-2023-country-consultations/ Percentage of children under 5 years of age falling above 2 standard deviations (moderate and severe) from the median weight-for-height of the reference population.
Definitions:	=>+2SD

% Adults living with obesity, 1975-2019

Women



Survey type: Measured

References:


- 1975: Monteiro CA, Conde WL, Popkin BM. Is obesity replacing or adding to undernutrition? Evidence from different social classes in Brazil. 2002. *Public Health Nutrition*:51(1A), 105-112
- 1997: Filozof C, Gonzales C, Sereday M, Mazza C, Braguinsky J. Obesity prevalence and trends in Latin American countries. *Obesity Reviews*, 2001;2:99-196
- 2002: Monteiro CA, Conde WL and Popkin BA. (2007). Income-specific trends in obesity in Brazil: 1975 - 2003. *American Journal of Public Health*, 97 (10): 1808 - 1812.
- 2008: Ministério da Saúde. Pesquisa de Orcamentos Familiares 2008-9. Antropometria E Esado Nutricional de Crianças, Adolescentes E Adultos No Brasil. Instituto Brasileiro de Geografia e Estatística (IBGE) 2010.
- 2013: FERREIRA, Arthur Pate de Souza; SZWARCWALD, Célia Landmann and DAMACENA, Giseli Nogueira. Prevalence of obesity and associated factors in the Brazilian population: a study of data from the 2013 National Health Survey. *Rev. bras. epidemiol.* [online]. 2019, vol.22 [cited 2019-05-16], e190024. Available from: http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1415-790X2019000100420&lng=en&nrm=iso. Epub Apr 01, 2019. ISSN 1415-790X. <http://dx.doi.org/10.1590/1980-549720190024>.
- 2019: Pesquisa nacional de saúde : 2019 : ciclos de vida : Brasil / IBGE, Coordenação de Trabalho e Rendimento. - Rio de Janeiro : IBGE, 2021. Available at <https://www.ibge.gov.br/en/statistics/social/health/16840-national-survey-of-health.html?=&t=resultados> (last accessed 04.04.23)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Men

● Obesity



Survey type:

Measured

References:


- 1975: Monteiro CA, Conde WL, Popkin BM. Is obesity replacing or adding to undernutrition? Evidence from different social classes in Brazil. 2002. Public Health Nutrition:51(1A), 105-112
- 1997: Filozof C, Gonzales C, Sereday M, Mazza C, Braguinsky J. Obesity prevalence and trends in Latin American countries. Obesity Reviews, 2001;2:99-196
- 2002: Monteiro CA, Conde WL and Popkin BA. (2007). Income-specific trends in obesity in Brazil: 1975 - 2003. American Journal of Public Health, 97 (10): 1808 - 1812.
- 2008: Ministério da Saúde. Pesquisa de Orcamentos Familiares 2008-9. Antropometria E Esado Nutricional de Crianças, Adolescentes E Adultos No Brasil. Instituto Brasileiro de Geografi a e Estatística (IBGE) 2010.
- 2013: FERREIRA, Arthur Pate de Souza; SZWARCWALD, Célia Landmann and DAMACENA, Giseli Nogueira. Prevalence of obesity and associated factors in the Brazilian population: a study of data from the 2013 National Health Survey. Rev. bras. epidemiol. [online]. 2019, vol.22 [cited 2019-05-16], e190024. Available from: http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1415-790X2019000100420&lng=en&nrm=iso. Epub Apr 01, 2019. ISSN 1415-790X. <http://dx.doi.org/10.1590/1980-549720190024>.
- 2019: Pesquisa nacional de saúde : 2019 : ciclos de vida : Brasil / IBGE, Coordenação de Trabalho e Rendimento. - Rio de Janeiro : IBGE, 2021. Available at <https://www.ibge.gov.br/en/statistics/social/health/16840-national-survey-of-health.html?=&t=resultados> (last accessed 04.04.23)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with overweight or obesity, 1975-2019

Women



Survey type:

Measured


References:

- 1975: Monteiro CA, Conde WL, Popkin BM. Is obesity replacing or adding to undernutrition? Evidence from different social classes in Brazil. 2002. Public Health Nutrition:51(1A), 105-112
- 1997: Filozof C, Gonzales C, Sereday M, Mazza C, Braguinsky J. Obesity prevalence and trends in Latin American countries. Obesity Reviews, 2001;2:99-196
- 2002: Monteiro CA, Conde WL and Popkin BA. (2007). Income-specific trends in obesity in Brazil: 1975 - 2003. American Journal of Public Health, 97 (10): 1808 - 1812.
- 2008: Ministério da Saúde. Pesquisa de Orcamentos Familiares 2008-9. Antropometria E Esado Nutricional de Crianças, Adolescentes E Adultos No Brasil. Instituto Brasileiro de Geografia e Estatística (IBGE) 2010.
- 2013: FERREIRA, Arthur Pate de Souza; SZWARCWALD, Célia Landmann and DAMACENA, Giseli Nogueira. Prevalence of obesity and associated factors in the Brazilian population: a study of data from the 2013 National Health Survey. Rev. bras. epidemiol. [online]. 2019, vol.22 [cited 2019-05-16], e190024. Available from: http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1415-790X2019000100420&lng=en&nrm=iso. Epub Apr 01, 2019. ISSN 1415-790X. <http://dx.doi.org/10.1590/1980-549720190024>.
- 2019: Pesquisa nacional de saúde : 2019 : ciclos de vida : Brasil / IBGE, Coordenação de Trabalho e Rendimento. - Rio de Janeiro : IBGE, 2021. Available at <https://www.ibge.gov.br/en/statistics/social/health/16840-national-survey-of-health.html?=&t=resultados> (last accessed 04.04.23)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Men



Survey type:

Measured

References:

- 1975: Monteiro CA, Conde WL, Popkin BM. Is obesity replacing or adding to undernutrition? Evidence from different social classes in Brazil. 2002. Public Health Nutrition:51(1A), 105-112
- 1997: Filozof C, Gonzales C, Sereday M, Mazza C, Braguinsky J. Obesity prevalence and trends in Latin American countries. Obesity Reviews, 2001;2:99-196
- 2002: Monteiro CA, Conde WL and Popkin BA. (2007). Income-specific trends in obesity in Brazil: 1975 - 2003. American Journal of Public Health, 97 (10): 1808 - 1812.
- 2008: Ministério da Saúde. Pesquisa de Orcamentos Familiares 2008-9. Antropometria E Esado Nutricional de Crianças, Adolescentes E Adultos No Brasil. Instituto Brasileiro de Geografi a e Estatística (IBGE) 2010.
- 2013: FERREIRA, Arthur Pate de Souza; SZWARCWALD, Célia Landmann and DAMACENA, Giseli Nogueira. Prevalence of obesity and associated factors in the Brazilian population: a study of data from the 2013 National Health Survey. Rev. bras. epidemiol. [online]. 2019, vol.22 [cited 2019-05-16], e190024. Available from: http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1415-790X2019000100420&lng=en&nrm=iso. Epub Apr 01, 2019. ISSN 1415-790X. <http://dx.doi.org/10.1590/1980-549720190024>.
- 2019: Pesquisa nacional de saúde : 2019 : ciclos de vida : Brasil / IBGE, Coordenação de Trabalho e Rendimento. - Rio de Janeiro : IBGE, 2021. Available at <https://www.ibge.gov.br/en/statistics/social/health/16840-national-survey-of-health.html?=&t=resultados> (last accessed 04.04.23)


Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with obesity, 2006-2019

Women

● Obesity




Survey type: Self-reported

References: 2006: BRASIL. Ministério da Saúde. Vigitel Brasil 2006: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília: Ministério da Saúde, 2007. Data published in Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil
 2009: BRASIL. Ministério da Saúde. Vigitel Brasil 2009: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília: Ministério da Saúde, 2010.
 2012: BRASIL. Ministério da Saúde. Vigitel Brasil 2012: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília, DF: Ministério da Saúde, 2013.
 2015: BRASIL. Ministério da Saúde. Vigitel Brasil 2015: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília, DF: Ministério da Saúde, 2016.
 2019: Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Men and women

● Obesity



Survey
type:


Self-reported

References: 2006: BRASIL. Ministério da Saúde. Vigitel Brasil 2006: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília: Ministério da Saúde, 2007. Data published in Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil 2009: BRASIL. Ministério da Saúde. Vigitel Brasil 2009: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília: Ministério da Saúde, 2010. 2012: BRASIL. Ministério da Saúde. Vigitel Brasil 2012: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília, DF: Ministério da Saúde, 2013. 2015: BRASIL. Ministério da Saúde. Vigitel Brasil 2015: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília, DF: Ministério da Saúde, 2016. 2019: Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Men

● Obesity



Survey type:

Self-reported


References: 2006: BRASIL. Ministério da Saúde. Vigitel Brasil 2006: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília: Ministério da Saúde, 2007. Data published in Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil 2009: BRASIL. Ministério da Saúde. Vigitel Brasil 2009: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília: Ministério da Saúde, 2010. 2012: BRASIL. Ministério da Saúde. Vigitel Brasil 2012: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília, DF: Ministério da Saúde, 2013. 2015: BRASIL. Ministério da Saúde. Vigitel Brasil 2015: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília, DF: Ministério da Saúde, 2016. 2019: Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with overweight or obesity, 2006-2019

Women

● Overweight or obesity




Survey type: Self-reported

References: 2006: BRASIL. Ministério da Saúde. Vigitel Brasil 2006: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília: Ministério da Saúde, 2007. Data published in Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil
 2009: BRASIL. Ministério da Saúde. Vigitel Brasil 2009: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília: Ministério da Saúde, 2010.
 2012: BRASIL. Ministério da Saúde. Vigitel Brasil 2012: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília, DF: Ministério da Saúde, 2013.
 2015: BRASIL. Ministério da Saúde. Vigitel Brasil 2015: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília, DF: Ministério da Saúde, 2016.
 2019: Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Men and women

● Overweight or obesity



Survey
type:


Self-reported

References: 2006: BRASIL. Ministério da Saúde. Vigitel Brasil 2006: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília: Ministério da Saúde, 2007. Data published in Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil 2009: BRASIL. Ministério da Saúde. Vigitel Brasil 2009: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília: Ministério da Saúde, 2010. 2012: BRASIL. Ministério da Saúde. Vigitel Brasil 2012: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília, DF: Ministério da Saúde, 2013. 2015: BRASIL. Ministério da Saúde. Vigitel Brasil 2015: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília, DF: Ministério da Saúde, 2016. 2019: Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Men

● Overweight or obesity



Survey type:


Self-reported

References: 2006: BRASIL. Ministério da Saúde. Vigitel Brasil 2006: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília: Ministério da Saúde, 2007. Data published in Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil 2009: BRASIL. Ministério da Saúde. Vigitel Brasil 2009: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília: Ministério da Saúde, 2010. 2012: BRASIL. Ministério da Saúde. Vigitel Brasil 2012: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília, DF: Ministério da Saúde, 2013. 2015: BRASIL. Ministério da Saúde. Vigitel Brasil 2015: vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico. Brasília, DF: Ministério da Saúde, 2016. 2019: Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with obesity, selected countries, 1960-2022


Men



- References:
- 1960, 1971, 1973, 1976, 1988, 1991: Flegal KM, Carroll MD, Kuczmarski RJ, Johnson CL. Overweight and obesity in the United States: prevalence and trends, 1960-1994. *International Journal of Obesity* (1998);22:39-47
 - 1975: Monteiro CA, Conde WL, Popkin BM. Is obesity replacing or adding to undernutrition? Evidence from different social classes in Brazil. 2002. *Public Health Nutrition*:51(1A), 105-112
 - 1992: Arroyo et al. Prevalence of Pre-Obesity and Obesity in Urban Adult Mexicans in Comparison with other Large Surveys. *Obesity Research*. 2000;8:179-185
 - 1994, 1995: Martorell R, Khan LK, Hughes ML, Grummer Strawn LM. Obesity in women from developing countries. *EJCN* (2000) 54;247-252
 - 1997: Filozof C, Gonzales C, Sereday M, Mazza C, Braguinsky J. Obesity prevalence and trends in Latin American countries. *Obesity Reviews*, 2001;2:99-196
 - 1998: Instituto Nacional de Estadística - INE/Guatemala and Macro International. 1999. Guatemala Encuesta Nacional de Salud Materno Infantil 1998-1999. Calverton, Maryland, USA: Instituto Nacional de Estadística - INE/Guatemala and Macro International.
 - 1999: Centres for Disease Control and Prevention. <http://www.cdc.gov/>
 - 2000: SCN (2004). 5th Report on the World Nutrition Situation. Nutrition for Improved Development Outcomes. Appendix 11
 - 2001: N Brathwaite, A Brathwaite, M Taylor. The Socio-economic Determinants of Obesity in Adults in the Bahamas. *West Indian Med J* 2011; 60 (4): 434
 - 2002: Monteiro CA, Conde WL and Popkin BA. (2007). Income-specific trends in obesity in Brazil: 1975 - 2003. *American Journal of Public Health*, 97 (10): 1808 - 1812.
 - 2003: Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, & Flegal KM. (2006). Prevalence of Overweight and Obesity in the United States, 1999-2004. *The Journal of the American Medical Association*, Vol 295(13):1549 - 1555.
 - 2005: 1st Argentinian National Survey of Risk Factors (Encuesta Nacional de Factores de Riesgo). Results from 1-3rd survey reported in the 4th survey report: https://www.indec.gob.ar/ftp/cuadros/publicaciones/enfr_2018_resultados_definitivos.pdf
 - 2006: Olaiz-Fernández G, Rivera-Dommarco J, Shamah-Levy T, Rojas R, Villalpando-Hernández S, Hernández-Avila M, Sepúlveda-Amor J. Encuesta Nacional de Salud y Nutrición 2006. Cuernavaca, México: Instituto Nacional de Salud Pública, 2006. (National Health and Nutrition Survey 2006)

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women




- References:
- 1960, 1971, 1973, 1976, 1988, 1991: Flegal KM, Carroll MD, Kuczmarski RJ, Johnson CL. Overweight and obesity in the United States: prevalence and trends, 1960-1994. *International Journal of Obesity* (1998);22:39-47
 - 1975: Monteiro CA, Conde WL, Popkin BM. Is obesity replacing or adding to undernutrition? Evidence from different social classes in Brazil. 2002. *Public Health Nutrition*:51(1A), 105-112
 - 1992: Arroyo et al. Prevalence of Pre-Obesity and Obesity in Urban Adult Mexicans in Comparison with other Large Surveys. *Obesity Research*. 2000;8:179-185
 - 1994, 1995: Martorell R, Khan LK, Hughes ML, Grummer Strawn LM. Obesity in women from developing countries. *EJCN* (2000) 54;247-252
 - 1997: Filozof C, Gonzales C, Sereday M, Mazza C, Braguinsky J. Obesity prevalence and trends in Latin American countries. *Obesity Reviews*, 2001;2:99-196
 - 1998: Instituto Nacional de Estadística - INE/Guatemala and Macro International. 1999. *Guatemala Encuesta Nacional de Salud Materno Infantil 1998-1999*. Calverton, Maryland, USA: Instituto Nacional de Estadística - INE/Guatemala and Macro International.
 - 1999: Centres for Disease Control and Prevention. <http://www.cdc.gov/>
 - 2000: SCN (2004). 5th Report on the World Nutrition Situation. Nutrition for Improved Development Outcomes. Appendix 11
 - 2001: N Brathwaite, A Brathwaite, M Taylor. The Socio-economic Determinants of Obesity in Adults in the Bahamas. *West Indian Med J* 2011; 60 (4): 434
 - 2002: Monteiro CA, Conde WL and Popkin BA. (2007). Income-specific trends in obesity in Brazil: 1975 - 2003. *American Journal of Public Health*, 97 (10): 1808 - 1812.
 - 2003: Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, & Flegal KM. (2006). Prevalence of Overweight and Obesity in the United States, 1999-2004. *The Journal of the American Medical Association*, Vol 295(13):1549 - 1555.
 - 2005: 1st Argentinian National Survey of Risk Factors (Encuesta Nacional de Factores de Riesgo). Results from 1-3rd survey reported in the 4th survey report: https://www.indec.gob.ar/ftp/cuadros/publicaciones/enfr_2018_resultados_definitivos.pdf
 - 2006: Olaiz-Fernández G, Rivera-Dommarco J, Shamah-Levy T, Rojas R, Villalpando-Hernández S, Hernández-Avila M, Sepúlveda-Amor J. *Encuesta Nacional de Salud y Nutrición 2006*. Cuernavaca, México: Instituto Nacional de Salud Pública, 2006. (National Health and Nutrition Survey 2006)

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with obesity, selected countries, 1960-2023


Men



- References:
- 1960, 1971, 1973, 1976, 1988, 1991: Flegal KM, Carroll MD, Kuczmarski RJ, Johnson CL. Overweight and obesity in the United States: prevalence and trends, 1960-1994. *International Journal of Obesity* (1998);22:39-47
 - 1975: Monteiro CA, Conde WL, Popkin BM. Is obesity replacing or adding to undernutrition? Evidence from different social classes in Brazil. 2002. *Public Health Nutrition*:51(1A), 105-112
 - 1981, 1986: Yoshiike N, Seino F, Tajima S, Arai Y, Kawano M, Furuhata T, Inoue S. Twenty-year changes in the prevalence of overweight in Japanese adults: The National Nutrition Survey 1976-95. *Obesity Reviews* 2002;3:183-190
 - 1982, 1993: Rodriguez-Ojea A, Jimenez S, Berdasco A, Esquivel M. The nutrition transition in Cuba in the nineties:an overview. *Public health Nutrition* 2002;5(1A), 129-133
 - 1985: Berg C, Rosengren A, Aires N, appas G, Toren K, Thelle D, Lissner L. Trends in overweight and obesity from 1985 to 2002 in Goteborg, West Sweden. *IJO* 2005 Aug;29(8):916-24
 - 1990: Berg C, Rosengren A, Aires N, appas G, Toren K, Thelle D, Lissner L. Trends in overweight and obesity from 1985 to 2002 in Goteborg, West Sweden. *IJO* 2005 online published ahead of print.
 - 1992: Uauy R, Albal C, Kain J. Obesity Trends in Latin America: Transiting from Under-to Overweight. *Journal of Nutrition* 2001;131:S893-S899
 - 1995: Health Survey for England 1995.
 - 1996: Health Survey for England 1996.
 - 1997: Filozof C, Gonzales C, Sereday M, Mazza C, Braguinsky J. Obesity prevalence and trends in Latin American countries. *Obesity Reviews*, 2001;2:99-196
 - 1998: Scottish Health Survey 1998
 - 1999: Health Survey for England 1999.
 - 2000: Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of Overweight and Obesity in the United States, 1999-2004. *JAMA* 2006;295(13):1549-1555
 - 2001: Health Survey for England 2001.
 - 2002: Monteiro CA, Conde WL and Popkin BA. (2007). Income-specific trends in obesity in Brazil: 1975 - 2003. *American Journal of Public Health*, 97 (10): 1808 - 1812.
 - 2003: 2002 ENS Report. Final results on the National Health Survey.

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women




- References:
- 1960, 1971, 1973, 1976, 1988, 1991: Flegal KM, Carroll MD, Kuczmarski RJ, Johnson CL. Overweight and obesity in the United States: prevalence and trends, 1960-1994. *International Journal of Obesity* (1998);22:39-47
 - 1975: Monteiro CA, Conde WL, Popkin BM. Is obesity replacing or adding to undernutrition? Evidence from different social classes in Brazil. 2002. *Public Health Nutrition*:51(1A), 105-112
 - 1981, 1986: Yoshiike N, Seino F, Tajima S, Arai Y, Kawano M, Furuhata T, Inoue S. Twenty-year changes in the prevalence of overweight in Japanese adults: The National Nutrition Survey 1976-95. *Obesity Reviews* 2002;3:183-190
 - 1982, 1993: Rodriguez-Ojea A, Jimenez S, Berdasco A, Esquivel M. The nutrition transition in Cuba in the nineties:an overview. *Public health Nutrition* 2002;5(1A), 129-133
 - 1985: Berg C, Rosengren A, Aires N, :appas G, Toren K, Thelle D, Lissner L. Trends in overweight and obesity from 1985 to 2002 in Goteborg, West Sweden. *IJO* 2005 Aug;29(8):916-24
 - 1990: Berg C, Rosengren A, Aires N, :appas G, Toren K, Thelle D, Lissner L. Trends in overweight and obesity from 1985 to 2002 in Goteborg, West Sweden. *IJO* 2005 online published ahead of print.
 - 1992: Uauy R, Albal C, Kain J. Obesity Trends in Latin America: Transiting from Under-to Overweight. *Journal of Nutrition* 2001;131:S893-S899
 - 1995: Health Survey for England 1995.
 - 1996: Health Survey for England 1996.
 - 1997: Filozof C, Gonzales C, Sereday M, Mazza C, Braguinsky J. Obesity prevalence and trends in Latin American countries. *Obesity Reviews*, 2001;2:99-196
 - 1998: Scottish Health Survey 1998
 - 1999: Health Survey for England 1999.
 - 2000: Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of Overweight and Obesity in the United States, 1999-2004. *JAMA* 2006;295(13):1549-1555
 - 2001: Health Survey for England 2001.
 - 2002: Monteiro CA, Conde WL and Popkin BA. (2007). Income-specific trends in obesity in Brazil: 1975 - 2003. *American Journal of Public Health*, 97 (10): 1808 - 1812.
 - 2002: 2002 ENS Report. Final results on the National Health Survey.

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Overweight/obesity by education

Men, 2019

Obesity Overweight



Survey type: Self-reported

Age: 18+

Sample size: 52443

Area covered: Regional


References: Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil

Notes: Data from Capitals of 26 Brazilian States and the Federal District Education based on years of education

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2019

■ Obesity ■ Overweight



Survey type: Self-reported

Age: 18+

Sample size: 52443

Area covered: Regional


References: Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil

Notes: Data from Capitals of 26 Brazilian States and the Federal District Education based on years of education

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Boys, 2007


■ Overweight or obesity



Survey type:	Measured
Age:	7-14
Sample size:	2826
Area covered:	Subnational - Municipality of Florianópolis
References:	Bernardo, Carla de Oliveira, & Vasconcelos, Francisco de Assis Guedes de. (2012). Association of parents' nutritional status, and sociodemographic and dietary factors with overweight/obesity in schoolchildren 7 to 14 years old. <i>Cadernos de Saúde Pública</i> , 28(2), 291-304. Retrieved April 04, 2016, from http://www.scielosp.org/scielo.php?script=sci_arttext&pid=S0102-311X2012000200008&lng=en&tlang=en .
Notes:	Prevalence of obesity based on Father's schooling years. For the analyses, the schoolchildren were classified into two groups: not overweight or obese (values equivalent to BMI < 25kg/m ² in adults) and overweight or obese (values equivalent to BMI ≥ 25kg/m ² in adults)
Cutoffs:	IOTF


Girls, 2007

■ Overweight or obesity



Survey type:	Measured
Age:	7-14
Sample size:	2826
Area covered:	Subnational - Municipality of Florianópolis
References:	Bernardo, Carla de Oliveira, & Vasconcelos, Francisco de Assis Guedes de. (2012). Association of parents' nutritional status, and sociodemographic and dietary factors with overweight/obesity in schoolchildren 7 to 14 years old. <i>Cadernos de Saúde Pública</i> , 28(2), 291-304. Retrieved April 04, 2016, from http://www.scielosp.org/scielo.php?script=sci_arttext&pid=S0102-311X2012000200008&lng=en&tlang=en .
Notes:	Prevalence of obesity based on Father's schooling years. For the analyses, the schoolchildren were classified into two groups: not overweight or obese (values equivalent to BMI < 25kg/m ² in adults) and overweight or obese (values equivalent to BMI ≥ 25kg/m ² in adults)
Cutoffs:	IOTF

Infants, 1996



Sample size:

3854

References: DHS: Pesquisa nacional sobre demografia e saude 1996. Demographic and Health Surveys. Rio de Janeiro, Brasil: Litografia Tucano Ltda., 1997


Notes: UNICEF/WHO/World Bank Joint Child Malnutrition Estimates Expanded Database: Overweight (Survey Estimates), May 2023, New York. For more information about the methodology, please consult <https://data.unicef.org/resources/jme-2023-country-consultations/> Percentage of children under 5 years of age falling above 2 standard deviations (moderate and severe) from the median weight-for-height of the reference population.

Definitions:

=>+2SD

Overweight/obesity by age

Adults, 2019



Survey

Self-reported

type:

Sample

52443

size:

Area

Regional

covered:

References: Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil


Notes:

Data from Capitals of 26 Brazilian States and the Federal District

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Children, 2013-2014


█ Obesity █ Overweight



Survey type:	Measured
Sample size:	73399
Area covered:	National
References:	Bloch KV, Klein CH, Szklo M, Kuschnir MCC, Abreu GA, Barufaldi LA et al. ERICA: prevalences of hypertension and obesity in Brazilian adolescents. Rev Saude Publica. 2016;50(suppl 1):9s.
Definitions:	WHO
Cutoffs:	WHO

Overweight/obesity by region

Men, 2019



Survey

Self-reported

type:

Age:

18+

Sample
size:

52443

Area
covered:

Regional


References: Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil

Notes:

Data from Capitals of 26 Brazilian States and the Federal District

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2019



Survey type: Self-reported

Age: 18+

Sample size: 52443

Area covered: Regional


References: Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2019 - report translation provided to WOF by Dra.Andrea Pereira, Presidente e co-fundadora da ONG Obesidade Brasil

Notes: Data from Capitals of 26 Brazilian States and the Federal District

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Boys, 2004-2005

■ Obesity ■ Overweight



Survey type:	Measured
Age:	10-15
Sample size:	36976
Area covered:	National
References:	Pelegrini, Andreia, Luiz Petroski, Edio, da Silva Coqueiro, Raildo, & Araujo Gaya, Adroaldo César. (2008). Overweight and obesity in brazilian schoolchildren aged 10 to 15 years: data from a Brazilian sports project. <i>Archivos Latinoamericanos de Nutrición</i> , 58(4), 343-349. Recuperado en 04 de abril de 2016, de http://www.scielo.org.ve/scielo.php?script=sci_arttext&pid=S0004-06222008000400004&lng=es&tlang=en .
Notes:	International cut-off BMI values were used for the evaluation of nutritional status (Cole et al)
Cutoffs:	IOTF


Girls, 2004-2005



Survey type:	Measured
Age:	10-15
Sample size:	36976
Area covered:	National
References:	Pelegrini, Andreia, Luiz Petroski, Edio, da Silva Coqueiro, Raildo, & Araujo Gaya, Adroaldo César. (2008). Overweight and obesity in brazilian schoolchildren aged 10 to 15 years: data from a Brazilian sports project. <i>Archivos Latinoamericanos de Nutrición</i> , 58(4), 343-349. Recuperado en 04 de abril de 2016, de http://www.scielo.org.ve/scielo.php?script=sci_arttext&pid=S0004-06222008000400004&lng=es&tlang=en .
Notes:	International cut-off BMI values were used for the evaluation of nutritional status (Cole et al)
Cutoffs:	IOTF

Infants, 2006-2007

Overweight or obesity



Sample
size:

12374534

References: Other: Pesquisa nacional de demografia e saúde da criança e da mulher - PNDS 2006. Relatório da pesquisa. Sao Paulo:
CEBRAP, 2008

Notes: UNICEF/WHO/World Bank Joint Child Malnutrition Estimates Expanded Database: Overweight (Survey Estimates), May 2023,
New York. For more information about the methodology, please consult <https://data.unicef.org/resources/jme-2023-country-consultations/> Percentage of children under 5 years of age falling above 2 standard deviations (moderate and severe) from the
median weight-for-height of the reference population.


Definitions:

=>+2SD

Overweight/obesity by socio-economic group

Adults, 2008-2010

■ Obesity ■ Overweight



Survey type: Measured

Age: 20-59

Sample size: 527

Area covered: Regional - N. Eastern Brazil


References: Lima R P A, Pereira D C, Luna R C P, et al. BMI, Overweight Status and Obesity Adjusted by Various Factors in All Age Groups in the Population of a City in Northeastern Brazil. *Int. J. Environ. Res. Public Health* 2015, 12, 4422-4438; doi:10.3390/ijerph120404422

Notes: WHO BMI classification of 1995 used for Adults. Median family income, R\$1000.00, or \$492.02.

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Children, 2010

■ Obesity ■ Overweight




Survey type:	Measured
Age:	6-10
Sample size:	939
Area covered:	Municipality of Cruzeiro do Oeste, Southern Brazil.
References: Azambuja, Ana Paula de O., Netto-Oliveira, Edna Regina, Oliveira, Amauri Aparecido B. de, Azambuja, Maximiliano dos Anjos, & Rinaldi, Wilson. (2013). Prevalence of overweight/obesity and economical status of schoolchildren. Revista Paulista de Pediatria, 31(2), 166-171. https://dx.doi.org/10.1590/S0103-05822013000200006	

Notes: The socioeconomic status was defined through a questionnaire of economic classification by the Brazilian Association of Research Companies – Associação Brasileira de Empresas de Pesquisa, ABEP. This classification is based on items such as ownership of goods (television, radio, car, vacuum cleaner, VCR and/or DVD player, fridge, freezer and washing machine), services (housemaid), household characteristics (number of bathrooms) and educational level of the head of the household. The total score for each item results in the classification of respondents into seven strata identified as "social classes" A1, A2, B1, B2, C, D, and E(17). For analysis purposes, the eight economic levels, proposed by ABEP, were regrouped and named as follows: A1, A2, and B1 in High Economic Level (H); B2, C1 and C2 in Medium Economic Level (M) and, D and E in Low Economic Level (L).

Cutoffs: Other

Infants, 2019

Overweight or obesity



Sample size:

10780287

References:

NNS: Estudo Nacional de Alimentação e Nutrição Infantil (ENANI), Brasil 2019


Notes: UNICEF/WHO/World Bank Joint Child Malnutrition Estimates Expanded Database: Overweight (Survey Estimates), May 2023, New York. For more information about the methodology, please consult <https://data.unicef.org/resources/jme-2023-country-consultations/> Percentage of children under 5 years of age falling above 2 standard deviations (moderate and severe) from the median weight-for-height of the reference population.

Definitions:

=>+2SD


Insufficient physical activity

Adults, 2016




References: Guthold R, Stevens GA, Riley LM, Bull FC. Worldwide trends in insufficient physical activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1.9 million participants. Lancet 2018 [http://dx.doi.org/10.1016/S2214-109X\(18\)30357-7](http://dx.doi.org/10.1016/S2214-109X(18)30357-7)

Men, 2016




References: Guthold R, Stevens GA, Riley LM, Bull FC. Worldwide trends in insufficient physical activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1.9 million participants. Lancet 2018 [http://dx.doi.org/10.1016/S2214-109X\(18\)30357-7](http://dx.doi.org/10.1016/S2214-109X(18)30357-7)

Women, 2016



References: Guthold R, Stevens GA, Riley LM, Bull FC. Worldwide trends in insufficient physical activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1.9 million participants. Lancet 2018 [http://dx.doi.org/10.1016/S2214-109X\(18\)30357-7](http://dx.doi.org/10.1016/S2214-109X(18)30357-7)

Children, 2016



Survey type:

Self-reported

Age:

11-17

References:

Global Health Observatory data repository, World Health Organisation,

<https://apps.who.int/gho/data/node.main.A893ADO?lang=en> (last accessed 16.03.21)


Notes:

% of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.

Definitions:

% Adolescents insufficiently active (age standardised estimate)

Boys, 2016



Survey type:

Self-reported

Age:

11-17

References:

Global Health Observatory data repository, World Health Organisation,

<https://apps.who.int/gho/data/node.main.A893ADO?lang=en> (last accessed 16.03.21)


Notes:

% of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.

Definitions:

% Adolescents insufficiently active (age standardised estimate)

Girls, 2016



Survey type:

Self-reported

Age:

11-17

References:

Global Health Observatory data repository, World Health Organisation,

<https://apps.who.int/gho/data/node.main.A893ADO?lang=en> (last accessed 16.03.21)

Notes:


% of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.

Definitions:

% Adolescents insufficiently active (age standardised estimate)

Estimated per capita fruit intake

Adults, 2017



Survey type:

Measured

Age:

25+

References:


Global Burden of Disease, the Institute for Health Metrics and Evaluation <http://ghdx.healthdata.org/>

Definitions:

Estimated per-capita fruit intake (g/day)

Estimated per-capita processed meat intake

Adults, 2017



Survey type:

Measured

Age:

25+

References:


Global Burden of Disease, the Institute for Health Metrics and Evaluation <http://ghdx.healthdata.org/>

Definitions:

Estimated per-capita processed meat intake (g per day)

Estimated per capita whole grains intake

Adults, 2017



Survey type:

Measured

Age:

25+

References:


Global Burden of Disease, the Institute for Health Metrics and Evaluation <http://ghdx.healthdata.org/>

Definitions:

Estimated per-capita whole grains intake (g/day)

Mental health - depression disorders

Adults, 2015




References: Prevalence data from Global Burden of Disease study 2015 (<http://ghdx.healthdata.org>) published in: Depression and Other Common Mental Disorders: Global Health Estimates. Geneva:World Health Organization; 2017. Licence: CC BY-NC-SA 3.0 IGO.

Definitions:

% of population with depression disorders

Mental health - anxiety disorders

Adults, 2015




References: Prevalence data from Global Burden of Disease study 2015 (<http://ghdx.healthdata.org>) published in: Depression and Other Common Mental Disorders: Global Health Estimates. Geneva:World Health Organization; 2017. Licence: CC BY-NC-SA 3.0 IGO.

Definitions:

% of population with anxiety disorders

% Infants exclusively breastfed 0-5 months

Children, 1998-2019



Area covered: National


References: Pesquisa nacional de demografia e saúde da criança e da mulher - PNDS 2006. Relatório da pesquisa. Sao Paulo: CEBRAP, 2008

Notes: See UNICEF website for further survey information. Available at : <https://data.unicef.org/resources/dataset/infant-young-child-feeding/> (last accessed 28.9.21) Citation: United Nations Children's Fund, Division of Data, Analysis, Planning and Monitoring (2021). Global UNICEF Global Databases: Infant and Young Child Feeding: Exclusive breastfeeding, New York, September 2021.

Definitions: % exclusively breastfed 0-5 months

Oesophageal cancer

Men, 2020



Age:

20+

Area

National

covered:


References:

Ferlay J, Ervik M, Lam F, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2020). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.fr/today>, accessed 10.01.2065

Definitions:

Age-standardized incidence rates per 100 000

Women, 2020



Age:

20+

Area covered:

National

References:


Ferlay J, Ervik M, Lam F, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2020). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.fr/today>, accessed 10.01.2065

Definitions:

Age-standardized incidence rates per 100 000

Breast cancer

Women, 2020



Age:

20+

Area

National

covered:

References:


Ferlay J, Ervik M, Lam F, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2020). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.fr/today>, accessed 10.01.2065

Definitions:

Age-standardized incidence rates per 100 000

Colorectal cancer

Men, 2020



Age:

20+

Area

National

covered:


References:

Ferlay J, Ervik M, Lam F, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2020). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.fr/today>, accessed 10.01.2065

Definitions:

Age-standardized incidence rates per 100 000

Women, 2020



Age:

20+

Area covered:

National

References:


Ferlay J, Ervik M, Lam F, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2020). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.fr/today>, accessed 10.01.2065

Definitions:

Age-standardized incidence rates per 100 000

Pancreatic cancer

Men, 2020



Age:

20+

Area

National

covered:


References:

Ferlay J, Ervik M, Lam F, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2020). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.fr/today>, accessed 10.01.2065

Definitions:

Age-standardized incidence rates per 100 000

Women, 2020



Age:

20+

Area covered:

National

References:


Ferlay J, Ervik M, Lam F, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2020). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.fr/today>, accessed 10.01.2065

Definitions:

Age-standardized incidence rates per 100 000

Gallbladder cancer

Men, 2020



Age:

20+

Area

National

covered:


References:

Ferlay J, Ervik M, Lam F, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2020). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.fr/today>, accessed 10.01.2065

Definitions:

Age-standardized incidence rates per 100 000

Women, 2020



Age:

20+

Area covered:

National

References:


Ferlay J, Ervik M, Lam F, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2020). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.fr/today>, accessed 10.01.2065

Definitions:

Age-standardized incidence rates per 100 000

Kidney cancer

Men, 2020



Age:

20+

Area

National

covered:


References:

Ferlay J, Ervik M, Lam F, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2020). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.fr/today>, accessed 10.01.2065

Definitions:

Age-standardized incidence rates per 100 000

Women, 2020



Age:

20+

Area covered:

National

References:


Ferlay J, Ervik M, Lam F, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2020). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.fr/today>, accessed 10.01.2065

Definitions:

Age-standardized incidence rates per 100 000

Cancer of the uterus

Women, 2020



Age:

20+

Area

National

covered:

References:


Ferlay J, Ervik M, Lam F, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2020). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.fr/today>, accessed 10.01.2065

Definitions:

Age-standardized incidence rates per 100 000

Raised blood pressure

Adults, 2015



References:


Global Health Observatory data repository, World Health Organisation,

<http://apps.who.int/gho/data/node.main.A875?lang=en>

Definitions:

Age Standardised estimated % Raised blood pressure 2015 (SBP>=140 OR DBP>=90).

Men, 2015



References:


Global Health Observatory data repository, World Health Organisation,

<http://apps.who.int/gho/data/node.main.A875?lang=en>

Definitions:

Age Standardised estimated % Raised blood pressure 2015 (SBP>=140 OR DBP>=90).

Women, 2015



References:

Global Health Observatory data repository, World Health Organisation,


<http://apps.who.int/gho/data/node.main.A875?lang=en>

Definitions:

Age Standardised estimated % Raised blood pressure 2015 (SBP>=140 OR DBP>=90).

Raised cholesterol

Adults, 2008




References:

Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A885>

Definitions:

% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).


Men, 2008



References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A885>

Definitions: % Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).

Women, 2008



References:


Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A885>

Definitions:

% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).

Raised fasting blood glucose

Men, 2014



References:


Global Health Observatory data repository, World Health Organisation,

<http://apps.who.int/gho/data/node.main.A869?lang=en>

Definitions:

Age Standardised % raised fasting blood glucose (≥ 7.0 mmol/L or on medication).

Women, 2014



References:

Global Health Observatory data repository, World Health Organisation,


<http://apps.who.int/gho/data/node.main.A869?lang=en>

Definitions:

Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication).

Diabetes prevalence

Adults, 2021



Age: 20-79

Area covered: National

References: Reproduced with kind permission International Diabetes Federation. IDF Diabetes Atlas, 10th edn. Brussels, Belgium:International Diabetes Federation, 2021. <http://www.diabetesatlas.org>

Definitions: Age-adjusted comparative prevalence of diabetes, %

Contextual factors

Disclaimer: These contextual factors should be interpreted with care. Results are updated as regularly as possible and use very specific criteria. The criteria used and full definitions are available for download at the bottom of this page.



Labelling

Is there mandatory nutrition labelling? ✓

Front-of-package labelling? ✓

Back-of-pack nutrition declaration? ✓

Color coding? ✗

Warning label? ✓



Regulation and marketing

Are there fiscal policies on unhealthy products?



Tax on unhealthy foods?



Tax on unhealthy drinks?



Are there fiscal policies on healthy products?



Subsidy on fruits?



Subsidy on vegetables?



Subsidy on other healthy products?



Mandatory limit or ban of trans fat (all settings)?



Mandatory limit of trans fats in place (all settings)?



Ban on trans-fats or phos in place (all settings)?



Are there any mandatory policies/marketing restrictions on the promotion of unhealthy food/drinks to children?



Mandatory restriction on broadcast media?



Mandatory restriction on non-broadcast media?



Voluntary policies/marketing restrictions on the promotion of unhealthy food/drinks to children?



Are there mandatory standards for food in schools?



Are there any mandatory nutrient limits in any manufactured food products?



Nutrition standards for public sector procurement?





Political will and support

National obesity strategy or nutrition and physical activity national strategy? ✓

National obesity strategy? ✓

National childhood obesity strategy? ✗

Comprehensive nutrition strategy? ✓

Comprehensive physical activity strategy? ✓

Evidence-based dietary guidelines and/or RDAs? ✓

National target(s) on reducing obesity? ✓

Guidelines/policy on obesity treatment? ✓

Promotion of breastfeeding? ✓



Monitoring and surveillance

Monitoring of the prevalence and incidence for the main obesity-related NCDs and risk factors? ✓

Within 5 years? ✓



Governance and resource

Multi-sectoral national co-ordination mechanism for obesity or nutrition (including obesity)? ✓

Key

✓ Present

✓ v Present

✓ Incoming

✗ Absent

? Unknown

(voluntary)