

Report card

China


Upper-middle income

	Page
Obesity prevalence	3
Trend: % Adults living with obesity, 2002-2019	6
Trend: % Adults living with obesity, 2002-2019	8
Trend: % Adults living with overweight or obesity by urbanisation in China 1993-2009	10
Trend: % Adults living with overweight or obesity in China 1993-2009	11
Trend: % Adults living with overweight or obesity, 2002-2019	12
Trend: % Adults living with overweight or obesity, 2002-2019	14
Trend: Children living with Overweight or obesity in China	16
Overweight/obesity by education	17
Overweight/obesity by age	19
Overweight/obesity by region	21
Overweight/obesity by age and region	25
Overweight/obesity by socio-economic group	27
Overweight/obesity by ethnicity	29
Insufficient physical activity	30
Estimated per capita fruit intake	36
Estimated per-capita processed meat intake	37
Estimated per capita whole grains intake	38
Mental health - depression disorders	39
Mental health - anxiety disorders	40
% Infants exclusively breastfed 0-5 months	41
Oesophageal cancer	42
Breast cancer	44
Colorectal cancer	45
Pancreatic cancer	47
Gallbladder cancer	49
Kidney cancer	51
Cancer of the uterus	53
Raised blood pressure	54
Raised cholesterol	57
Raised fasting blood glucose	60
Diabetes prevalence	62
Ovarian Cancer	63
Leukemia	64
Leukemia	66
Liver and intrahepatic bile duct Cancer	68
Multiple Myeloma	70

Contents	Page
Non Hodgkin Lymphoma	%%
Thyroid Cancer	%%
Contextual factors	%%

Obesity prevalence


Adults, 2018-2019



Survey type:	Measured
Age:	18+
Sample size:	155413
Area covered:	National
References:	Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. The Lancet Regional Health: Western Pacific. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809
Notes:	Chinese Cut Offs applied
Definitions:	Overweight BMI ≥ 24 - < 28 Kg/m ² Obesity BMI ≥ 28 Kg/m ²

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Children, 2015



Survey type:	Measured
Age:	7-18
Sample size:	1617
Area covered:	National
References:	Zhang J, Wang H, Wang Z, Du W, Su C, Zhang J, Jiang H, Jia X, Huang F, Ouyang Y, Wang Y, Zhang B. Prevalence and stabilizing trends in overweight and obesity among children and adolescents in China, 2011-2015. BMC Public Health. 2018 May 2;18(1):571. doi: 10.1186/s12889-018-5483-9.
Definitions:	IOTF International Cut off. WHO & WGO also available in paper.
Cutoffs:	IOTF

Infants, 2016-2017

■ Overweight or obesity



Survey type: Measured

Age: 0-5

Area covered: National


References: Other: China Nutrition and Health Surveillance of Children and Lactating Women 2016–2017

Notes: UNICEF/WHO/World Bank Joint Child Malnutrition Estimates Expanded Database: Overweight (Survey Estimates), May 2023, New York. For more information about the methodology, please consult <https://data.unicef.org/resources/jme-2023-country-consultations/> Percentage of children under 5 years of age falling above 2 standard deviations (moderate and severe) from the median weight-for-height of the reference population.

Definitions: =>+2SD

% Adults living with obesity, 2002-2019

Men



Survey type:

Measured


References: Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809
 Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809
 Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809

Definitions:

Overweight BMI ≥ 24 - < 28 Kg/m² Obesity BMI ≥ 28 Kg/m²

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women



Survey
type:

Measured

References: Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809
 Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809
 Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809


Definitions:

Overweight BMI ≥ 24 – < 28 Kg/m² Obesity BMI ≥ 28 Kg/m²

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with obesity, 2002-2019

Men




References: Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809
 Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809
 Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809

Definitions: Overweight BMI ≥ 24 - < 28 Kg/m² Obesity BMI ≥ 28 Kg/m²

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women



References: Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809

Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809


Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809

Definitions: Overweight BMI ≥ 24 - < 28 Kg/m² Obesity BMI ≥ 28 Kg/m²

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with overweight or obesity by urbanisation in China 1993-2009



Survey type:


Measured

References: Xi, B., Liang, Y., He, T., Reilly, K. H., Hu, Y., Wang, Q., Yan, Y. and Mi, J. (2012), Secular trends in the prevalence of general and abdominal obesity among Chinese adults, 1993-2009. *Obesity Reviews*, 13: 287-296. doi: 10.1111/j.1467-789X.2011.00944.x

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with overweight or obesity in China 1993-2009



Survey
type:

Measured


References: Xi, B., Liang, Y., He, T., Reilly, K. H., Hu, Y., Wang, Q., Yan, Y. and Mi, J. (2012), Secular trends in the prevalence of general and abdominal obesity among Chinese adults, 1993-2009. *Obesity Reviews*, 13: 287-296. doi: 10.1111/j.1467-789X.2011.00944.x

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with overweight or obesity, 2002-2019

Men



Survey type:

Measured


References: Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809
 Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809
 Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809

Definitions:

Overweight BMI ≥ 24 - < 28 Kg/m² Obesity BMI ≥ 28 Kg/m²

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women



Survey type:

Measured

References: Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809
 Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809
 Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809


Definitions:

Overweight BMI ≥ 24 – < 28 Kg/m² Obesity BMI ≥ 28 Kg/m²

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with overweight or obesity, 2002-2019

Men




References: Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809
 Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809
 Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809

Definitions: Overweight BMI ≥ 24 - < 28 Kg/m² Obesity BMI ≥ 28 Kg/m²

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women



References: Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809

Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809


Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809

Definitions: Overweight BMI ≥ 24 – < 28 Kg/m² Obesity BMI ≥ 28 Kg/m²

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Children living with Overweight or obesity in China



Survey
type:


Measured

- References:
- 1982: Li Y., Schouten EG, Hu X et al. Obesity prevalence and time trend among youngsters in China, 1982-2002. Published in Asia Pacific Journal of Clinical Nutrition. Our version from Book - Li Y. Childhood Obesity in China: prevalence, determinants and health.
 - 1992: Li Y., Schouten EG, Hu X et al. Obesity prevalence and time trend among youngsters in China, 1982-2002. Published in Asia Pacific Journal of Clinical Nutrition. Our version from Book - Li Y. Childhood Obesity in China: prevalence, determinants and health. Chapter 2
 - 2002: Yanping L, Evert GS, Xiaoqi H, Zhaohui C, Dechun L and Guansheng M. 2008. Obesity prevalence and time trend among youngsters in china, 1982 - 2002. Asia Pac Journal of Clinical Nutrition, 17(1):131 - 137.

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Overweight/obesity by education

Adults, 2015



Survey type: Measured

Age: 18-59

Sample size: 6602


Area covered: National

References: Huang, Q., Wang, L., Jiang, H., Wang, H., Zhang, B., Zhang, J., Jia, X. and Wang, Z., 2020. Intra-Individual Double Burden of Malnutrition among Adults in China: Evidence from the China Health and Nutrition Survey 2015. *Nutrients*, 12(9), p.2811.

Notes: 3699 men and 2903 women

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Children, 2014



Survey type: Measured

Age: 5-12

Sample size: 9917

Area covered: Regional - Guangzhou (urban setting)


References: Liu W, Liu W, Lin R, et al. Socioeconomic determinants of childhood obesity among primary school children in Guangzhou, China. BMC Public Health. 2016;16:482. doi:10.1186/s12889-016-3171-1.

Notes: BMI standard deviation scores (BMI z-score) were derived using the age (calculated by subtracting the date of birth from the date of examination) and sex specific WHO growth reference for school-aged children, which were further classified as non-overweight ($\leq 1SD$), overweight ($>1SD$) and obese ($>2SD$). Overweight and obesity prevalence by Father's education. Education was categorised into low (primary and junior high school level), medium (senior high and vocational school level) or high (university level or higher).

Cutoffs: WHO

Overweight/obesity by age

Adults, 2018-2019



Survey type: Measured

Sample size: 155413

Area covered: National


References: Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809

Notes: Chinese Cut Off applied

Definitions: Overweight BMI ≥ 24 -<28 Kg/m² Obesity BMI ≥ 28 Kg/m²

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Children, 2015



Survey type:	Measured
Sample size:	1617
Area covered:	National
References:	Zhang J, Wang H, Wang Z, Du W, Su C, Zhang J, Jiang H, Jia X, Huang F, Ouyang Y, Wang Y, Zhang B. Prevalence and stabilizing trends in overweight and obesity among children and adolescents in China, 2011-2015. BMC Public Health. 2018 May 2;18(1):571. doi: 10.1186/s12889-018-5483-9.
Notes:	International Cut off used, WHO & WGOC also available
Cutoffs:	IOTF


Overweight/obesity by region

Men, 2018-2019



Survey type:	Measured
Age:	18+
Sample size:	155413
Area covered:	National
References:	Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. <i>The Lancet Regional Health: Western Pacific</i> . 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809
Definitions:	Overweight BMI ≥ 24 -<28 Kg/m ² Obesity BMI ≥ 28 Kg/m ²
<i>Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².</i>	

Women, 2018-2019



Survey type: Measured

Age: 18+

Sample size: 155413


Area covered: National

References: Wen Peng, Shiqi Chen and Xinguang Chen et al. Trends in major non-communicable diseases and related risk factors in China 2002–2019: an analysis of nationally representative survey data. *The Lancet Regional Health: Western Pacific*. 2023. Vol. 43. DOI: 10.1016/j.lanwpc.2023.100809

Definitions: Overweight BMI ≥ 24 -<28 Kg/m² Obesity BMI ≥ 28 Kg/m²


Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Children, 2015



Survey type:	Measured
Age:	7-18
Sample size:	1617
Area covered:	National
References:	Zhang J, Wang H, Wang Z, Du W, Su C, Zhang J, Jiang H, Jia X, Huang F, Ouyang Y, Wang Y, Zhang B. Prevalence and stabilizing trends in overweight and obesity among children and adolescents in China, 2011-2015. BMC Public Health. 2018 May 2;18(1):571. doi: 10.1186/s12889-018-5483-9.
Notes:	International Cut off used, WHO & WGO also available
Cutoffs:	IOTF

Infants, 2010



Sample size: 15399


References: Surveillance: Nutritional status of children aged 0-5 years old in China (2010) - National (38 nutrition surveillance sites from 25 provinces). Beijing, China: Chinese Center for Disease Control and Prevention, 2012

Notes: UNICEF/WHO/World Bank Joint Child Malnutrition Estimates Expanded Database: Overweight (Survey Estimates), May 2023, New York. For more information about the methodology, please consult <https://data.unicef.org/resources/jme-2023-country-consultations/> Percentage of children under 5 years of age falling above 2 standard deviations (moderate and severe) from the median weight-for-height of the reference population.

Definitions: =>+2SD

Overweight/obesity by age and region

Men, 2005



Survey type: Measured

Age: 18

Sample size: 30447


Area covered: Regional

References: Zhang YX, Wang SR. Distribution of body mass index and the prevalence changes of overweight and obesity among adolescents in Shandong, China from 1985 to 2005. *Annals of Human Biology*, Volume 35, Issue 5 September 2008 , pages 547 - 555

Cutoffs: Other

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2005




Survey type:	Measured
Age:	18
Sample size:	30447
Area covered:	Regional
References:	Zhang YX, Wang SR. Distribution of body mass index and the prevalence changes of overweight and obesity among adolescents in Shandong, China from 1985 to 2005. <i>Annals of Human Biology</i> , Volume 35, Issue 5 September 2008 , pages 547 - 555
Cutoffs:	Other

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Overweight/obesity by socio-economic group

Adults, 2015



Survey type: Measured

Age: 18-59

Sample size: 6602


Area covered: National

References: Huang, Q., Wang, L., Jiang, H., Wang, H., Zhang, B., Zhang, J., Jia, X. and Wang, Z., 2020. Intra-Individual Double Burden of Malnutrition among Adults in China: Evidence from the China Health and Nutrition Survey 2015. *Nutrients*, 12(9), p.2811.

Notes: 3699 men and 2903 women

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Children, 2014



Survey type: Measured

Age: 5-12

Sample size: 9917

Area covered: Regional - Guangzhou (urban setting)

References: Liu W, Liu W, Lin R, et al. Socioeconomic determinants of childhood obesity among primary school children in Guangzhou, China. BMC Public Health. 2016;16:482. doi:10.1186/s12889-016-3171-1.


Notes: BMI standard deviation scores (BMI z-score) were derived using the age (calculated by subtracting the date of birth from the date of examination) and sex specific WHO growth reference for school-aged children, which were further classified as non-overweight ($\leq 1SD$), overweight ($>1SD$) and obese ($>2SD$). Overweight and obesity prevalence by Father's education. Education was categorised into low (primary and junior high school level), medium (senior high and vocational school level) or high (university level or higher).

Cutoffs: WHO

Overweight/obesity by ethnicity

Ethnic groups are as defined by publication of origin and are not as defined by WOF. In some instances ethnicity is conflated with nationality and/or race.

Children, 2014



Survey type: Measured

Age: 7-18

Sample size: 80,821

Area covered: National


References: Dong, Yanhui, et al. "Prevalence of Excess Body Weight and Underweight among 26 Chinese Ethnic Minority Children and Adolescents in 2014: A Cross-Sectional Observational Study." *BMC Public Health*, vol. 18, no. 1, 27 Apr. 2018, 10.1186/s12889-018-5352-6.

Notes: Data from Chinese National Survey on Students Constitution and Health 2014

Cutoffs: Overweight and obesity was defined as \geq the referent age-and sex- specific 85th centile according to the reference developed by Working Group on Obesity in China (WGO)

Insufficient physical activity

Adults, 2022



Survey type: Self-reported


Age: 18+

Area covered: National

References: WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-\(age-standardized-estimate\)-\(-\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-))

Definitions: Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.

Men, 2022



Survey type: Self-reported


Age: 18+

Area covered: National

References: WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-\(age-standardized-estimate\)-\(-\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-))

Definitions: Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.

Women, 2022



Survey type: Self-reported


Age: 18+

Area covered: National

References: WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-\(age-standardized-estimate\)-\(-\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-))

Definitions: Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.

Children, 2016



Survey type: Self-reported


Age: 11-17

References: Global Health Observatory data repository, World Health Organisation, <https://apps.who.int/gho/data/node.main.A893ADO?lang=en> (last accessed 16.03.21)

Notes: % of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.

Definitions: % Adolescents insufficiently active (age standardised estimate)

Boys, 2016



Survey type: Self-reported


Age: 11-17

References: Global Health Observatory data repository, World Health Organisation, <https://apps.who.int/gho/data/node.main.A893ADO?lang=en> (last accessed 16.03.21)

Notes: % of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.

Definitions: % Adolescents insufficiently active (age standardised estimate)

Girls, 2016



Survey type: Self-reported

Age: 11-17


References: Global Health Observatory data repository, World Health Organisation, <https://apps.who.int/gho/data/node.main.A893ADO?lang=en> (last accessed 16.03.21)

Notes: % of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.

Definitions: % Adolescents insufficiently active (age standardised estimate)

Estimated per capita fruit intake

Adults, 2017



Survey type: Measured


Age: 25+

References: Global Burden of Disease, the Institute for Health Metrics and Evaluation <http://ghdx.healthdata.org/>

Definitions: Estimated per-capita fruit intake (g/day)

Estimated per-capita processed meat intake

Adults, 2017



Survey type: Measured


Age: 25+

References: Global Burden of Disease, the Institute for Health Metrics and Evaluation <http://ghdx.healthdata.org/>

Definitions: Estimated per-capita processed meat intake (g per day)

Estimated per capita whole grains intake

Adults, 2017



Survey type:

Measured

Age:

25+

References:


Global Burden of Disease, the Institute for Health Metrics and Evaluation <http://ghdx.healthdata.org/>

Definitions:

Estimated per-capita whole grains intake (g/day)

Mental health - depression disorders

Adults, 2015




References: Prevalence data from Global Burden of Disease study 2015 (<http://ghdx.healthdata.org>) published in: Depression and Other Common Mental Disorders: Global Health Estimates. Geneva:World Health Organization; 2017. Licence: CC BY-NC-SA 3.0 IGO.

Definitions: % of population with depression disorders

Mental health - anxiety disorders

Adults, 2015




References: Prevalence data from Global Burden of Disease study 2015 (<http://ghdx.healthdata.org>) published in: Depression and Other Common Mental Disorders: Global Health Estimates. Geneva:World Health Organization; 2017. Licence: CC BY-NC-SA 3.0 IGO.

Definitions: % of population with anxiety disorders

% Infants exclusively breastfed 0-5 months

Infants, 2007-2022



References:


China Nutrition and Health Surveillance of Children and Lactating Women 2016–2017

Notes:

Full details are available. Original citation United Nations Children’s Fund, Division of Data, Analysis, Planning and Monitoring (2023). Global UNICEF Global Databases: Infant and Young Child Feeding: Exclusive breastfeeding, New York, October 2023.

Oesophageal cancer

Men, 2022




Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Women, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Breast cancer

Women, 2022




Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Colorectal cancer

Men, 2022



Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Women, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Pancreatic cancer

Men, 2022




Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Women, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Gallbladder cancer

Men, 2022




Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Women, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Kidney cancer

Men, 2022




Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Women, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Cancer of the uterus

Women, 2022




Age: 20+

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Raised blood pressure

Adults, 2015




References:

Global Health Observatory data repository, World Health Organisation,
<http://apps.who.int/gho/data/node.main.A875?lang=en>

Definitions:

Age Standardised estimated % Raised blood pressure 2015 (SBP \geq 140 OR DBP \geq 90).

Men, 2015




References:

Global Health Observatory data repository, World Health Organisation,
<http://apps.who.int/gho/data/node.main.A875?lang=en>

Definitions:

Age Standardised estimated % Raised blood pressure 2015 (SBP \geq 140 OR DBP \geq 90).

Women, 2015



References:


Global Health Observatory data repository, World Health Organisation,
<http://apps.who.int/gho/data/node.main.A875?lang=en>

Definitions:

Age Standardised estimated % Raised blood pressure 2015 (SBP>=140 OR DBP>=90).

Raised cholesterol


Adults, 2008



References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A885>

Definitions: % Raised total cholesterol (≥ 5.0 mmol/L) (age-standardized estimate).


Men, 2008



References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A885>

Definitions: % Raised total cholesterol (≥ 5.0 mmol/L) (age-standardized estimate).

Women, 2008




References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A885>

Definitions: % Raised total cholesterol (≥ 5.0 mmol/L) (age-standardized estimate).

Raised fasting blood glucose

Men, 2014




References:

Global Health Observatory data repository, World Health Organisation,
<http://apps.who.int/gho/data/node.main.A869?lang=en>

Definitions:

Age Standardised % raised fasting blood glucose (≥ 7.0 mmol/L or on medication).

Women, 2014



References:


Global Health Observatory data repository, World Health Organisation,
<http://apps.who.int/gho/data/node.main.A869?lang=en>

Definitions:

Age Standardised % raised fasting blood glucose (≥ 7.0 mmol/L or on medication).

Diabetes prevalence

Adults, 2021



Age: 20-79


Area covered: National

References: Reproduced with kind permission International Diabetes Federation. IDF Diabetes Atlas, 10th edn. Brussels, Belgium:International Diabetes Federation, 2021. <http://www.diabetesatlas.org>

Definitions: Age-adjusted comparative prevalence of diabetes, %

Ovarian Cancer

Women, 2022



Age: 20+

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Leukemia

Men, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022



Age: 20+

Area covered: National


References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Leukemia

Men, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022



Age: 20+

Area covered: National


References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Liver and intrahepatic bile duct Cancer

Men, 2022




Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Women, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Multiple Myeloma

Men, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022



Age: 20+

Area covered: National


References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Non Hodgkin Lymphoma

Men, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022



Age: 20+

Area covered: National


References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Thyroid Cancer

Men, 2022




Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Women, 2022



Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]



Definitions: Incidence per 100,000

Contextual factors

Disclaimer: These contextual factors should be interpreted with care. Results are updated as regularly as possible and use very specific criteria. The criteria used and full definitions are available for download at the bottom of this page.



Labelling

Is there mandatory nutrition labelling?	
Front-of-package labelling?	
Back-of-pack nutrition declaration?	
Color coding?	
Warning label?	



Regulation and marketing

Are there fiscal policies on unhealthy products?	X
Tax on unhealthy foods?	X
Tax on unhealthy drinks?	X
Are there fiscal policies on healthy products?	X
Subsidy on fruits?	X
Subsidy on vegetables?	X
Subsidy on other healthy products?	X
Mandatory limit or ban of trans fat (all settings)?	X
Mandatory limit of trans fats in place (all settings)?	X
Ban on trans-fats or phos in place (all settings)?	X
Are there any mandatory policies/marketing restrictions on the promotion of unhealthy food/drinks to children?	X
Mandatory restriction on broadcast media?	X
Mandatory restriction on non-broadcast media?	X
Voluntary policies/marketing restrictions on the promotion of unhealthy food/drinks to children?	X
Are there mandatory standards for food in schools?	X
Are there any mandatory nutrient limits in any manufactured food products?	X
Nutrition standards for public sector procurement?	X



Political will and support

National obesity strategy or nutrition and physical activity national strategy?	✓
National obesity strategy?	✓
National childhood obesity strategy?	✓
Comprehensive nutrition strategy?	✓
Comprehensive physical activity strategy?	✓
Evidence-based dietary guidelines and/or RDAs?	✓
National target(s) on reducing obesity?	✓
Guidelines/policy on obesity treatment?	✓
Promotion of breastfeeding?	✓



Monitoring and surveillance

Monitoring of the prevalence and incidence for the main obesity-related NCDs and risk factors?	✓
Within 5 years?	✓



Governance and resource

Multi-sectoral national co-ordination mechanism for obesity or nutrition (including obesity)?	✓
---	---

Key

✓ Present

✓_v Present

(voluntary)

✓ Incoming

✗ Absent

? Unknown

Last updated September 13, 2022

PDF created on September 11, 2024