Cuba



Country report card - adults


This report card contains the latest data available on the Global Obesity Observatory on overweight and obesity for adults. Where available, data on common and relevant obesity drivers and comorbidities are also presented.

View the latest version of this report on the Global Obesity Observatory at https://data.worldobesity.org/country/cuba-52/.

Contents	Page
Obesity prevalence	3
Trend: % Adults living with obesity in Cuba 1982-2019	6
Trend: % Adults living with overweight or obesity in Cuba 1982-2019	11
Trend: % Adults living with obesity in selected countries worldwide 1976-2018, selected countries	16
Overweight/obesity by region	21
Double burden of underweight & obesity	24
Insufficient physical activity	27
Mental health - depression disorders	34
Mental health - anxiety disorders	38
Oesophageal cancer	42
Breast cancer	45
Colorectal cancer	47
Pancreatic cancer	50
Gallbladder cancer	53
Kidney cancer	56
Cancer of the uterus	59
Raised blood pressure	61
Raised cholesterol	65
Raised fasting blood glucose	68
Diabetes prevalence	70
Ovarian Cancer	71
Leukemia	73
Liver and intrahepatic bile duct Cancer	78
Multiple Myeloma	81
Non Hodgkin Lymphoma	86
Thyroid Cancer	91

Obesity prevalence

Adults, 2018-2019




Survey type:	Measured
Age:	15+
Sample size:	14,339
Area covered:	National
References:	Díaz, E., Maldonado, G., Medina, R.S. and Pérez, P.V. (2022). Nuevos datos sobre el sobrepeso y la obesidad en Cuba. Nuevos datos del sobrepeso y la obesidad en Cuba. https://www.researchgate.net/publication/364715998_Nuevos_datos_sobre_el_sobrepeso_y_la_obesidad_en_Cuba [Accessed 09.01.26]

Notes:	Data from the Cuba National Health Survey 2018-2019
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

% Adults living with obesity in Cuba 1982-2019

Men




Survey type:	Measured
References:	1982, 1993: Rodriguez-Ojea A, Jimenez S, Berdasco A, Esquivel M. The nutrition transition in Cuba in the nineties:an overview. Public health Nutrition 2002:5(1A), 129-133 1998: Rodriguez-Ojea A, Jimenez S, Berdasco A and Esquivel M. (2002). The Nutrition transition in Cuba in the nineties: an overview. Public Health Nutrition, 5 (1A): 129 - 133. 2010: Nie, Peng & León, Alina & Díaz, María & Sousa-Poza, Alfonso. (2017). The rise in obesity in Cuba from 2001 to 2010: An analysis of National Survey on Risk Factors and Chronic Diseases data. Economics & Human Biology. 28. 10.1016/j.ehb.2017.11.003. 2018: Díaz, E., Maldonado, G., Medina, R.S. and Pérez, P.V. (2022). Nuevos datos sobre el sobrepeso y la obesidad en Cuba. Nuevos datos del sobrepeso y la obesidad en Cuba. https://www.researchgate.net/publication/364715998_Nuevos_datos_sobre_el_sobrepeso_y_la_obesidad_en_Cuba [Accessed

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women




Survey type:	Measured
References:	1982, 1993: Rodriguez-Ojea A, Jimenez S, Berdasco A, Esquivel M. The nutrition transition in Cuba in the nineties:an overview. Public health Nutrition 2002:5(1A), 129-133 1998: Rodriguez-Ojea A, Jimenez S, Berdasco A and Esquivel M. (2002). The Nutrition transition in Cuba in the nineties: an overview. Public Health Nutrition, 5 (1A): 129 - 133. 2010: Nie, Peng & León, Alina & Díaz, María & Sousa-Poza, Alfonso. (2017). The rise in obesity in Cuba from 2001 to 2010: An analysis of National Survey on Risk Factors and Chronic Diseases data. Economics & Human Biology. 28. 10.1016/j.ehb.2017.11.003. 2018: Díaz, E., Maldonado, G., Medina, R.S. and Pérez, P.V. (2022). Nuevos datos sobre el sobrepeso y la obesidad en Cuba. Nuevos datos del sobrepeso y la obesidad en Cuba. https://www.researchgate.net/publication/364715998_Nuevos_datos_sobre_el_sobrepeso_y_la_obesidad_en_Cuba [Accessed

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with overweight or obesity in Cuba 1982-2019

Men




Survey type:	Measured
References:	1982, 1993: Rodriguez-Ojea A, Jimenez S, Berdasco A, Esquivel M. The nutrition transition in Cuba in the nineties:an overview. Public health Nutrition 2002:5(1A), 129-133 1998: Rodriguez-Ojea A, Jimenez S, Berdasco A and Esquivel M. (2002). The Nutrition transition in Cuba in the nineties: an overview. Public Health Nutrition, 5 (1A): 129 - 133. 2010: Nie, Peng & León, Alina & Díaz, María & Sousa-Poza, Alfonso. (2017). The rise in obesity in Cuba from 2001 to 2010: An analysis of National Survey on Risk Factors and Chronic Diseases data. Economics & Human Biology. 28. 10.1016/j.ehb.2017.11.003. 2018: Díaz, E., Maldonado, G., Medina, R.S. and Pérez, P.V. (2022). Nuevos datos sobre el sobrepeso y la obesidad en Cuba. Nuevos datos del sobrepeso y la obesidad en Cuba. https://www.researchgate.net/publication/364715998_Nuevos_datos_sobre_el_sobrepeso_y_la_obesidad_en_Cuba [Accessed

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women




Survey type:	Measured
References:	1982, 1993: Rodriguez-Ojea A, Jimenez S, Berdasco A, Esquivel M. The nutrition transition in Cuba in the nineties:an overview. Public health Nutrition 2002:5(1A), 129-133 1998: Rodriguez-Ojea A, Jimenez S, Berdasco A and Esquivel M. (2002). The Nutrition transition in Cuba in the nineties: an overview. Public Health Nutrition, 5 (1A): 129 - 133. 2010: Nie, Peng & León, Alina & Díaz, María & Sousa-Poza, Alfonso. (2017). The rise in obesity in Cuba from 2001 to 2010: An analysis of National Survey on Risk Factors and Chronic Diseases data. Economics & Human Biology. 28. 10.1016/j.ehb.2017.11.003. 2018: Díaz, E., Maldonado, G., Medina, R.S. and Pérez, P.V. (2022). Nuevos datos sobre el sobrepeso y la obesidad en Cuba. Nuevos datos del sobrepeso y la obesidad en Cuba. https://www.researchgate.net/publication/364715998_Nuevos_datos_sobre_el_sobrepeso_y_la_obesidad_en_Cuba [Accessed

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with obesity in selected countries worldwide 1976-2018, selected countries

Men



References:

1960, 1971, 1973, 1976, 1991: Flegal KM, Carroll MD, Kuczmarski RJ, Johnson CL. Overweight and obesity in the United States: prevalence and trends, 1960-1994. International Journal of Obesity (1998);22:39-47

1975: Monteiro CA, Conde WL, Popking BM. Is obesity replacing or adding to undernutrition? Evidence from different social classes in Brazil. 2002. Public Health Nutrition:51(1A), 105-112

1981: Royal College of Physicians (1983). Obesity. Reprinted from the Journal of the Royal College of Physicians of London Vol 17 (No 1) January 1983


1982: Rodriguez-Ojea A, Jimenez S, Berdasco A, Esquivel M. The nutrition transition in Cuba in the nineties:an overview. Public health Nutrition 2002:5(1A), 129-133

1985: Berg C, Rosengren A, Aires N, :appas G, Toren K, Thelle D, Lissner L. Trends in overweight and obesity from 1985 to 2002 in Goteborg, West Sweden. IJO 2005 Aug;29(8):916-24

1986: Yoshiike N, Seino F, Tajima S, Arai Y, Kawano M, Furuhata T, Inoue S. Twenty-year changes in the prevalence of overweight in

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women



References:

1960, 1971, 1973, 1976, 1991: Flegal KM, Carroll MD, Kuczmarski RJ, Johnson CL. Overweight and obesity in the United States: prevalence and trends, 1960-1994. International Journal of Obesity (1998);22:39-47

1975: Monteiro CA, Conde WL, Popking BM. Is obesity replacing or adding to undernutrition? Evidence from different social classes in Brazil. 2002. Public Health Nutrition:51(1A), 105-112

1981: Royal College of Physicians (1983). Obesity. Reprinted from the Journal of the Royal College of Physicians of London Vol 17 (No 1) January 1983

1982: Rodriguez-Ojea A, Jimenez S, Berdasco A, Esquivel M. The nutrition transition in Cuba in the nineties:an overview. Public health Nutrition 2002:5(1A), 129-133


1985: Berg C, Rosengren A, Aires N, :appas G, Toren K, Thelle D, Lissner L. Trends in overweight and obesity from 1985 to 2002 in Goteborg, West Sweden. IJO 2005 Aug;29(8):916-24

1986: Yoshiike N, Seino F, Tajima S, Arai Y, Kawano M, Furuhata T, Inoue S. Twenty-year changes in the prevalence of overweight in

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Overweight/obesity by region

Adults, 2018-2019



Survey type:	Measured
Age:	15+
Sample size:	14,339
Area covered:	National


References: Díaz, E., Maldonado, G., Medina, R.S. and Pérez, P.V. (2022). Nuevos datos sobre el sobrepeso y la obesidad en Cuba. Nuevos datos del sobrepeso y la obesidad en Cuba. https://www.researchgate.net/publication/364715998_Nuevos_datos_sobre_el_sobrepeso_y_la_obesidad_en_Cuba [Accessed 09.01.26]

Notes: Data from the Cuba National Health Survey 2018-2019

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Double burden of underweight & obesity

Adults, 2022




Survey type:	Measured
Age:	20+
References:	NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population representative studies with 222 million children, adolescents, and adults. Lancet 2024; published online Feb 29. https://doi.org/10.1016/S0140-6736(23)02750-2.
Notes:	Age Standardised estimates

Definitions:	Combined prevalence of BMI<18.5 kg/m² and BMI>=30 kg/m² (double burden of underweight and obesity)

Insufficient physical activity


Adults, 2022



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.


Men, 2022



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.

Women, 2022




Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.


Mental health - depression disorders

Adults, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)

Men, 2023



Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)


Women, 2023



Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)


Mental health - anxiety disorders

Adults, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)

Men, 2023



Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)


Women, 2023



Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)


Oesophageal cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Breast cancer


Women, 2022



Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Colorectal cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Pancreatic cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Gallbladder cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Kidney cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Cancer of the uterus


Women, 2022



Age:	20+
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Raised blood pressure

Adults, 2019




Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.

Men, 2019



Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.


Women, 2019



Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.


Raised cholesterol

Adults, 2008




References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).

Men, 2008



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).


Women, 2008



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).


Raised fasting blood glucose

Men, 2014



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A869?lang=en
Definitions:	Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication).


Women, 2014



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A869?lang=en
Definitions:	Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication).

Diabetes prevalence


Adults, 2024



References: Reproduced with kind permission International Diabetes Federation. IDF Diabetes Atlas, 11th edn. Brussels, Belgium:International Diabetes Federation, 2025. http://www.diabetesatlas.org

Ovarian Cancer


Women, 2022



Age:	20+
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Leukemia


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Liver and intrahepatic bile duct Cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Multiple Myeloma


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Non Hodgkin Lymphoma


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Thyroid Cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

PDF created on November 2, 2026