
India



Country report card - adults


This report card contains the latest data available on the Global Obesity Observatory on overweight and obesity for adults. Where available, data on common and relevant obesity drivers and comorbidities are also presented.

View the latest version of this report on the Global Obesity Observatory at https://data.worldobesity.org/country/india-95/.

Contents	Page
Obesity prevalence	3
Trend: % Adults living with obesity in India 2005-2019	6
Trend: % Adults living with overweight or obesity in India, 2005-2023	11
Overweight/obesity by education	16
Overweight/obesity by age	21
Overweight/obesity by region	23
Overweight/obesity by socio-economic group	28
Double burden of underweight & obesity	33
Insufficient physical activity	36
Average weekly frequency of fast food consumption	43
Mental health - depression disorders	50
Mental health - anxiety disorders	54
Percent of population who cannot afford a healthy diet	58
Oesophageal cancer	59
Breast cancer	62
Colorectal cancer	64
Pancreatic cancer	67
Gallbladder cancer	70
Kidney cancer	73
Cancer of the uterus	76
Raised blood pressure	78
Raised cholesterol	82
Raised fasting blood glucose	85
Diabetes prevalence	87
Ovarian Cancer	88
Leukemia	90
Liver and intrahepatic bile duct Cancer	95
Multiple Myeloma	98
Non Hodgkin Lymphoma	103
Thyroid Cancer	108

Obesity prevalence

Adults, 2023-2024




Survey type:	Measured
Age:	15-49
Sample size:	817,734
Area covered:	National
References:	International Institute for Population Sciences (IIPS) 2026. National Family Health Survey (NFHS-6), 2023-24: India and State/UT Fact Sheets. Mumbai: IIPS. Available at: https://www.nfhsiips.in/nfhsuser/assets/National%20Family%20Health%20Survey%20%28NFHS-6%29%202023-2024%20Fact%20Sheets.pdf [Accessed: 29.05.26]

Notes:	716,397 women and 100,977 men.
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

% Adults living with obesity in India 2005-2019


Men



Survey type:	Measured
References:	2005: International Institute for Population Sciences - IIPS/India and Macro International. 2007. National Family Health Survey NFHS-3, 2005–06: India: Volume I. Mumbai, India: IIPS and Macro International. 2015: International Institute for Population Sciences - IIPS/India and ICF. 2017. National Family Health Survey NFHS-4, 2015-16: India. Mumbai: IIPS. Available at: https://dhsprogram.com/pubs/pdf/FR339/FR339.pdf (Accessed 6 March 2019). 2019: International Institute for Population Sciences (IIPS) and ICF. 2021. National Family Health Survey (NFHS-5), 2019-21. Available at: https://dhsprogram.com/pubs/pdf/FR375/FR375.pdf [Accessed 24.07.24]

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women




Survey type:	Measured
References:	2005: International Institute for Population Sciences - IIPS/India and Macro International. 2007. National Family Health Survey NFHS-3, 2005–06: India: Volume I. Mumbai, India: IIPS and Macro International. 2015: International Institute for Population Sciences - IIPS/India and ICF. 2017. National Family Health Survey NFHS-4, 2015-16: India. Mumbai: IIPS. Available at: https://dhsprogram.com/pubs/pdf/FR339/FR339.pdf (Accessed 6 March 2019). 2019: International Institute for Population Sciences (IIPS) and ICF. 2021. National Family Health Survey (NFHS-5), 2019-21. Available at: https://dhsprogram.com/pubs/pdf/FR375/FR375.pdf [Accessed 24.07.24]

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with overweight or obesity in India, 2005-2023


Men



Survey type:	Measured
References:	2005: International Institute for Population Sciences - IIPS/India and Macro International. 2007. National Family Health Survey NFHS-3, 2005–06: India: Volume I. Mumbai, India: IIPS and Macro International. 2015: International Institute for Population Sciences - IIPS/India and ICF. 2017. National Family Health Survey NFHS-4, 2015-16: India. Mumbai: IIPS. Available at: https://dhsprogram.com/pubs/pdf/FR339/FR339.pdf (Accessed 6 March 2019). 2019: International Institute for Population Sciences (IIPS) and ICF. 2021. National Family Health Survey (NFHS-5), 2019-21. Available at: https://dhsprogram.com/pubs/pdf/FR375/FR375.pdf [Accessed 24.07.24] 2023: International Institute for Population Sciences (IIPS) 2026. National Family Health Survey (NFHS-6), 2023-24: India and State/UT Fact Sheets. Mumbai: IIPS. Available at: https://www.nfhsiips.in/nfhsuser/assets/National%20Family%20Health%20Survey%20%28NFHS-6%29%202023-

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women




Survey type:	Measured
References:	2005: International Institute for Population Sciences - IIPS/India and Macro International. 2007. National Family Health Survey NFHS-3, 2005–06: India: Volume I. Mumbai, India: IIPS and Macro International. 2015: International Institute for Population Sciences - IIPS/India and ICF. 2017. National Family Health Survey NFHS-4, 2015-16: India. Mumbai: IIPS. Available at: https://dhsprogram.com/pubs/pdf/FR339/FR339.pdf (Accessed 6 March 2019). 2019: International Institute for Population Sciences (IIPS) and ICF. 2021. National Family Health Survey (NFHS-5), 2019-21. Available at: https://dhsprogram.com/pubs/pdf/FR375/FR375.pdf [Accessed 24.07.24] 2023: International Institute for Population Sciences (IIPS) 2026. National Family Health Survey (NFHS-6), 2023-24: India and State/UT Fact Sheets. Mumbai: IIPS. Available at: https://www.nfhsiips.in/nfhsuser/assets/National%20Family%20Health%20Survey%20%28NFHS-6%29%202023-

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Overweight/obesity by education


Men, 2019-2021



Survey type:	Measured
Age:	15-49
Sample size:	752983
Area covered:	National
References:	International Institute for Population Sciences (IIPS) and ICF. 2021. National Family Health Survey (NFHS-5), 2019-21 India: Volume I. Mumbai: IIPS. Available at https://dhsprogram.com/pubs/pdf/FR375/FR375.pdf (last accessed 05.09.24)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2019-2021




Survey type:	Measured
Age:	15-49
Sample size:	752983
Area covered:	National
References:	International Institute for Population Sciences (IIPS) and ICF. 2021. National Family Health Survey (NFHS-5), 2019-21 India: Volume I. Mumbai: IIPS. Available at https://dhsprogram.com/pubs/pdf/FR375/FR375.pdf (last accessed 05.09.24)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Overweight/obesity by age

Adults, 2019-2021




Survey type:	Measured
Sample size:	752983
Area covered:	National
References:	International Institute for Population Sciences (IIPS) and ICF. 2021. National Family Health Survey (NFHS-5), 2019-21 India: Volume I. Mumbai: IIPS. Available at https://dhsprogram.com/pubs/pdf/FR375/FR375.pdf (last accessed 05.09.24)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Overweight/obesity by region


Men, 2023-2024



Survey type:	Measured
Age:	15-49
Sample size:	817,734
Area covered:	National

References:	International Institute for Population Sciences (IIPS) 2026. National Family Health Survey (NFHS-6), 2023-24: India and State/UT Fact Sheets. Mumbai: IIPS. Available at: https://www.nfhsiips.in/nfhsuser/assets/National%20Family%20Health%20Survey%20%28NFHS-6%29%202023-2024%20Fact%20Sheets.pdf [Accessed: 29.05.26]
Notes:	716,397 women and 100,977 men.
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2023-2024




Survey type:	Measured
Age:	15-49
Sample size:	817,734
Area covered:	National

References:	International Institute for Population Sciences (IIPS) 2026. National Family Health Survey (NFHS-6), 2023-24: India and State/UT Fact Sheets. Mumbai: IIPS. Available at: https://www.nfhsiips.in/nfhsuser/assets/National%20Family%20Health%20Survey%20%28NFHS-6%29%202023-2024%20Fact%20Sheets.pdf [Accessed: 29.05.26]
Notes:	716,397 women and 100,977 men.
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Overweight/obesity by socio-economic group


Men, 2019-2021



Survey type:	Measured
Age:	15-49
Sample size:	752983
Area covered:	National
References:	International Institute for Population Sciences (IIPS) and ICF. 2021. National Family Health Survey (NFHS-5), 2019-21 India: Volume I. Mumbai: IIPS. Available at https://dhsprogram.com/pubs/pdf/FR375/FR375.pdf (last accessed 05.09.24)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2019-2021




Survey type:	Measured
Age:	15-49
Sample size:	752983
Area covered:	National
References:	International Institute for Population Sciences (IIPS) and ICF. 2021. National Family Health Survey (NFHS-5), 2019-21 India: Volume I. Mumbai: IIPS. Available at https://dhsprogram.com/pubs/pdf/FR375/FR375.pdf (last accessed 05.09.24)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Double burden of underweight & obesity

Adults, 2022




Survey type:	Measured
Age:	20+
References:	NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population representative studies with 222 million children, adolescents, and adults. Lancet 2024; published online Feb 29. https://doi.org/10.1016/S0140-6736(23)02750-2.
Notes:	Age Standardised estimates

Definitions:	Combined prevalence of BMI<18.5 kg/m² and BMI>=30 kg/m² (double burden of underweight and obesity)

Insufficient physical activity


Adults, 2022



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.


Men, 2022



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.

Women, 2022




Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.

Average weekly frequency of fast food consumption


Adults, 2021-2024



Survey type:	Self-reported
Age:	15+
Sample size:	2997
Area covered:	National
References:	Global Diet Quality Project. 2024. "DQQ Results Dataset 2021-2024." Harvard Dataverse. https://doi.org/doi:10.7910/DVN/KY3W8A Licence: CC BY-NC-SA 4.0. Accessed at dietquality.org on October 15, 2025.

Notes:	For more global dietary survey data, visit the Global Diet Quality Project website at https://www.dietquality.org/
Definitions:	Proportion of the population who consumed fast food during the previous day or night (self-reported)


Men, 2021-2024



Survey type:	Self-reported
Age:	15+
Sample size:	2997
Area covered:	National
References:	Global Diet Quality Project. 2024. "DQQ Results Dataset 2021-2024." Harvard Dataverse. https://doi.org/doi:10.7910/DVN/KY3W8A Licence: CC BY-NC-SA 4.0. Accessed at dietquality.org on October 15, 2025.

Notes:	For more global dietary survey data, visit the Global Diet Quality Project website at https://www.dietquality.org/
Definitions:	Proportion of the population who consumed fast food during the previous day or night (self-reported)

Women, 2021-2024




Survey type:	Self-reported
Age:	15+
Sample size:	2997
Area covered:	National
References:	Global Diet Quality Project. 2024. "DQQ Results Dataset 2021-2024." Harvard Dataverse. https://doi.org/doi:10.7910/DVN/KY3W8A Licence: CC BY-NC-SA 4.0. Accessed at dietquality.org on October 15, 2025.

Notes:	For more global dietary survey data, visit the Global Diet Quality Project website at https://www.dietquality.org/
Definitions:	Proportion of the population who consumed fast food during the previous day or night (self-reported)


Mental health - depression disorders

Adults, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)

Men, 2023



Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)


Women, 2023



Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)


Mental health - anxiety disorders

Adults, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)

Men, 2023



Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)


Women, 2023



Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)

Percent of population who cannot afford a healthy diet


Adults, 2025



Area covered:	National
References:	FAO. 2026. FAOSTAT: Cost and Affordability of a Healthy Diet (CoAHD). https://www.fao.org/faostat/en/#data/CAHD. Licence: CC-BY-4.0. [Last Accessed 12.10.26]
Notes:	The percent of the population whose food budget is below the cost of a healthy diet.


Oesophageal cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Breast cancer


Women, 2022



Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Colorectal cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Pancreatic cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Gallbladder cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Kidney cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Cancer of the uterus


Women, 2022



Age:	20+
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Raised blood pressure

Adults, 2019




Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.

Men, 2019



Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.


Women, 2019



Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.


Raised cholesterol

Adults, 2008




References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).

Men, 2008



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).


Women, 2008



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).


Raised fasting blood glucose

Men, 2014



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A869?lang=en
Definitions:	Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication).


Women, 2014



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A869?lang=en
Definitions:	Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication).

Diabetes prevalence


Adults, 2024



References: Reproduced with kind permission International Diabetes Federation. IDF Diabetes Atlas, 11th edn. Brussels, Belgium:International Diabetes Federation, 2025. http://www.diabetesatlas.org

Ovarian Cancer


Women, 2022



Age:	20+
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Leukemia


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Liver and intrahepatic bile duct Cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Multiple Myeloma


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Non Hodgkin Lymphoma


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Thyroid Cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

PDF created on November 7, 2026