Ireland



Country report card

This report card contains the latest data available on the Global Obesity Observatory on overweight and obesity . Where available, data on common and relevant obesity drivers and comorbidities are also presented.


View the latest version of this report on the Global Obesity Observatory at https://data.worldobesity.org/country/ireland-99/.

	Page
Obesity prevalence	4
Trend: % Adults living with obesity in Ireland (Measured) 1990-2022	11
Trend: % Adults living with obesity in Ireland (Self-reported), 2000-2022	16
Trend: % Adults living with overweight or obesity in Ireland (Measured) 1990-2008	23
Trend: % Adults living with overweight or obesity in Ireland (Self-reported), 2000-2022	28
Trend: % Children (age 6-7) living with overweight or obesity in Ireland 2008-2022	35
Trend: % Children (aged 7) living with obesity in Ireland, 2007-2022 (WHO)	37
Trend: % Children (aged 7) living with overweight or obesity in Ireland, 2007-2022 (WHO)	42
Trend: % Children living with overweight or obesity by urbanisation in Ireland 2008-2022	47
Trend: % Adults living with obesity in Europe 1976-2018, selected countries	49
Overweight/obesity by age and education	54
Overweight/obesity by education	57
Overweight/obesity by age	66
Overweight/obesity by region	70
Overweight/obesity by age and region	75
Overweight/obesity by age and socio-economic group	79
Overweight/obesity by socio-economic group	83
Overweight/obesity by age and limited activity	88
Double burden of underweight & obesity	92
Insufficient physical activity	99
Sugar consumption	112
Estimated per capita sugar sweetened beverages intake	113
Prevalence of at least daily soft drink consumption	114
Prevalence of confectionery consumption	121
Prevalence of sweet/savoury snack consumption	122
Prevalence of less than daily fruit consumption	123
Prevalence of less than daily vegetable consumption	130
Mental health - depression disorders	137
Mental health - anxiety disorders	144
Percent of population who cannot afford a healthy diet	151
Oesophageal cancer	152
Breast cancer	155
Colorectal cancer	157
Pancreatic cancer	160
Gallbladder cancer	163
Kidney cancer	166
Cancer of the uterus	169
Raised blood pressure	171
Raised cholesterol	175
Raised fasting blood glucose	178
Diabetes prevalence	180
Ovarian Cancer	181
Leukemia	183

Contents	Page
Liver and intrahepatic bile duct Cancer	%%
Multiple Myeloma	%%
Non Hodgkin Lymphoma	%%
Thyroid Cancer	%%

Obesity prevalence


Adults, 2023-2024



Survey type:	Self-reported
Age:	15+
Sample size:	7398
Area covered:	National
References:	Healthy Ireland Survey 2024: Summary Report. Available at: https://www.gov.ie/en/healthy-ireland/publications/healthy-ireland-survey-2024/ (Accessed: 13.06.25)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Children, 2022-2024



Survey type:	Measured
Age:	7
Sample size:	1285
Area covered:	National
References:	WHO European Childhood Obesity Surveillance Initiative (COSI): report on the sixth round of data collection, 2022– 2024. Copenhagen: WHO Regional Office for Europe; 2025. Licence: CC BY-NC-SA 3.0 IGO.

Cutoffs:	WHO 2007

0-5 years, 2010-2011




Survey type:	Measured
Age:	1-5
Sample size:	494
Area covered:	National
References:	Ireland 2010-2011 National Pre-School Nutrition Survey
Notes:	UNICEF/WHO/World Bank Joint Child Malnutrition Estimates Expanded Database: Overweight estimates: National and Disaggregated, July 2025, New York. Weighted sample size. For more information, please consult https://data.unicef.org/resources/jme/. NB. Age interval 12-59 months: unadjusted

Definitions:	Percentage of children under 5 years of age falling above 2 standard deviations (moderate and severe) from the median weight-for-height of the reference population.
Cutoffs:	=>+2SD

% Adults living with obesity in Ireland (Measured) 1990-2022


Men



Survey type:	Measured
References:	1990: North/South Ireland Food Consumption Survey, 2000. Irish Universities Nutrition Alliance Summary Report 1997: McCarthy SN, Harrington KE, Kiely M et al. Analyses of the anthropometric data from the North/South Ireland Food Consumption Survey. Public Health Nutrition 2001;4 (5(A):1099-1106 2008: National Adult Nutrition Survey, Irish Universities Nutrition Alliance, Published 2011 http://www.iuna.net/wp-content/uploads/2010/12/National-Adult-Nutrition-Survey-Summary-Report-March-2011.pdf 2021: Irish Universities Nutrition Alliance (IUNA). National Adult Nutrition Survey (NANS) II: Main Report. https://www.iuna.net/surveyreports [Accessed 06.08.25]

Notes:	Adults aged 18/19-64
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².
	Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women




Survey type:	Measured
References:	1990: North/South Ireland Food Consumption Survey, 2000. Irish Universities Nutrition Alliance Summary Report 1997: McCarthy SN, Harrington KE, Kiely M et al. Analyses of the anthropometric data from the North/South Ireland Food Consumption Survey. Public Health Nutrition 2001;4 (5(A):1099-1106 2008: National Adult Nutrition Survey, Irish Universities Nutrition Alliance, Published 2011 http://www.iuna.net/wp-content/uploads/2010/12/National-Adult-Nutrition-Survey-Summary-Report-March-2011.pdf 2021: Irish Universities Nutrition Alliance (IUNA). National Adult Nutrition Survey (NANS) II: Main Report. https://www.iuna.net/surveyreports [Accessed 06.08.25]

Notes:	Adults aged 18/19-64
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².
	Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with obesity in Ireland (Self-reported), 2000-2022


Men



Survey type:	Self-reported
References:	2000: Eurostat Yearbook 2006/07. A goldmine of statistical information. Available at https://ec.europa.eu/eurostat/documents/2995521/5059290/1-20022007-BP-EN.PDF.pdf/edab8c31-b9f3-4c8e-b4db-4137bd045efa?t=1414683510000 (last accessed 04.11.21) 2014: Eurostat Database:http://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=hlth_ehis_bm1e&lang=en (last accessed 25.08.20) 2019: Eurostat 2019. Data available at https://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=hlth_ehis_bm1i&lang=en (last accessed 09.08.21) 2022: Eurostat 2022 Data. Available at https://ec.europa.eu/eurostat/databrowser/view/ilc_hch10/default/table?lang=en (last accessed 19.03.25)

Notes:	Eurostat
Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².	
Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.	


Women



Survey type:	Self-reported
References:	2000: Eurostat Yearbook 2006/07. A goldmine of statistical information. Available at https://ec.europa.eu/eurostat/documents/2995521/5059290/1-20022007-BP-EN.PDF.pdf/edab8c31-b9f3-4c8e-b4db-4137bd045efa?t=1414683510000 (last accessed 04.11.21) 2014: Eurostat Database:http://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=hlth_ehis_bm1e&lang=en (last accessed 25.08.20) 2019: Eurostat 2019. Data available at https://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=hlth_ehis_bm1i&lang=en (last accessed 09.08.21) 2022: Eurostat 2022 Data. Available at https://ec.europa.eu/eurostat/databrowser/view/ilc_hch10/default/table?lang=en (last accessed 19.03.25)

Notes:	Eurostat
Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².	
Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.	

Men and women




Survey type:	Self-reported
References:	2000: Eurostat Yearbook 2006/07. A goldmine of statistical information. Available at https://ec.europa.eu/eurostat/documents/2995521/5059290/1-20022007-BP-EN.PDF.pdf/edab8c31-b9f3-4c8e-b4db-4137bd045efa?t=1414683510000 (last accessed 04.11.21) 2014: Eurostat Database:http://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=hlth_ehis_bm1e&lang=en (last accessed 25.08.20) 2019: Eurostat 2019. Data available at https://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=hlth_ehis_bm1i&lang=en (last accessed 09.08.21) 2022: Eurostat 2022 Data. Available at https://ec.europa.eu/eurostat/databrowser/view/ilc_hch10/default/table?lang=en (last accessed 19.03.25)

Notes:	Eurostat
Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².	
Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.	

% Adults living with overweight or obesity in Ireland (Measured) 1990-2008


Men



Survey type:	Measured
References:	1990: North/South Ireland Food Consumption Survey, 2000. Irish Universities Nutrition Alliance Summary Report 1997: McCarthy SN, Harrington KE, Kiely M et al. Analyses of the anthropometric data from the North/South Ireland Food Consumption Survey. Public Health Nutrition 2001;4 (5(A):1099-1106 2008: National Adult Nutrition Survey, Irish Universities Nutrition Alliance, Published 2011 http://www.iuna.net/wp-content/uploads/2010/12/National-Adult-Nutrition-Survey-Summary-Report-March-2011.pdf 2021: Irish Universities Nutrition Alliance (IUNA). National Adult Nutrition Survey (NANS) II: Main Report. https://www.iuna.net/surveyreports [Accessed 06.08.25]

Notes:	Adults aged 18/19-64
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².
	Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women




Survey type:	Measured
References:	1990: North/South Ireland Food Consumption Survey, 2000. Irish Universities Nutrition Alliance Summary Report 1997: McCarthy SN, Harrington KE, Kiely M et al. Analyses of the anthropometric data from the North/South Ireland Food Consumption Survey. Public Health Nutrition 2001;4 (5(A):1099-1106 2008: National Adult Nutrition Survey, Irish Universities Nutrition Alliance, Published 2011 http://www.iuna.net/wp-content/uploads/2010/12/National-Adult-Nutrition-Survey-Summary-Report-March-2011.pdf 2021: Irish Universities Nutrition Alliance (IUNA). National Adult Nutrition Survey (NANS) II: Main Report. https://www.iuna.net/surveyreports [Accessed 06.08.25]

Notes:	Adults aged 18/19-64
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².
	Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with overweight or obesity in Ireland (Self-reported), 2000-2022


Men



Survey type:	Self-reported
References:	Eurostat Yearbook 2006/07. A goldmine of statistical information. Available at https://ec.europa.eu/eurostat/documents/2995521/5059290/1-20022007-BP-EN.PDF.pdf/edab8c31-b9f3-4c8e-b4db-4137bd045efa?t=1414683510000 (last accessed 04.11.21)
Notes:	Eurostat
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.


Women



Survey type:	Self-reported
References:	Eurostat Yearbook 2006/07. A goldmine of statistical information. Available at https://ec.europa.eu/eurostat/documents/2995521/5059290/1-20022007-BP-EN.PDF.pdf/edab8c31-b9f3-4c8e-b4db-4137bd045efa?t=1414683510000 (last accessed 04.11.21)
Notes:	Eurostat
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.


Men and women



Survey type:	Self-reported
References:	Eurostat Yearbook 2006/07. A goldmine of statistical information. Available at https://ec.europa.eu/eurostat/documents/2995521/5059290/1-20022007-BP-EN.PDF.pdf/edab8c31-b9f3-4c8e-b4db-4137bd045efa?t=1414683510000 (last accessed 04.11.21)
Notes:	Eurostat
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Children (age 6-7) living with overweight or obesity in Ireland 2008-2022




Survey type:	Measured
References:	Bel-Serrat S, Heinen MM, Murrin CM, Daly L, Mehegan J, Concannon M, Flood C, Farrell D, O'Brien S, Eldin N, Kelleher CC [2017]. The Childhood Obesity Surveillance Initiative (COSI) in the Republic of Ireland: Findings from 2008, 2010, 2012 and 2015. Dublin: Health Service Executive.
Notes:	Aged 6-7. IOTF International Cut off point

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Children (aged 7) living with obesity in Ireland, 2007-2022 (WHO)

Boys




Survey type:	Measured
References:	2007: Wijnhoven, T.M.A., van Raaij, J.M.A., Spinelli, A., Rito, A.I., Hovengen, R., Kunesova, M., Starc, G., Rutter, H., Sjoberg, A., Petrauskiene, A., O'Dwyer, U., Petrova, S., Farrugia Sant'Angelo, V., Wauters, M., Yngve, A., Rubana, I.-M. and Breda, J. (2013), WHO-COSI 2008: weight, height and BMI. Pediatric Obesity, 8: 79-97. https://doi.org/10.1111/j.2047-6310.2012.00090.x 2009: Wijnhoven, T.M., van Raaij, J.M., Spinelli, A. et al. WHO European Childhood Obesity Surveillance Initiative: body mass index and level of overweight among 6-9-year-old children from school year 2007/2008 to school year 2009/2010. BMC Public Health 14, 806 (2014). https://doi.org/10.1186/1471-2458-14-806 2012: World Health Organization. Regional Office for Europe. 2018. WHO European Childhood Obesity Surveillance Initiative: overweight and obesity among 6-9-year-old children: report of the third round of data collection 2012-2013.World Health Organization. Regional Office for Europe. https://apps.who.int/iris/handle/10665/346108

Notes:	COSI
Definitions:	WHO 2007 cutoffs

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Girls




Survey type:	Measured
References:	2007: Wijnhoven, T.M.A., van Raaij, J.M.A., Spinelli, A., Rito, A.I., Hovengen, R., Kunesova, M., Starc, G., Rutter, H., Sjoberg, A., Petrauskiene, A., O'Dwyer, U., Petrova, S., Farrugia Sant'Angelo, V., Wauters, M., Yngve, A., Rubana, I.-M. and Breda, J. (2013), WHO-COSI 2008: weight, height and BMI. Pediatric Obesity, 8: 79-97. https://doi.org/10.1111/j.2047-6310.2012.00090.x 2009: Wijnhoven, T.M., van Raaij, J.M., Spinelli, A. et al. WHO European Childhood Obesity Surveillance Initiative: body mass index and level of overweight among 6-9-year-old children from school year 2007/2008 to school year 2009/2010. BMC Public Health 14, 806 (2014). https://doi.org/10.1186/1471-2458-14-806 2012: World Health Organization. Regional Office for Europe. 2018. WHO European Childhood Obesity Surveillance Initiative: overweight and obesity among 6-9-year-old children: report of the third round of data collection 2012-2013.World Health Organization. Regional Office for Europe. https://apps.who.int/iris/handle/10665/346108

Notes:	COSI
Definitions:	WHO 2007 cutoffs

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Children (aged 7) living with overweight or obesity in Ireland, 2007-2022 (WHO)

Boys




Survey type:	Measured
References:	2007: Wijnhoven, T.M.A., van Raaij, J.M.A., Spinelli, A., Rito, A.I., Hovengen, R., Kunesova, M., Starc, G., Rutter, H., Sjoberg, A., Petrauskiene, A., O'Dwyer, U., Petrova, S., Farrugia Sant'Angelo, V., Wauters, M., Yngve, A., Rubana, I.-M. and Breda, J. (2013), WHO-COSI 2008: weight, height and BMI. Pediatric Obesity, 8: 79-97. https://doi.org/10.1111/j.2047-6310.2012.00090.x 2009: Wijnhoven, T.M., van Raaij, J.M., Spinelli, A. et al. WHO European Childhood Obesity Surveillance Initiative: body mass index and level of overweight among 6-9-year-old children from school year 2007/2008 to school year 2009/2010. BMC Public Health 14, 806 (2014). https://doi.org/10.1186/1471-2458-14-806 2012: World Health Organization. Regional Office for Europe. 2018. WHO European Childhood Obesity Surveillance Initiative: overweight and obesity among 6-9-year-old children: report of the third round of data collection 2012-2013.World Health Organization. Regional Office for Europe. https://apps.who.int/iris/handle/10665/346108

Notes:	COSI
Definitions:	WHO 2007 cutoffs

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Girls




Survey type:	Measured
References:	2007: Wijnhoven, T.M.A., van Raaij, J.M.A., Spinelli, A., Rito, A.I., Hovengen, R., Kunesova, M., Starc, G., Rutter, H., Sjoberg, A., Petrauskiene, A., O'Dwyer, U., Petrova, S., Farrugia Sant'Angelo, V., Wauters, M., Yngve, A., Rubana, I.-M. and Breda, J. (2013), WHO-COSI 2008: weight, height and BMI. Pediatric Obesity, 8: 79-97. https://doi.org/10.1111/j.2047-6310.2012.00090.x 2009: Wijnhoven, T.M., van Raaij, J.M., Spinelli, A. et al. WHO European Childhood Obesity Surveillance Initiative: body mass index and level of overweight among 6-9-year-old children from school year 2007/2008 to school year 2009/2010. BMC Public Health 14, 806 (2014). https://doi.org/10.1186/1471-2458-14-806 2012: World Health Organization. Regional Office for Europe. 2018. WHO European Childhood Obesity Surveillance Initiative: overweight and obesity among 6-9-year-old children: report of the third round of data collection 2012-2013.World Health Organization. Regional Office for Europe. https://apps.who.int/iris/handle/10665/346108

Notes:	COSI
Definitions:	WHO 2007 cutoffs

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Children living with overweight or obesity by urbanisation in Ireland 2008-2022




Survey type:	Measured
References:	Bel-Serrat S, Heinen MM, Murrin CM, Daly L, Mehegan J, Concannon M, Flood C, Farrell D, O'Brien S, Eldin N, Kelleher CC [2017]. The Childhood Obesity Surveillance Initiative (COSI) in the Republic of Ireland: Findings from 2008, 2010, 2012 and 2015. Dublin: Health Service Executive.
Notes:	Aged 6-7. IOTF International Cut off point

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with obesity in Europe 1976-2018, selected countries

Men



References:

1976, 1987: Visscher TLS, Kromhout D, Seidell J. Long term and recent time trends in the prevalence of obesity among Dutch men and women. IJO 2002;26:1218-24

1981: Royal College of Physicians (1983). Obesity. Reprinted from the Journal of the Royal College of Physicians of London Vol 17 (No 1) January 1983

1982: Lahti-Koski M, Vartiainen E, Mannisto S, Pietinen P. Age, education and occupation as determinants of trends in body mass index in Finland from 1982 to 1997. International Journal of Obesity (2000);24:1669-1676


1985: Berg C, Rosengren A, Aires N, :appas G, Toren K, Thelle D, Lissner L. Trends in overweight and obesity from 1985 to 2002 in Goteborg, West Sweden. IJO 2005 Aug;29(8):916-24

1990: Berg C, Rosengren A, Aires N, :appas G, Toren K, Thelle D, Lissner L. Trends in overweight and obesity from 1985 to 2002 in Goteborg, West Sweden. IJO 2005 online published ahead of print.

1992: Lahti-Koski M, Pietinen P, Mannisto S, Vartiainen E. Trends in waist-to-hip ratio and its determinants in adults in Finland from 1987

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women



References:

1976, 1987: Visscher TLS, Kromhout D, Seidell J. Long term and recent time trends in the prevalence of obesity among Dutch men and women. IJO 2002;26:1218-24

1981: Royal College of Physicians (1983). Obesity. Reprinted from the Journal of the Royal College of Physicians of London Vol 17 (No 1) January 1983

1982: Lahti-Koski M, Vartiainen E, Mannisto S, Pietinen P. Age, education and occupation as determinants of trends in body mass index in Finland from 1982 to 1997. International Journal of Obesity (2000);24:1669-1676

1985: Berg C, Rosengren A, Aires N, :appas G, Toren K, Thelle D, Lissner L. Trends in overweight and obesity from 1985 to 2002 in Goteborg, West Sweden. IJO 2005 Aug;29(8):916-24


1990: Berg C, Rosengren A, Aires N, :appas G, Toren K, Thelle D, Lissner L. Trends in overweight and obesity from 1985 to 2002 in Goteborg, West Sweden. IJO 2005 online published ahead of print.

1992: Lahti-Koski M, Pietinen P, Mannisto S, Vartiainen E. Trends in waist-to-hip ratio and its determinants in adults in Finland from 1987

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Overweight/obesity by age and education


Men, 2017



Survey type:	Self-reported
Area covered:	National
References:	Eurostat http://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=ilc_hch10&lang=en (last accessed 25.08.20)
Notes:	Some Irish data considered to have low reliability, see EUROSTAT database for details

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2017




Survey type:	Self-reported
Area covered:	National
References:	Eurostat http://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=ilc_hch10&lang=en (last accessed 25.08.20)
Notes:	Some Irish data considered to have low reliability, see EUROSTAT database for details

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Overweight/obesity by education


Adults, 2014



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	Eurostat 2014. https://doi.org/10.2908/HLTH_EHIS_BM1E [Accessed 07.08.25]
Definitions:	Level 0-2 = Less than primary, primary and lower secondary education, Level 3-4 = Upper secondary and non-tertiary post-secondary education, Level 5-8 = Tertiary education

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Men, 2014



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	Eurostat 2014. https://doi.org/10.2908/HLTH_EHIS_BM1E [Accessed 07.08.25]
Definitions:	Level 0-2 = Less than primary, primary and lower secondary education, Level 3-4 = Upper secondary and non-tertiary post-secondary education, Level 5-8 = Tertiary education

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Women, 2014



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	Eurostat 2014. https://doi.org/10.2908/HLTH_EHIS_BM1E [Accessed 07.08.25]
Definitions:	Level 0-2 = Less than primary, primary and lower secondary education, Level 3-4 = Upper secondary and non-tertiary post-secondary education, Level 5-8 = Tertiary education

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Children, 2022-2024




Survey type:	Measured
Age:	6-9
Sample size:	1019
Area covered:	National
References:	WHO European Childhood Obesity Surveillance Initiative (COSI): report on the sixth round of data collection, 2022– 2024. Copenhagen: WHO Regional Office for Europe; 2025. Licence: CC BY-NC-SA 3.0 IGO.

Definitions:	Level of parental education: (i) low to medium, where both parents have lower education or one parent has lower education and the other has higher education; and (ii) high, where both parents have higher education. Lower education was defined as “primary school or less”, “lower secondary education”, or “upper secondary and post-secondary non-tertiary education”. Higher education included “short-cycle tertiary education or bachelor’s or equivalent level” and “master’s or doctoral or equivalent level”. In single-parent families, parental education was determined based on the educational attainment of the single parent.
Cutoffs:	WHO 2007

Overweight/obesity by age


Adults, 2023-2024



Survey type:	Self-reported
Sample size:	7308
Area covered:	National
References:	Healthy Ireland Survey 2024: Summary Report. Available at: https://www.gov.ie/en/healthy-ireland/publications/healthy-ireland-survey-2024/ (Accessed: 13.06.25)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Children, 2022-2023




Survey type:	Measured
Sample size:	5072
Area covered:	National
References:	Kilduff O, Slattery J, Lee C, O' Brien S, Murrin C, Kelleher C (2024). The Childhood Obesity Surveillance Initiative (COSI) in the Republic of Ireland - Findings from 2022 and 2023.
Notes:	Data from COSI Round 6 (2022-24). Children surveyed from 1st, 2nd, 5th and 6th classes

Cutoffs:	IOTF

Overweight/obesity by region


Adults, 2024



Survey type:	Self-reported
Age:	18+
Sample size:	5101
Area covered:	National
References:	Central Statistics Office, Ireland. 2025. Irish Health Survey 2024 - Main Results. https://www.cso.ie/en/statistics/health/irishhealthsurvey/ [Accessed 06.08.25]

Definitions:	HSE Regions
Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².	

Children, 2022-2023




Survey type:	Measured
Age:	6-12
Sample size:	5072
Area covered:	National
References:	Kilduff O, Slattery J, Lee C, O' Brien S, Murrin C, Kelleher C (2024). The Childhood Obesity Surveillance Initiative (COSI) in the Republic of Ireland - Findings from 2022 and 2023.

Notes:	Data from COSI Round 6 (2022-24). Children surveyed from 1st, 2nd, 5th and 6th classes
Cutoffs:	IOTF

Overweight/obesity by age and region


Men, 2014



Survey type:	Self-reported
Area covered:	National
References:	Eurostat http://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=hlth_ehis_bm1u&lang=en (last accessed 25.08.20)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Women, 2014



Survey type:	Self-reported
Area covered:	National
References:	Eurostat http://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=hlth_ehis_bm1u&lang=en (last accessed 25.08.20)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Children, 2022-2023




Survey type:	Measured
Age:	6-7
Sample size:	1279
Area covered:	National
References:	Kilduff O, Slattery J, Lee C, O' Brien S, Murrin C, Kelleher C (2024). The Childhood Obesity Surveillance Initiative (COSI) in the Republic of Ireland - Findings from 2022 and 2023.

Notes:	Data from COSI Round 6 (2022-24). Children surveyed from 1st, 2nd, 5th and 6th classes
Cutoffs:	IOTF


Overweight/obesity by age and socio-economic group

Adults, 2019




Survey type:	Self-reported
Area covered:	National
References:	Eurostat 2019. Available at https://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=hlth_ehis_bm1i&lang=en (last accessed 09.08.21).
Notes:	NB. Some age ranges missing due to insufficient data to display by age & income quintile
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2019



Survey type:	Self-reported
Area covered:	National
References:	Eurostat 2019. Available at https://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=hlth_ehis_bm1i&lang=en (last accessed 09.08.21).
Notes:	NB. Some age ranges missing due to insufficient data to display by age & income quintile
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Men, 2019



Survey type:	Self-reported
Area covered:	National
References:	Eurostat 2019. Available at https://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=hlth_ehis_bm1i&lang=en (last accessed 09.08.21).
Notes:	NB. Some age ranges missing due to insufficient data to display by age & income quintile
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Overweight/obesity by socio-economic group


Adults, 2019



Survey type:	Self-reported
Age:	15+
Sample size:	7621
Area covered:	National
References:	Central Statistics Office, Ireland (2019). Results of Irish Health Survey 2019. Data available at: https://data.cso.ie/#. Last accessed: 02.06.21.

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Children, 2022-2023




Survey type:	Measured
Age:	7-12
Sample size:	5072
Area covered:	National
References:	Kilduff O, Slattery J, Lee C, O' Brien S, Murrin C, Kelleher C (2024). The Childhood Obesity Surveillance Initiative (COSI) in the Republic of Ireland - Findings from 2022 and 2023.

Notes:	Data from COSI Round 6 (2022-24). Children surveyed from 1st, 2nd, 5th and 6th classes
Definitions:	Disadvantaged schools are those in the Delivering Equality of Opportunity in Schools (DEIS) programme
Cutoffs:	IOTF

Overweight/obesity by age and limited activity


Adults, 2014



Survey type:	Self-reported
Area covered:	National
References:	Eurostat 2014 available at https://appsso.eurostat.ec.europa.eu/nui/submitViewTableAction.do (last accessed 06.10.21)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Men, 2014



Survey type:	Self-reported
Area covered:	National
References:	Eurostat 2014 available at https://appsso.eurostat.ec.europa.eu/nui/submitViewTableAction.do (last accessed 06.10.21)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2014




Survey type:	Self-reported
Area covered:	National
References:	Eurostat 2014 available at https://appsso.eurostat.ec.europa.eu/nui/submitViewTableAction.do (last accessed 06.10.21)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Double burden of underweight & obesity


Adults, 2022



Survey type:	Measured
Age:	20+
References:	NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population representative studies with 222 million children, adolescents, and adults. Lancet 2024; published online Feb 29. https://doi.org/10.1016/S0140-6736(23)02750-2.
Notes:	Age Standardised estimates

Definitions:	Combined prevalence of BMI<18.5 kg/m² and BMI>=30 kg/m² (double burden of underweight and obesity)


Children, 2022



Survey type:	Measured
Age:	5-19
References:	NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population representative studies with 222 million children, adolescents, and adults. Lancet 2024; published online Feb 29. https://doi.org/10.1016/S0140-6736(23)02750-2.
Notes:	Age standardised estimates

Definitions:	Combined prevalence of BMI < -2SD and BMI > 2SD (double burden of thinness and obesity)
Cutoffs:	BMI < -2SD and BMI > 2SD

0-5 years, 2010-2011




Survey type:	Measured
Age:	0-5
References:	United Nations Children's Fund, Division of Data, Analysis, Planning and Monitoring (2025). Global UNICEF Global Databases: Overlapping Stunting, Wasting and Overweight (Survey Estimates), 2025, New York. For more information, visit: https://data.unicef.org/topic/nutrition/malnutrition/ [Accessed 06.08.25]
Notes:	Ireland 2010-2011 National Pre-School Nutrition Survey

Definitions:	Combined percentage of children under 5 years of age who are either wasted, stunted, or overweight (falling below -2 standard deviations from the median weight-for-height, falling below -2 standard deviations from the median height for age, or falling at or above +2 standard deviations from the median weight-for-height of the reference population)
Cutoffs:	WHZ <-2, HAZ <-2 and WHZ >+2

Insufficient physical activity


Adults, 2022



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.


Men, 2022



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.


Women, 2022



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.


Children, 2022-2024



Survey type:	Self-reported
Age:	6-9
Sample size:	1019
Area covered:	National
References:	WHO European Childhood Obesity Surveillance Initiative (COSI): report on the sixth round of data collection, 2022– 2024. Copenhagen: WHO Regional Office for Europe; 2025. Licence: CC BY-NC-SA 3.0 IGO.

Notes:	National data excluding Germany (Bremen only) and Kazakhstan (Almaty only).
Definitions:	Percentage of children not meeting WHO recommendations for physical activity, i.e. not playing actively/vigorously for at least one hour a day on an average day, as reported by their parent


Boys, 2022-2024



Survey type:	Self-reported
Age:	6-9
Sample size:	1019
Area covered:	National
References:	WHO European Childhood Obesity Surveillance Initiative (COSI): report on the sixth round of data collection, 2022– 2024. Copenhagen: WHO Regional Office for Europe; 2025. Licence: CC BY-NC-SA 3.0 IGO.

Notes:	National data excluding Germany (Bremen only) and Kazakhstan (Almaty only).
Definitions:	Percentage of children not meeting WHO recommendations for physical activity, i.e. not playing actively/vigorously for at least one hour a day on an average day, as reported by their parent

Girls, 2022-2024




Survey type:	Self-reported
Age:	6-9
Sample size:	1019
Area covered:	National
References:	WHO European Childhood Obesity Surveillance Initiative (COSI): report on the sixth round of data collection, 2022– 2024. Copenhagen: WHO Regional Office for Europe; 2025. Licence: CC BY-NC-SA 3.0 IGO.

Notes:	National data excluding Germany (Bremen only) and Kazakhstan (Almaty only).
Definitions:	Percentage of children not meeting WHO recommendations for physical activity, i.e. not playing actively/vigorously for at least one hour a day on an average day, as reported by their parent

Sugar consumption


Adults, 2016



References:	Source: Euromonitor International
Definitions:	Sugar consumption (Number of 500g sugar portions/person/month)

Estimated per capita sugar sweetened beverages intake

Adults, 2016




References:

Source: Euromonitor International

Prevalence of at least daily soft drink consumption


Children, 2022-2024



Survey type:	Self-reported
Age:	6-9
Sample size:	1078
Area covered:	National
References:	WHO European Childhood Obesity Surveillance Initiative (COSI): report on the sixth round of data collection, 2022– 2024. Copenhagen: WHO Regional Office for Europe; 2025. Licence: CC BY-NC-SA 3.0 IGO.

Notes:	National data excluding Germany (Bremen only) and Kazakhstan (Almaty only)
Definitions:	Prevalence of daily consumption of soft drinks containing sugar, as reported by the child's parent (%)


Boys, 2022-2024



Survey type:	Self-reported
Age:	6-9
Sample size:	1078
Area covered:	National
References:	WHO European Childhood Obesity Surveillance Initiative (COSI): report on the sixth round of data collection, 2022– 2024. Copenhagen: WHO Regional Office for Europe; 2025. Licence: CC BY-NC-SA 3.0 IGO.

Notes:	National data excluding Germany (Bremen only) and Kazakhstan (Almaty only)
Definitions:	Prevalence of daily consumption of soft drinks containing sugar, as reported by the child's parent (%)

Girls, 2022-2024




Survey type:	Self-reported
Age:	6-9
Sample size:	1078
Area covered:	National
References:	WHO European Childhood Obesity Surveillance Initiative (COSI): report on the sixth round of data collection, 2022– 2024. Copenhagen: WHO Regional Office for Europe; 2025. Licence: CC BY-NC-SA 3.0 IGO.

Notes:	National data excluding Germany (Bremen only) and Kazakhstan (Almaty only)
Definitions:	Prevalence of daily consumption of soft drinks containing sugar, as reported by the child's parent (%)

Prevalence of confectionery consumption


Adults, 2016



References:	Source: Euromonitor International
Definitions:	Prevalence of confectionery consumption (Number of 50g confectionery portions/person/month)

Prevalence of sweet/savoury snack consumption


Adults, 2016



References:	Source: Euromonitor International
Definitions:	Prevalence of sweet/savoury snack consumption (Number of 35g sweet/savoury snack portions/person/month)

Prevalence of less than daily fruit consumption


Children, 2022-2024



Survey type:	Self-reported
Age:	6-9
Sample size:	1070
Area covered:	National
References:	WHO European Childhood Obesity Surveillance Initiative (COSI): report on the sixth round of data collection, 2022– 2024. Copenhagen: WHO Regional Office for Europe; 2025. Licence: CC BY-NC-SA 3.0 IGO.

Notes:	National data excluding Germany (Bremen only) and Kazakhstan (Almaty only).
Definitions:	Proportion of children whose parents reported that their child consumes fruit less than daily


Boys, 2022-2024



Survey type:	Self-reported
Age:	6-9
Sample size:	1070
Area covered:	National
References:	WHO European Childhood Obesity Surveillance Initiative (COSI): report on the sixth round of data collection, 2022– 2024. Copenhagen: WHO Regional Office for Europe; 2025. Licence: CC BY-NC-SA 3.0 IGO.

Notes:	National data excluding Germany (Bremen only) and Kazakhstan (Almaty only).
Definitions:	Proportion of children whose parents reported that their child consumes fruit less than daily

Girls, 2022-2024




Survey type:	Self-reported
Age:	6-9
Sample size:	1070
Area covered:	National
References:	WHO European Childhood Obesity Surveillance Initiative (COSI): report on the sixth round of data collection, 2022– 2024. Copenhagen: WHO Regional Office for Europe; 2025. Licence: CC BY-NC-SA 3.0 IGO.

Notes:	National data excluding Germany (Bremen only) and Kazakhstan (Almaty only).
Definitions:	Proportion of children whose parents reported that their child consumes fruit less than daily

Prevalence of less than daily vegetable consumption


Children, 2022-2024



Survey type:	Self-reported
Age:	6-9
Sample size:	1071
Area covered:	National
References:	WHO European Childhood Obesity Surveillance Initiative (COSI): report on the sixth round of data collection, 2022– 2024. Copenhagen: WHO Regional Office for Europe; 2025. Licence: CC BY-NC-SA 3.0 IGO.

Notes:	National data excluding Germany (Bremen only) and Kazakhstan (Almaty only).
Definitions:	Proportion of children whose parents reported that their child consumes vegetables less than daily


Boys, 2022-2024



Survey type:	Self-reported
Age:	6-9
Sample size:	1071
Area covered:	National
References:	WHO European Childhood Obesity Surveillance Initiative (COSI): report on the sixth round of data collection, 2022– 2024. Copenhagen: WHO Regional Office for Europe; 2025. Licence: CC BY-NC-SA 3.0 IGO.

Notes:	National data excluding Germany (Bremen only) and Kazakhstan (Almaty only).
Definitions:	Proportion of children whose parents reported that their child consumes vegetables less than daily

Girls, 2022-2024




Survey type:	Self-reported
Age:	6-9
Sample size:	1071
Area covered:	National
References:	WHO European Childhood Obesity Surveillance Initiative (COSI): report on the sixth round of data collection, 2022– 2024. Copenhagen: WHO Regional Office for Europe; 2025. Licence: CC BY-NC-SA 3.0 IGO.

Notes:	National data excluding Germany (Bremen only) and Kazakhstan (Almaty only).
Definitions:	Proportion of children whose parents reported that their child consumes vegetables less than daily


Mental health - depression disorders

Adults, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)

Men, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)

Women, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)

Children, 2023




Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (under 20 years of age)

Boys, 2023



Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (under 20 years of age)


Girls, 2023



Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (under 20 years of age)


Mental health - anxiety disorders

Adults, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)

Men, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)

Women, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)

Children, 2023




Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (under 20 years of age)

Boys, 2023



Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (under 20 years of age)


Girls, 2023



Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (under 20 years of age)

Percent of population who cannot afford a healthy diet


Adults, 2025



Area covered:	National
References:	FAO. 2026. FAOSTAT: Cost and Affordability of a Healthy Diet (CoAHD). https://www.fao.org/faostat/en/#data/CAHD. Licence: CC-BY-4.0. [Last Accessed 12.10.26]
Notes:	The percent of the population whose food budget is below the cost of a healthy diet.


Oesophageal cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Breast cancer


Women, 2022



Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Colorectal cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Pancreatic cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Gallbladder cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Kidney cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Cancer of the uterus


Women, 2022



Age:	20+
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Raised blood pressure

Adults, 2019




Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.

Men, 2019



Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.


Women, 2019



Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.


Raised cholesterol

Adults, 2008




References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).

Men, 2008



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).


Women, 2008



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).


Raised fasting blood glucose

Men, 2014



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A869?lang=en
Definitions:	Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication).


Women, 2014



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A869?lang=en
Definitions:	Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication).

Diabetes prevalence


Adults, 2024



References: Reproduced with kind permission International Diabetes Federation. IDF Diabetes Atlas, 11th edn. Brussels, Belgium:International Diabetes Federation, 2025. http://www.diabetesatlas.org

Ovarian Cancer


Women, 2022



Age:	20+
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Leukemia


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Liver and intrahepatic bile duct Cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Multiple Myeloma


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Non Hodgkin Lymphoma


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Thyroid Cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

PDF created on November 4, 2026