

Report card Jordan




Lower-middle income

Contents	Page
Obesity prevalence	2
Trend: % Adults living with obesity in Jordan 2002-2017	5
Trend: % Adults living with overweight or obesity in Jordan 2002-2017	6
Trend: % Adults living with obesity in selected countries in the EMRO Region 1984-2017, selected countries	7
Overweight/obesity by education	10
Overweight/obesity by age	13
Overweight/obesity by region	15
Overweight/obesity by socio-economic group	18
Overweight/obesity by ethnicity	21
Insufficient physical activity	24
Estimated per capita fruit intake	30
Estimated per-capita processed meat intake	31
Estimated per capita whole grains intake	32
Mental health - depression disorders	33
Mental health - anxiety disorders	34
% Infants exclusively breastfed 0-5 months	35
Oesophageal cancer	36
Breast cancer	38
Colorectal cancer	39
Pancreatic cancer	41
Gallbladder cancer	43
Kidney cancer	45
Cancer of the uterus	47
Raised blood pressure	48
Raised cholesterol	51
Raised fasting blood glucose	54
Diabetes prevalence	56
Ovarian Cancer	57
Leukemia	58
Liver and intrahepatic bile duct Cancer	60
Multiple Myeloma	62
Non Hodgkin Lymphoma	64
Thyroid Cancer	66
Contextual factors	68




Obesity prevalence

Women, 2023






Girls, 2023





0-5 years, 2023

Overweight or obesity




	o o years boys	o o years gires	, it o o years
Survey type:			Measured
Age:			0-5
Sample size:			5274
Area covered:			National
References:	Department of Statistics	(DoS) [Jordan] and ICF. 2024. Jordan Population Jorda	and Family and Health Survey 2023. Amman, in, and Rockville, Maryland, USA: DoS and ICF.
Definitions:	Children whose weight-for-h	neight z score is more than two standard deviation	ons (+2 SD) above the median of the reference population are considered overweight
Cutoffs:			Weight for height



% Adults living with obesity in Jordan 2002-2017

Obesity, Women



Survey Measured

type:

References:

2002: WHO Infobase, Macro international data

2007: Department of Statistics [Jordan] and Macro International. 2008. Jordan Population and Family Health Survey 2007: Key Findings. Calverton, Maryland, USA: Department of Statistics and Macro International.

2009: Department of Statistics [Jordan] and ICF Macro. 2010. Jordan Population and Family Health Survey 2009. Calverton, Maryland, USA: Department of Statistics and ICF Macro.

2012: Department of Statistics/Jordan and ICF International. 2013. Jordan Population and Family Health Survey 2012.

Calverton, Maryland, USA: Department of Statistics/Jordan and ICF International.

2017: Department of Statistics/DOS and ICF. 2019. Jordan Population and Family and Health Survey 2017-18. Amman, Jordan, and Rockville, Maryland, USA: DOS and ICF. Available from: https://dhsprogram.com/publications/publication-FR346-DHS-Final-Reports.cfm [Accessed 27 March 2019].


Notes: Adults aged 15-49

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.



% Adults living with overweight or obesity in Jordan 2002-2017



Survey Measured

type:

References:

2002: WHO Infobase, Macro international data

2007: Department of Statistics [Jordan] and Macro International. 2008. Jordan Population and Family Health Survey 2007: Key Findings. Calverton, Maryland, USA: Department of Statistics and Macro International.

2009: Department of Statistics [Jordan] and ICF Macro. 2010. Jordan Population and Family Health Survey 2009. Calverton,

Maryland, USA: Department of Statistics and ICF Macro.

2012: Department of Statistics/Jordan and ICF International. 2013. Jordan Population and Family Health Survey 2012.

Calverton, Maryland, USA: Department of Statistics/Jordan and ICF International.

2017: Department of Statistics/DOS and ICF. 2019. Jordan Population and Family and Health Survey 2017-18. Amman, Jordan, and Rockville, Maryland, USA: DOS and ICF. Available from: https://dhsprogram.com/publications/publication-FR346-DHS-Final-Reports.cfm [Accessed 27 March 2019].

Notes: Adults aged 15-49

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.



% Adults living with obesity in selected countries in the EMRO Region 1984-2017, selected countries



Women



References:

1984, 1998: Benjelloun S. Nutrition transition in Morocco. Public Health Nutrition. 2002;5(1A):135-140

1991: Al-Lawati JA, Jousilahti PJ. Prevalence and 10year secular trend of obesity in Oman. Saudi Med J 2004;25:346-351 1997: Sibai AM, Hwalla N, Adra N, Rahal B. Prevalence and Covariates of Obesity in Lebanon: Findings from the First Epidemiolgical Study. Obesity Research 203;11:1353-1361

2000: Personal Communication from Morocco Minister of Health. National Survey carried out in 2000. Details in press for Journal of Hypertension (Arpil 03)

2002: WHO Infobase, Macro international data

2007: Department of Statistics [Jordan] and Macro International. 2008. Jordan Population and Family Health Survey 2007: Key Findings. Calverton, Maryland, USA: Department of Statistics and Macro International.

2008: Rhazi K El, Nejjari C, Zidouh A et al. Prevalence of obesity and associated sociodemographic and lifestyle factors in Morocco. Public Health Nutrition 2010:14(1)160-167

2009: WHO STEPS Report 2009 Lebanon

2011: WHO EMRO Egypt STEPS Survey 2011-12

2012: Department of Statistics/Jordan and ICF International. 2013. Jordan Population and Family Health Survey 2012.

Calverton, Maryland, USA: Department of Statistics/Jordan and ICF International.

2015: Ministry of Health and Population/Egypt, El-Zanaty and Associates/Egypt, and ICF International. 2015. Egypt Health Issues Survey 2015. Cairo, Egypt: Ministry of Health and Population/Egypt and ICF International. Available at

http://dhsprogram.com/publications/publication-FR313-DHS-Final-Reports.cfm#sthash.StgV9s6X.dpuf

2016: WHO STEPWISE APPROACH FOR NON-COMMUNICABLE DISEASESRISK FACTOR SURVEILLANCE, Lebanon, 2016-2017 https://www.who.int/ncds/surveillance/steps/Lebanon_STEPS_report_2016-2017.pdf?ua=1 (last accessed 16.10.19)


2017: Executive Summary. Oman National Non-Communicable Disease & their Risk Factors Survey 2017

https://mohcsr.gov.om/wp-content/uploads/2019/01/Executive-Summary_NCDsurvey2017_En.pdf (last accessed 17.10.19)

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.



Men



References:

1984, 1998: Benjelloun S. Nutrition transition in Morocco. Public Health Nutrition. 2002;5(1A):135-140

1991: Al-Lawati JA, Jousilahti PJ. Prevalence and 10year secular trend of obesity in Oman. Saudi Med J 2004;25:346-351 1997: Sibai AM, Hwalla N, Adra N, Rahal B. Prevalence and Covariates of Obesity in Lebanon: Findings from the First Epidemiolgical Study. Obesity Research 203;11:1353-1361

2000: Personal Communication from Morocco Minister of Health. National Survey carried out in 2000. Details in press for Journal of Hypertension (Arpil 03)

2002: WHO Infobase, Macro international data

2007: Department of Statistics [Jordan] and Macro International. 2008. Jordan Population and Family Health Survey 2007: Key Findings. Calverton, Maryland, USA: Department of Statistics and Macro International.

2008: Rhazi K El, Nejjari C, Zidouh A et al. Prevalence of obesity and associated sociodemographic and lifestyle factors in Morocco. Public Health Nutrition 2010:14(1)160-167

2009: WHO STEPS Report 2009 Lebanon

2011: WHO EMRO Egypt STEPS Survey 2011-12

2012: Department of Statistics/Jordan and ICF International. 2013. Jordan Population and Family Health Survey 2012.

Calverton, Maryland, USA: Department of Statistics/Jordan and ICF International.

2015: Ministry of Health and Population/Egypt, El-Zanaty and Associates/Egypt, and ICF International. 2015. Egypt Health Issues Survey 2015. Cairo, Egypt: Ministry of Health and Population/Egypt and ICF International. Available at

http://dhsprogram.com/publications/publication-FR313-DHS-Final-Reports.cfm#sthash.StgV9s6X.dpuf

2016: WHO STEPWISE APPROACH FOR NON-COMMUNICABLE DISEASESRISK FACTOR SURVEILLANCE, Lebanon, 2016-2017 https://www.who.int/ncds/surveillance/steps/Lebanon_STEPS_report_2016-2017.pdf?ua=1 (last accessed 16.10.19)

2017: Executive Summary. Oman National Non-Communicable Disease & their Risk Factors Survey 2017

https://mohcsr.gov.om/wp-content/uploads/2019/01/Executive-Summary_NCDsurvey2017_En.pdf (last accessed 17.10.19)

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.




Overweight/obesity by education

Women, 2023

Survey type:





Age:	20-49
Sample size:	8034
Area covered:	National
References:	Department of Statistics (DoS) [Jordan] and ICF. 2024. Jordan Population and Family and Health Survey 2023. Amman, Jordan, and Rockville, Maryland, USA: DoS and ICF.


Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Measured



Girls, 2023






Survey type:	Measured
Age:	15-19
Sample size:	2383
Area covered:	National
References:	Department of Statistics (DoS) [Jordan] and ICF. 2024. Jordan Population and Family and Health Survey 2023. Amman, Jordan, and Rockville, Maryland, USA: DoS and ICF.
Notes:	Insufficent sample size to demonstrate 'no education' category.
Cutoffs:	WHO



0-5 years, 2023






Overweight/obesity by age

Women, 2023





Sample size:

Area
Covered:


References:

Department of Statistics (DoS) [Jordan] and ICF. 2024. Jordan Population and Family and Health Survey 2023. Amman, Jordan, and Rockville, Maryland, USA: DoS and ICF.

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².



Children, 2006






Overweight/obesity by region

Women, 2023

Survey type:





Age:	20-49
Sample size:	8034
Area covered:	National
References:	Department of Statistics (DoS) [Jordan] and ICF. 2024. Jordan Population and Family and Health Survey 2023. Amman, Jordan, and Rockville, Maryland, USA: DoS and ICF.


 $Unless \ otherwise \ noted, overweight \ refers \ to \ a \ BMI \ between \ 25kg \ and \ 29.9kg/m^2, obesity \ refers \ to \ a \ BMI \ greater \ than \ 30kg/m^2.$

Measured



Girls, 2023

Cutoffs:




WHO



0-5 years, 2023

Overweight or obesity




Survey type:	Measured
Sample size:	5274
Area covered:	National
References:	Department of Statistics (DoS) [Jordan] and ICF. 2024. Jordan Population and Family and Health Survey 2023. Amman, Jordan, and Rockville, Maryland, USA: DoS and ICF.
Definitions:	Children whose weight-for-height z score is more than two standard deviations (+2 SD) above the median of the reference population are considered overweight
Cutoffs:	Weight for height



Overweight/obesity by socio-economic group

Women, 2023



Area National covered:


References: Department of Statistics (DoS) [Jordan] and ICF. 2024. Jordan Population and Family and Health Survey 2023. Amman, Jordan, and Rockville, Maryland, USA: DoS and ICF.

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².



Girls, 2023







Survey type:	Measured
Age:	15-19
Sample size:	2383
Area covered:	National
References:	Department of Statistics (DoS) [Jordan] and ICF. 2024. Jordan Population and Family and Health Survey 2023. Amman, Jordan, and Rockville, Maryland, USA: DoS and ICF.
Cutoffs:	who



0-5 years, 2023






Survey type:	Measured
Sample size:	5274
Area covered:	National
References:	Department of Statistics (DoS) [Jordan] and ICF. 2024. Jordan Population and Family and Health Survey 2023. Amman, Jordan, and Rockville, Maryland, USA: DoS and ICF.
Definitions:	Children whose weight-for-height z score is more than two standard deviations (+2 SD) above the median of the reference population are considered overweight



Overweight/obesity by ethnicity


Ethnic groups are as defined by publication of origin and are not as defined by WOF. In some instances ethnicity is conflated with nationality and/or race.

Women, 2023






Girls, 2023





0-5 years, 2023

Overweight or obesity




Survey type:	Measured
Sample size:	5274
Area covered:	National
References:	Department of Statistics (DoS) [Jordan] and ICF. 2024. Jordan Population and Family and Health Survey 2023. Amman, Jordan, and Rockville, Maryland, USA: DoS and ICF.
Definitions:	Children whose weight-for-height z score is more than two standard deviations (+2 SD) above the median of the reference population are considered overweight
Cutoffs:	Weight for height




Insufficient physical activity

Adults, 2022





Men, 2022



Survey	Seti-reported
type:	
Age:	18+
Area	National
covered:	


References: WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at <a href="https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicator-data/gho/data/indicator-data/gho/data/gh

<u>among-adults-aged-18-years-</u>(age-standardized-estimate)-(-)

Definitions: Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.



Women, 2022



Julycy	Sett reported
type:	
Age:	18+
Area	National
covered:	

References: WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at <a href="https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicator-details/GHO/prevalence-of-insufficient-physical-activity-data/gho/data/indicator-data/gho/data/indicator-data/gho/data/gh


among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions: Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.



Children, 2016

Definitions:




Survey type:	Self-reported
Age:	11-17
References:	Global Health Observatory data repository, World Health Organisation, https://apps.who.int/gho/data/node.main.A893ADO?lang=en (last accessed 16.03.21)
Notes:	% of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.

% Adolescents insufficiently active (age standardised estimate)



Boys, 2016



Survey	Self-reported
type:	
Age:	11-17
References:	Global Health Observatory data repository, World Health Organisation, https://apps.who.int/gho/data/node.main.A893ADO?lang=en (last accessed 16.03.21)
Notes:	% of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60


Notes: % of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.

Definitions: % Adolescents insufficiently active (age standardised estimate)



Girls, 2016

Definitions:




Survey type:	Self-reported
Age:	11-17
References:	Global Health Observatory data repository, World Health Organisation, https://apps.who.int/gho/data/node.main.A893ADO?lang=en (last accessed 16.03.21)
Notes:	% of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.

% Adolescents insufficiently active (age standardised estimate)



Estimated per capita fruit intake


Adults, 2017





Estimated per-capita processed meat intake

Adults, 2017



Survey type:

Age:

Clobal Burden of Disease, the Institute for Health Metrics and Evaluation http://ghdx.healthdata.org/


Definitions:

Estimated per-capita processed meat intake (g per day)



Estimated per capita whole grains intake


Adults, 2017





Mental health - depression disorders

Adults, 2015



References: Prevalence data from Global Burden of Disease study 2015 (http://ghdx.healthdata.org) published in: Depression and Other Common Mental Disorders: Global Health Estimates. Geneva: World Health Organization; 2017. Licence: CC BY-NC-SA 3.0 IGO.


Definitions:

% of population with depression disorders



Mental health - anxiety disorders

Adults, 2015



References: Prevalence data from Global Burden of Disease study 2015 (http://ghdx.healthdata.org) published in: Depression and Other Common Mental Disorders: Global Health Estimates. Geneva: World Health Organization; 2017. Licence: CC BY-NC-SA 3.0


IGO.

Definitions: % of population with anxiety disorders



% Infants exclusively breastfed 0-5 months

0-5 years, 2010-2023



References:

Jordan National Micronutrient and Nutritional Survey 2019. Amman, Jordan; 2021.


Notes:

Full details are available. Original citation United Nations Children's Fund, Division of Data, Analysis, Planning and Monitoring (2023). Global UNICEF Global Databases: Infant and Young Child Feeding: Exclusive breastfeeding, New York, October 2023.




Oesophageal cancer

Men, 2022










Breast cancer

Women, 2022





Area National covered:

References:

Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from:


https://gco.iarc.who.int/today, accessed [16.07.24]

Definitions: Incidence per 100,000




Colorectal cancer

Men, 2022











Pancreatic cancer

Men, 2022



Definitions: Incidence per 100,000







Gallbladder cancer

Men, 2022










Kidney cancer

Men, 2022










Cancer of the uterus

Women, 2022



Age: 20+

References:

Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from:


https://gco.iarc.who.int/today, accessed [16.07.24]

Definitions: Incidence per 100,000



Raised blood pressure

Adults, 2015



References:

Global Health Observatory data repository, World Health Organisation,


http://apps.who.int/gho/data/node.main.A875?lang=en

Definitions:

Age Standardised estimated % Raised blood pressure 2015 (SBP>=140 OR DBP>=90).



Men, 2015



References:


Global Health Observatory data repository, World Health Organisation,

http://apps.who.int/gho/data/node.main.A875?lang=en

Definitions:

Age Standardised estimated % Raised blood pressure 2015 (SBP>=140 OR DBP>=90).





References:

Global Health Observatory data repository, World Health Organisation,

http://apps.who.int/gho/data/node.main.A875?lang=en


Definitions:

Age Standardised estimated % Raised blood pressure 2015 (SBP>=140 OR DBP>=90).



Raised cholesterol

Adults, 2008



References:


Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885

Definitions:

% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).



Men, 2008




References:

Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885

Definitions:

% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).





References: Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885


Definitions:

% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).



Raised fasting blood glucose

Men, 2014



References:


Global Health Observatory data repository, World Health Organisation,

http://apps.who.int/gho/data/node.main.A869?lang=en

Definitions:

Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication).





References:

Global Health Observatory data repository, World Health Organisation,

http://apps.who.int/gho/data/node.main.A869?lang=en


Definitions:

Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication).



Diabetes prevalence


Adults, 2021





Ovarian Cancer

Women, 2022



Age: 20+

References:

Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from:

https://gco.iarc.who.int/today, accessed [16.07.24]

Definitions: Incidence per 100,000



Leukemia

Men, 2022	<u>.</u>						
4							
3							
2							
1							
0							
⁻¹ _1	0	1	2	3	4	5	6
Age:							20+
Area covered:							National
References:			nne M, Colombet M, ncer Today. Lyon, Fr	ance: International		ch on Cancer. Availa	able from:
Definitions:						Incidence per	100,000
Unle	ess otherwise note	ed, overweight refer	rs to a BMI between	25kg and 29.9kg/ı	m², obesity refers to	a BMI greater than	30kg/m².




Wome	n, 2022						
4							
3							
2							
1							
0							
⁻¹ 1	0	1	2	3	4	5	6
Age:							20+
Area covered:							National
References	•			rance: International	Znaor A, Soerjomat Agency for Researc tps://gco.iarc.who.ir	ch on Cancer. Availa	able from:
Definitions	s:					Incidence per	r 100,000
	Unless otherwise note	ed, overweight refer	s to a BMI between	25kg and 29.9kg/i	m², obesity refers to	a BMI greater than	30kg/m².




Liver and intrahepatic bile duct Cancer

Men, 2022










Multiple Myeloma

Men, 2022



Age: 20+

Area National


covered:
References:

Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from:

https://gco.iarc.who.int/today, accessed [16.07.24]

Definitions: Indicence per 100,000





References:

Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from:


https://gco.iarc.who.int/today, accessed [16.07.24]

Definitions: Indicence per 100,000



Non Hodgkin Lymphoma

Men, 2022



Area National


covered:
References:

Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from:

https://gco.iarc.who.int/today, accessed [16.07.24]

Definitions: Incidence per 100,000





References:

Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from:


https://gco.iarc.who.int/today, accessed [16.07.24]

Definitions: Incidence per 100,000



Thyroid Cancer

Men, 2022



References:

covered:


Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from:

https://gco.iarc.who.int/today, accessed [16.07.24]

Definitions: Incidence per 100,000



References:



Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from:

https://gco.iarc.who.int/today, accessed [16.07.24]

Definitions: Incidence per 100,000



Contextual factors

Disclaimer: These contextual factors should be interpreted with care. Results are updated as regularly as possible and use very specific criteria. The criteria used and full definitions are available for download at the bottom of this page.



Labelling

Is there mandatory nutrition labelling?	×
Front-of-package labelling?	×
Back-of-pack nutrition declaration?	?
Color coding?	×
Warning label?	×





Regulation and marketing

Are there fiscal policies on unhealthy products?	X
Tax on unhealthy foods?	×
Tax on unhealthy drinks?	×
Are there fiscal policies on healthy products?	X
Subsidy on fruits?	×
Subsidy on vegetables?	×
Subsidy on other healthy products?	×
Mandatory limit or ban of trans fat (all settings)?	X
Mandatory limit of trans fats in place (all settings)?	×
Ban on trans-fats or phos in place (all settings)?	×
Are there any mandatory policies/marketing restrictions on the promotion of unhealthy food/drinks to children?	X
Mandatory restriction on broadcast media?	X
Mandatory restriction on non-broadcast media?	×
Voluntary policies/marketing restrictions on the promotion of unhealthy food/drinks to children?	×
Are there mandatory standards for food in schools?	~
Are there any mandatory nutrient limits in any manufactured food products?	~
Nutrition standards for public sector procurement?	X





Political will and support

National obesity strategy or nutrition and physical activity national strategy?	×
National obesity strategy?	×
National childhood obesity strategy?	×
Comprehensive nutrition strategy?	~
Comprehensive physical activity strategy?	?
Evidence-based dietary guidelines and/or RDAs?	?
National target(s) on reducing obesity?	~
Guidelines/policy on obesity treatment?	~
Promotion of breastfeeding?	?
Monitoring and surveillance	
Monitoring of the prevalence and incidence for the main obesity-related NCDs and risk factors?	~
Within 5 years?	~
Governance and resource	
Multi-sectoral national co-ordination mechanism for obesity or nutrition (including obesity)?	~
Key ✓ Present ✓ Present ✓ Incoming ✓ Absent ? Un (voluntary)	known

Last updated October 24, 2022



PDF created on October 1, 2024