Mexico



Country report card


This report card contains the latest data available on the Global Obesity Observatory on overweight and obesity . Where available, data on common and relevant obesity drivers and comorbidities are also presented.

View the latest version of this report on the Global Obesity Observatory at https://data.worldobesity.org/country/mexico-139/.

Contents	Page
Obesity prevalence	3
Trend: % 12-19 year old children living with overweight or obesity in Mexico 2012-2023	10
Trend: % 5-11 year old children living with overweight or obesity in Mexico 2012-2023	17
Trend: % Adults living with obesity in Mexico 1992-2023	24
Trend: % Adults living with overweight or obesity in Mexico 1992-2023	31
Trend: % Adults living with obesity in selected countries in the Americas Region 1960-2018, selected countries	38
Overweight/obesity by education	43
Overweight/obesity by age	50
Overweight/obesity by region	55
Overweight/obesity by socio-economic group	66
Overweight/obesity by ethnicity	73
Double burden of underweight & obesity	76
Insufficient physical activity	83
Average weekly frequency of fast food consumption	96
Mental health - depression disorders	103
Mental health - anxiety disorders	110
% Infants exclusively breastfed 0-5 months	117
Percent of population who cannot afford a healthy diet	119
Oesophageal cancer	120
Breast cancer	123
Colorectal cancer	125
Pancreatic cancer	128
Gallbladder cancer	131
Kidney cancer	134
Cancer of the uterus	137
Raised blood pressure	139
Raised cholesterol	143
Raised fasting blood glucose	146
Diabetes prevalence	148
Ovarian Cancer	149
Leukemia	151
Liver and intrahepatic bile duct Cancer	156
Multiple Myeloma	159
Non Hodgkin Lymphoma	164
Thyroid Cancer	169

Obesity prevalence


Adults, 2023



Survey type:	Measured
Age:	20+
Sample size:	5474
Area covered:	National
References:	Shamah-Levy T, Lazcano-Ponce EC, Cuevas-Nasu L, Romero-Martínez M, Gaona-Pineda EB, Gómez-Acosta LM, Mendoza-Alvarado LR, Méndez-Gómez-Humarán I. Encuesta Nacional de Salud y Nutrición Continua 2023. Resultados Nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública, 2024.

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Children, 2023



Survey type:	Measured
Age:	12-19
Sample size:	1303
Area covered:	National
References:	Shamah-Levy T, Lazcano-Ponce EC, Cuevas-Nasu L, Romero-Martínez M, Gaona-Pineda EB, Gómez-Acosta LM, Mendoza-Alvarado LR, Méndez-Gómez-Humarán I. Encuesta Nacional de Salud y Nutrición Continua 2023. Resultados Nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública, 2024.

Cutoffs:	WHO

0-5 years, 2022




Survey type:	Measured
Age:	0-5
Sample size:	10122251
Area covered:	National
References:	Encuesta Nacional de Salud y Nutrición Continua, Mexico 2022
Notes:	UNICEF/WHO/World Bank Joint Child Malnutrition Estimates Expanded Database: Overweight estimates: National and Disaggregated, July 2025, New York. Weighted sample size. For more information, please consult https://data.unicef.org/resources/jme/.

Definitions:	Percentage of children under 5 years of age falling above 2 standard deviations (moderate and severe) from the median weight-for-height of the reference population.
Cutoffs:	=>+2SD

% 12-19 year old children living with overweight or obesity in Mexico 2012-2023


Boys



Survey type:	Measured
References:	2012: ENSANUT 2012, https://ensanut.insp.mx/encuestas/ensanut2012/doctos/informes/ENSANUT2012ResultadosNacionales.pdf (Accessed 15.12.20)
	2016: Shamah-Levy T, Cuevas-Nasu L, Gaona-Pineda EB,Gómez-Acosta LM, Morales-Ruán MC,Hernández-Ávila M, Rivera-Dommarco JA. Overweight and obesity in children and adolescents,2016 Halfway National Health and Nutrition Survey update. Salud Publica Mex 2018;60:244-253.
	2018: Encuesta Nacional de Salud y Nutrición 2018. https://ensanut.insp.mx/encuestas/ensanut2018/informes.php (Accessed 15.12.20)
	2020: Shamah-Levy T, Romero-Martínez M, Barrientos-Gutiérrez T, Cuevas-Nasu L, Bautista-Arredondo S, Colchero MA, GaonaPineda EB, Lazcano-Ponce E, Martínez-Barnetche J, Alpuche-Arana C, Rivera-Dommarco J. Encuesta Nacional de Salud y Nutrición 2020 sobre Covid-19. Resultados nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública, 2021.

Notes:	Aged 12-19
Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.	


Girls



Survey type:	Measured
References:	2012: ENSANUT 2012, https://ensanut.insp.mx/encuestas/ensanut2012/doctos/informes/ENSANUT2012ResultadosNacionales.pdf (Accessed 15.12.20)
	2016: Shamah-Levy T, Cuevas-Nasu L, Gaona-Pineda EB,Gómez-Acosta LM, Morales-Ruán MC,Hernández-Ávila M, Rivera-Dommarco JA. Overweight and obesity in children and adolescents,2016 Halfway National Health and Nutrition Survey update. Salud Publica Mex 2018;60:244-253.
	2018: Encuesta Nacional de Salud y Nutrición 2018. https://ensanut.insp.mx/encuestas/ensanut2018/informes.php (Accessed 15.12.20)
	2020: Shamah-Levy T, Romero-Martínez M, Barrientos-Gutiérrez T, Cuevas-Nasu L, Bautista-Arredondo S, Colchero MA, GaonaPineda EB, Lazcano-Ponce E, Martínez-Barnetche J, Alpuche-Arana C, Rivera-Dommarco J. Encuesta Nacional de Salud y Nutrición 2020 sobre Covid-19. Resultados nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública, 2021.

Notes:	Aged 12-19
Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.	

Boys and girls




Survey type:	Measured
References:	2012: ENSANUT 2012, https://ensanut.insp.mx/encuestas/ensanut2012/doctos/informes/ENSANUT2012ResultadosNacionales.pdf (Accessed 15.12.20) 2016: Shamah-Levy T, Cuevas-Nasu L, Gaona-Pineda EB,Gómez-Acosta LM, Morales-Ruán MC,Hernández-Ávila M, Rivera-Dommarco JA. Overweight and obesity in children and adolescents,2016 Halfway National Health and Nutrition Survey update. Salud Publica Mex 2018;60:244-253. 2018: Encuesta Nacional de Salud y Nutrición 2018. https://ensanut.insp.mx/encuestas/ensanut2018/informes.php (Accessed 15.12.20) 2020: Shamah-Levy T, Romero-Martínez M, Barrientos-Gutiérrez T, Cuevas-Nasu L, Bautista-Arredondo S, Colchero MA, GaonaPineda EB, Lazcano-Ponce E, Martínez-Barnetche J, Alpuche-Arana C, Rivera-Dommarco J. Encuesta Nacional de Salud y Nutrición 2020 sobre Covid-19. Resultados nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública, 2021.

Notes:	Aged 12-19
Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.	

% 5-11 year old children living with overweight or obesity in Mexico 2012-2023


Boys



Survey type:	Measured
References:	2012: ENSANUT 2012 http://ensanut.insp.mx/informes/ENSANUT2012ResultadosNacionales.pdf accessed 25th June 2014 2020: Shamah-Levy T, Romero-Martínez M, Barrientos-Gutiérrez T, Cuevas-Nasu L, Bautista-Arredondo S, Colchero MA, GaonaPineda EB, Lazcano-Ponce E, Martínez-Barnetche J, Alpuche-Arana C, Rivera-Dommarco J. Encuesta Nacional de Salud y Nutrición 2020 sobre Covid-19. Resultados nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública, 2021. https://ensanut.insp.mx/encuestas/ensanutcontinua2020/doctos/informes/ensanutCovid19ResultadosNacionales.pdf (Accessed 11.01.2022) 2021: Shamah-Levy T, Romero-Martínez M, Barrientos-Gutiérrez T, Cuevas-Nasu L, Bautista-Arredondo S, Colchero MA, Gaona-Pineda EB, Lazcano-Ponce E,Martínez-Barnetche J, Alpuche-Arana C, Rivera-Dommarco J. Encuesta Nacional de Salud y Nutrición 2021 sobre Covid-19. Resultados nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública, 2022.

Notes:	Aged 5-11
Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.	


Girls



Survey type:	Measured
References:	2012: ENSANUT 2012 http://ensanut.insp.mx/informes/ENSANUT2012ResultadosNacionales.pdf accessed 25th June 2014 2020: Shamah-Levy T, Romero-Martínez M, Barrientos-Gutiérrez T, Cuevas-Nasu L, Bautista-Arredondo S, Colchero MA, GaonaPineda EB, Lazcano-Ponce E, Martínez-Barnetche J, Alpuche-Arana C, Rivera-Dommarco J. Encuesta Nacional de Salud y Nutrición 2020 sobre Covid-19. Resultados nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública, 2021. https://ensanut.insp.mx/encuestas/ensanutcontinua2020/doctos/informes/ensanutCovid19ResultadosNacionales.pdf (Accessed 11.01.2022) 2021: Shamah-Levy T, Romero-Martínez M, Barrientos-Gutiérrez T, Cuevas-Nasu L, Bautista-Arredondo S, Colchero MA, Gaona-Pineda EB, Lazcano-Ponce E,Martínez-Barnetche J, Alpuche-Arana C, Rivera-Dommarco J. Encuesta Nacional de Salud y Nutrición 2021 sobre Covid-19. Resultados nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública, 2022.

Notes:	Aged 5-11
Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.	

Boys and girls




Survey type:	Measured
References:	2012: ENSANUT 2012 http://ensanut.insp.mx/informes/ENSANUT2012ResultadosNacionales.pdf accessed 25th June 2014 2020: Shamah-Levy T, Romero-Martínez M, Barrientos-Gutiérrez T, Cuevas-Nasu L, Bautista-Arredondo S, Colchero MA, GaonaPineda EB, Lazcano-Ponce E, Martínez-Barnetche J, Alpuche-Arana C, Rivera-Dommarco J. Encuesta Nacional de Salud y Nutrición 2020 sobre Covid-19. Resultados nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública, 2021. https://ensanut.insp.mx/encuestas/ensanutcontinua2020/doctos/informes/ensanutCovid19ResultadosNacionales.pdf (Accessed 11.01.2022) 2021: Shamah-Levy T, Romero-Martínez M, Barrientos-Gutiérrez T, Cuevas-Nasu L, Bautista-Arredondo S, Colchero MA, Gaona-Pineda EB, Lazcano-Ponce E,Martínez-Barnetche J, Alpuche-Arana C, Rivera-Dommarco J. Encuesta Nacional de Salud y Nutrición 2021 sobre Covid-19. Resultados nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública, 2022.

Notes:	Aged 5-11
Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.	

% Adults living with obesity in Mexico 1992-2023

Men




Survey type:	Measured
References:	1992: Arroyo et al. Prevalence of Pre-Obesity and Obesity in Urban Adult Mexicans in Comparison with other Large Surveys. Obesity Research. 2000;8:179-185 2000: Personal Communication from Dr C Sanchez Castillio, Body Mass Index cutoff points for Type 2 Diabetes Melliuts and Hypertension in the Mexican National Health Survey 2000. 2006: Olaiz-Fernández G, Rivera-Dommarco J, Shamah-Levy T, Rojas R, Villalpando-Hernández S, Hernández-Avila M, Sepúlveda-Amor J. Encuesta Nacional de Salud y Nutrición 2006. Cuernavaca, México: Instituto Nacional de Salud Pública, 2006. (National Health and Nutrition Survey 2006). 2012: ENSANUT 2012 http://ensanut.insp.mx/informes/ENSANUT2012ResultadosNacionales.pdf accessed 25th June 2014 2016: Encuesta Nacional de Salud y Nutrición de Medio Camino 2016

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women




Survey type:	Measured
References:	1992: Arroyo et al. Prevalence of Pre-Obesity and Obesity in Urban Adult Mexicans in Comparison with other Large Surveys. Obesity Research. 2000;8:179-185 2000: Personal Communication from Dr C Sanchez Castillio, Body Mass Index cutoff points for Type 2 Diabetes Melliuts and Hypertension in the Mexican National Health Survey 2000. 2006: Olaiz-Fernández G, Rivera-Dommarco J, Shamah-Levy T, Rojas R, Villalpando-Hernández S, Hernández-Avila M, Sepúlveda-Amor J. Encuesta Nacional de Salud y Nutrición 2006. Cuernavaca, México: Instituto Nacional de Salud Pública, 2006. (National Health and Nutrition Survey 2006). 2012: ENSANUT 2012 http://ensanut.insp.mx/informes/ENSANUT2012ResultadosNacionales.pdf accessed 25th June 2014 2016: Encuesta Nacional de Salud y Nutrición de Medio Camino 2016

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Men and women




Survey type:	Measured
References:	1992: Arroyo et al. Prevalence of Pre-Obesity and Obesity in Urban Adult Mexicans in Comparison with other Large Surveys. Obesity Research. 2000;8:179-185 2000: Personal Communication from Dr C Sanchez Castillio, Body Mass Index cutoff points for Type 2 Diabetes Melliuts and Hypertension in the Mexican National Health Survey 2000. 2006: Olaiz-Fernández G, Rivera-Dommarco J, Shamah-Levy T, Rojas R, Villalpando-Hernández S, Hernández-Avila M, Sepúlveda-Amor J. Encuesta Nacional de Salud y Nutrición 2006. Cuernavaca, México: Instituto Nacional de Salud Pública, 2006. (National Health and Nutrition Survey 2006). 2012: ENSANUT 2012 http://ensanut.insp.mx/informes/ENSANUT2012ResultadosNacionales.pdf accessed 25th June 2014 2016: Encuesta Nacional de Salud y Nutrición de Medio Camino 2016

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with overweight or obesity in Mexico 1992-2023

Men




Survey type:	Measured
References:	1992: Arroyo et al. Prevalence of Pre-Obesity and Obesity in Urban Adult Mexicans in Comparison with other Large Surveys. Obesity Research. 2000;8:179-185 2000: Personal Communication from Dr C Sanchez Castillio, Body Mass Index cutoff points for Type 2 Diabetes Melliuts and Hypertension in the Mexican National Health Survey 2000. 2006: Olaiz-Fernández G, Rivera-Dommarco J, Shamah-Levy T, Rojas R, Villalpando-Hernández S, Hernández-Avila M, Sepúlveda-Amor J. Encuesta Nacional de Salud y Nutrición 2006. Cuernavaca, México: Instituto Nacional de Salud Pública, 2006. (National Health and Nutrition Survey 2006). 2012: ENSANUT 2012 http://ensanut.insp.mx/informes/ENSANUT2012ResultadosNacionales.pdf accessed 25th June 2014 2016: Encuesta Nacional de Salud y Nutrición de Medio Camino 2016

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women




Survey type:	Measured
References:	1992: Arroyo et al. Prevalence of Pre-Obesity and Obesity in Urban Adult Mexicans in Comparison with other Large Surveys. Obesity Research. 2000;8:179-185 2000: Personal Communication from Dr C Sanchez Castillio, Body Mass Index cutoff points for Type 2 Diabetes Melliuts and Hypertension in the Mexican National Health Survey 2000. 2006: Olaiz-Fernández G, Rivera-Dommarco J, Shamah-Levy T, Rojas R, Villalpando-Hernández S, Hernández-Avila M, Sepúlveda-Amor J. Encuesta Nacional de Salud y Nutrición 2006. Cuernavaca, México: Instituto Nacional de Salud Pública, 2006. (National Health and Nutrition Survey 2006). 2012: ENSANUT 2012 http://ensanut.insp.mx/informes/ENSANUT2012ResultadosNacionales.pdf accessed 25th June 2014 2016: Encuesta Nacional de Salud y Nutrición de Medio Camino 2016

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Men and women




Survey type:	Measured
References:	1992: Arroyo et al. Prevalence of Pre-Obesity and Obesity in Urban Adult Mexicans in Comparison with other Large Surveys. Obesity Research. 2000;8:179-185 2000: Personal Communication from Dr C Sanchez Castillio, Body Mass Index cutoff points for Type 2 Diabetes Melliuts and Hypertension in the Mexican National Health Survey 2000. 2006: Olaiz-Fernández G, Rivera-Dommarco J, Shamah-Levy T, Rojas R, Villalpando-Hernández S, Hernández-Avila M, Sepúlveda-Amor J. Encuesta Nacional de Salud y Nutrición 2006. Cuernavaca, México: Instituto Nacional de Salud Pública, 2006. (National Health and Nutrition Survey 2006). 2012: ENSANUT 2012 http://ensanut.insp.mx/informes/ENSANUT2012ResultadosNacionales.pdf accessed 25th June 2014 2016: Encuesta Nacional de Salud y Nutrición de Medio Camino 2016

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with obesity in selected countries in the Americas Region 1960-2018, selected countries

Men



References:

1960, 1971, 1973, 1976, 1991: Flegal KM, Carroll MD, Kuczmarski RJ, Johnson CL. Overweight and obesity in the United States: prevalence and trends, 1960-1994. International Journal of Obesity (1998);22:39-47

1975: Monteiro CA, Conde WL, Popking BM. Is obesity replacing or adding to undernutrition? Evidence from different social classes in Brazil. 2002. Public Health Nutrition:51(1A), 105-112

1988: Berrios X, Jadue I, Zenteno J, Ross MI, Rodriguez H. Prevalencia de factores de riesgo para enfermedades cronicas. Estudio de la poblacion general de la region Metropolitana, 1986-1987. Rev. Med. Chile. 1990;118:597-604


1992, 1994, 1995: Martorell R, Khan LK, Hughes ML, Grummer Strawn LM. Obesity in women from developing countries. EJCN (2000) 54;247-252

1997: Filozof C, Gonzales C, Sereday M, Mazza C, Braguinsky J. Obesity prevalence and trends in Latin American countries. Obesity Reviews, 2001;2:99-196

1998: Instituto Nacional de Estadística - INE/Guatemala and Macro International. 1999. Guatemala Encuesta Nacional de Salud Materno

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women



References:

1960, 1971, 1973, 1976, 1991: Flegal KM, Carroll MD, Kuczmarski RJ, Johnson CL. Overweight and obesity in the United States: prevalence and trends, 1960-1994. International Journal of Obesity (1998);22:39-47

1975: Monteiro CA, Conde WL, Popking BM. Is obesity replacing or adding to undernutrition? Evidence from different social classes in Brazil. 2002. Public Health Nutrition:51(1A), 105-112

1988: Berrios X, Jadue I, Zenteno J, Ross MI, Rodriguez H. Prevalencia de factores de riesgo para enfermedades cronicas. Estudio de la poblacion general de la region Metropolitana, 1986-1987. Rev. Med. Chile. 1990;118:597-604

1992, 1994, 1995: Martorell R, Khan LK, Hughes ML, Grummer Strawn LM. Obesity in women from developing countries. EJCN (2000) 54;247-252


1997: Filozof C, Gonzales C, Sereday M, Mazza C, Braguinsky J. Obesity prevalence and trends in Latin American countries. Obesity Reviews, 2001;2:99-196

1998: Instituto Nacional de Estadística - INE/Guatemala and Macro International. 1999. Guatemala Encuesta Nacional de Salud Materno

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Overweight/obesity by education


Adults, 2022



Survey type:	Measured
Age:	20+
Sample size:	8563
Area covered:	National
References:	Campos-Nonato I, Galván-Valencia O,Hernández-Barrera L, Oviedo-Solís C, Barquera S.Prevalence of obesity and associated risk factors in Mexican adults: results of the Ensanut 2022.Salud Publica Mex. 2023;65(supl 1): S238-S247.https://doi.org/10.21149/14809. Available at https://saludpublica.mx/index.php/spm/article/view/14809/12424 (last accessed 11.12.23)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Children, 2020-2024



Survey type:	Measured
Age:	5-11
Sample size:	10,979
Area covered:	National
References:	Shamah-Levy T, Gaona-Pineda EB, Cuevas-Nasu L, Méndez-Gómez-Humarán I, Rodríguez-Ramírez S, Ávila-Arcos MA, Morales-Ruan C, Valenzuela-Bravo DG. National and state prevalence of overweight and obesity in Mexican school-age and adolescent population, and its associated factors. Salud Publica Mex. 2025;67:609-621. https://doi.org/10.21149/17311 [Accessed 15.01.26]

Definitions:	Mother's education
Cutoffs:	WHO 2007

0-5 years, 2020




Sample size:	9891242
References:	Other: Encuesta Nacional de Salud y Nutrición Continua COVID-19, Mexico 2020
Notes:	UNICEF/WHO/World Bank Joint Child Malnutrition Estimates Expanded Database: Overweight (Survey Estimates), May 2023, New York. For more information about the methodology, please consult https://data.unicef.org/resources/jme-2023-country-consultations/ Percentage of children under 5 years of age falling above 2 standard deviations (moderate and severe) from the median weight-for-height of the reference population.

Definitions:	=>+2SD

Overweight/obesity by age


Adults, 2022



Survey type:	Measured
Sample size:	8563
Area covered:	National
References:	Campos-Nonato I, Galván-Valencia O,Hernández-Barrera L, Oviedo-Solís C, Barquera S.Prevalence of obesity and associated risk factors in Mexican adults: results of the Ensanut 2022.Salud Publica Mex. 2023;65(supl 1): S238-S247.https://doi.org/10.21149/14809. Available at https://saludpublica.mx/index.php/spm/article/view/14809/12424 (last accessed 11.12.23)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Children, 2023




Survey type:	Measured
Sample size:	2716
Area covered:	National
References:	Shamah-Levy T, Lazcano-Ponce EC, Cuevas-Nasu L, Romero-Martínez M, Gaona-Pineda EB, Gómez-Acosta LM, Mendoza-Alvarado LR, Méndez-Gómez-Humarán I. Encuesta Nacional de Salud y Nutrición Continua 2023. Resultados Nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública, 2024.

Cutoffs:	WHO

Overweight/obesity by region


Adults, 2023



Survey type:	Measured
Age:	20+
Sample size:	5474
Area covered:	National
References:	Shamah-Levy T, Lazcano-Ponce EC, Cuevas-Nasu L, Romero-Martínez M, Gaona-Pineda EB, Gómez-Acosta LM, Mendoza-Alvarado LR, Méndez-Gómez-Humarán I. Encuesta Nacional de Salud y Nutrición Continua 2023. Resultados Nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública, 2024.

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Children, 2023



Survey type:	Measured
Age:	5-11
Sample size:	1413
Area covered:	National
References:	Shamah-Levy T, Lazcano-Ponce EC, Cuevas-Nasu L, Romero-Martínez M, Gaona-Pineda EB, Gómez-Acosta LM, Mendoza-Alvarado LR, Méndez-Gómez-Humarán I. Encuesta Nacional de Salud y Nutrición Continua 2023. Resultados Nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública, 2024.

Cutoffs:	WHO


Boys, 2023



Survey type:	Measured
Age:	5-11
Sample size:	1413
Area covered:	National
References:	Shamah-Levy T, Lazcano-Ponce EC, Cuevas-Nasu L, Romero-Martínez M, Gaona-Pineda EB, Gómez-Acosta LM, Mendoza-Alvarado LR, Méndez-Gómez-Humarán I. Encuesta Nacional de Salud y Nutrición Continua 2023. Resultados Nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública, 2024.

Cutoffs:	WHO


Girls, 2023



Survey type:	Measured
Age:	5-11
Sample size:	1413
Area covered:	National
References:	Shamah-Levy T, Lazcano-Ponce EC, Cuevas-Nasu L, Romero-Martínez M, Gaona-Pineda EB, Gómez-Acosta LM, Mendoza-Alvarado LR, Méndez-Gómez-Humarán I. Encuesta Nacional de Salud y Nutrición Continua 2023. Resultados Nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública, 2024.

Cutoffs:	WHO

0-5 years, 2022




Survey type:	Measured
Sample size:	10122251
Area covered:	National
References:	Encuesta Nacional de Salud y Nutrición Continua, Mexico 2022
Notes:	UNICEF/WHO/World Bank Joint Child Malnutrition Estimates Expanded Database: Overweight estimates: National and Disaggregated, July 2025, New York. Weighted sample size. For more information, please consult https://data.unicef.org/resources/jme/.

Definitions:	Percentage of children under 5 years of age falling above 2 standard deviations (moderate and severe) from the median weight-for-height of the reference population.
Cutoffs:	=>+2SD

Overweight/obesity by socio-economic group


Adults, 2022



Survey type:	Measured
Age:	20+
Sample size:	8563
Area covered:	National
References:	Campos-Nonato I, Galván-Valencia O,Hernández-Barrera L, Oviedo-Solís C, Barquera S.Prevalence of obesity and associated risk factors in Mexican adults: results of the Ensanut 2022.Salud Publica Mex. 2023;65(supl 1): S238-S247.https://doi.org/10.21149/14809. Available at https://saludpublica.mx/index.php/spm/article/view/14809/12424 (last accessed 11.12.23)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Children, 2020-2022



Survey type:	Measured
Sample size:	12371
Area covered:	National
References:	Shamah-Levy T, Gaona-Pineda EB, Cuevas-Nasu L, Morales-Ruan C, Valenzuela-Bravo DG, Méndez-Gómez Humaran I, Ávila-Arcos MA. Prevalence of overweight and obesity in the school-age and adolescent population in Mexico. Ensanut Continua 2020-2022. Salud Publica Mex [Internet]. June 14, 2023;65:s218-s224. Available from: https://saludpublica.mx/index.php/spm/article/view/14762 (Accessed 15.05.25)

Notes:	Age 5-11: 6,950, Age 12-19: 5,421
Definitions:	Socioeconomic status refers to welfare condition which is based on housing characteristics and goods and services available in the home, as in previous Ensanut surveys. Low = tertile 1, middle = tertile 2, high = tertile 3.
Cutoffs:	WHO 2007

0-5 years, 2020




Sample size:	9891242
References:	Other: Encuesta Nacional de Salud y Nutrición Continua COVID-19, Mexico 2020
Notes:	UNICEF/WHO/World Bank Joint Child Malnutrition Estimates Expanded Database: Overweight (Survey Estimates), May 2023, New York. For more information about the methodology, please consult https://data.unicef.org/resources/jme-2023-country-consultations/ Percentage of children under 5 years of age falling above 2 standard deviations (moderate and severe) from the median weight-for-height of the reference population.

Definitions:	=>+2SD

Overweight/obesity by ethnicity

Ethnic groups are as defined by publication of origin and are not as defined by WOF. In some instances ethnicity is conflated with nationality and/or race.

Children, 2020-2024




Survey type:	Measured
Age:	5-11
Sample size:	10,979
Area covered:	National

References:	Shamah-Levy T, Gaona-Pineda EB, Cuevas-Nasu L, Méndez-Gómez-Humarán I, Rodríguez-Ramírez S, Ávila-Arcos MA, Morales-Ruan C, Valenzuela-Bravo DG. National and state prevalence of overweight and obesity in Mexican school-age and adolescent population, and its associated factors. Salud Publica Mex. 2025;67:609-621. https://doi.org/10.21149/17311 [Accessed 15.01.26]
Definitions:	Ethnicity of the household
Cutoffs:	WHO 2007

Double burden of underweight & obesity


Adults, 2022



Survey type:	Measured
Age:	20+
References:	NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population representative studies with 222 million children, adolescents, and adults. Lancet 2024; published online Feb 29. https://doi.org/10.1016/S0140-6736(23)02750-2.
Notes:	Age Standardised estimates

Definitions:	Combined prevalence of BMI<18.5 kg/m² and BMI>=30 kg/m² (double burden of underweight and obesity)


Children, 2022



Survey type:	Measured
Age:	5-19
References:	NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population representative studies with 222 million children, adolescents, and adults. Lancet 2024; published online Feb 29. https://doi.org/10.1016/S0140-6736(23)02750-2.
Notes:	Age standardised estimates

Definitions:	Combined prevalence of BMI < -2SD and BMI > 2SD (double burden of thinness and obesity)
Cutoffs:	BMI < -2SD and BMI > 2SD

0-5 years, 2022




Survey type:	Measured
Age:	0-5
References:	United Nations Children's Fund, Division of Data, Analysis, Planning and Monitoring (2025). Global UNICEF Global Databases: Overlapping Stunting, Wasting and Overweight (Survey Estimates), 2025, New York. For more information, visit: https://data.unicef.org/topic/nutrition/malnutrition/ [Accessed 06.08.25]
Notes:	Encuesta Nacional de Salud y Nutrición Continua, Mexico 2022

Definitions:	Combined percentage of children under 5 years of age who are either wasted, stunted, or overweight (falling below -2 standard deviations from the median weight-for-height, falling below -2 standard deviations from the median height for age, or falling at or above +2 standard deviations from the median weight-for-height of the reference population)
Cutoffs:	WHZ <-2, HAZ <-2 and WHZ >+2

Insufficient physical activity


Adults, 2022



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.


Men, 2022



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.


Women, 2022



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.


Children, 2016



Survey type:	Self-reported
Age:	11-17
References:	Global Health Observatory data repository, World Health Organisation, https://apps.who.int/gho/data/node.main.A893ADO?lang=en (last accessed 16.03.21)
Notes:	% of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.

Definitions:	% Adolescents insufficiently active (age standardised estimate)


Boys, 2016



Survey type:	Self-reported
Age:	11-17
References:	Global Health Observatory data repository, World Health Organisation, https://apps.who.int/gho/data/node.main.A893ADO?lang=en (last accessed 16.03.21)
Notes:	% of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.

Definitions:	% Adolescents insufficiently active (age standardised estimate)

Girls, 2016




Survey type:	Self-reported
Age:	11-17
References:	Global Health Observatory data repository, World Health Organisation, https://apps.who.int/gho/data/node.main.A893ADO?lang=en (last accessed 16.03.21)
Notes:	% of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.

Definitions:	% Adolescents insufficiently active (age standardised estimate)

Average weekly frequency of fast food consumption

Adults, 2021-2024




Survey type:	Self-reported
Age:	15+
Sample size:	1003
Area covered:	National
References:	Global Diet Quality Project. 2024. "DQQ Results Dataset 2021-2024." Harvard Dataverse. https://doi.org/doi:10.7910/DVN/KY3W8A

Licence: CC BY-NC-SA 4.0. Accessed at dietquality.org on October 15, 2025.

Notes:	For more global dietary survey data, visit the Global Diet Quality Project website at https://www.dietquality.org/
Definitions:	Proportion of the population who consumed fast food during the previous day or night (self-reported)

Men, 2021-2024




Survey type:	Self-reported
Age:	15+
Sample size:	1003
Area covered:	National
References:	Global Diet Quality Project. 2024. "DQQ Results Dataset 2021-2024." Harvard Dataverse. https://doi.org/doi:10.7910/DVN/KY3W8A

Licence: CC BY-NC-SA 4.0. Accessed at dietquality.org on October 15, 2025.

Notes:	For more global dietary survey data, visit the Global Diet Quality Project website at https://www.dietquality.org/
Definitions:	Proportion of the population who consumed fast food during the previous day or night (self-reported)

Women, 2021-2024




Survey type:	Self-reported
Age:	15+
Sample size:	1003
Area covered:	National
References:	Global Diet Quality Project. 2024. "DQQ Results Dataset 2021-2024." Harvard Dataverse. https://doi.org/doi:10.7910/DVN/KY3W8A Licence: CC BY-NC-SA 4.0. Accessed at dietquality.org on October 15, 2025.

Notes:	For more global dietary survey data, visit the Global Diet Quality Project website at https://www.dietquality.org/
Definitions:	Proportion of the population who consumed fast food during the previous day or night (self-reported)


Mental health - depression disorders

Adults, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)

Men, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)

Women, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)

Children, 2023




Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (under 20 years of age)

Boys, 2023



Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (under 20 years of age)


Girls, 2023



Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (under 20 years of age)


Mental health - anxiety disorders

Adults, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)

Men, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)

Women, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)

Children, 2023




Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (under 20 years of age)

Boys, 2023



Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (under 20 years of age)


Girls, 2023



Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (under 20 years of age)

% Infants exclusively breastfed 0-5 months


0-5 years, 2011-2024



Area covered:	National
References:	Encuesta Nacional de Salud y Nutrición Continua, Mexico 2023
Notes:	Full details available at source: United Nations Children's Fund, Division of Data, Analysis, Planning and Monitoring (2025). Global UNICEF Global Databases: Infant and Young Child Feeding: Exclusive breastfeeding, New York, November 2025.
Definitions:	Infants 0-5 months of age who received only breast milk during the previous day. The definition allows for medicines, vitamins and oral re-hydration solution to have been given. These infants would not have received water-based liquids, milk based-liquids or food.

Percent of population who cannot afford a healthy diet


Adults, 2025



Area covered:	National
References:	FAO. 2026. FAOSTAT: Cost and Affordability of a Healthy Diet (CoAHD). https://www.fao.org/faostat/en/#data/CAHD. Licence: CC-BY-4.0. [Last Accessed 12.10.26]
Notes:	The percent of the population whose food budget is below the cost of a healthy diet.


Oesophageal cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Breast cancer


Women, 2022



Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Colorectal cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Pancreatic cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Gallbladder cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Kidney cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Cancer of the uterus


Women, 2022



Age:	20+
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Raised blood pressure

Adults, 2019




Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.

Men, 2019



Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.


Women, 2019



Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.


Raised cholesterol

Adults, 2008




References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).

Men, 2008



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).


Women, 2008



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).


Raised fasting blood glucose

Men, 2014



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A869?lang=en
Definitions:	Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication).


Women, 2014



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A869?lang=en
Definitions:	Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication).

Diabetes prevalence


Adults, 2024



References: Reproduced with kind permission International Diabetes Federation. IDF Diabetes Atlas, 11th edn. Brussels, Belgium:International Diabetes Federation, 2025. http://www.diabetesatlas.org

Ovarian Cancer


Women, 2022



Age:	20+
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Leukemia


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Liver and intrahepatic bile duct Cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Multiple Myeloma


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Non Hodgkin Lymphoma


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Thyroid Cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

PDF created on November 4, 2026