



Mongolia



Country report card


This report card contains the latest data available on the Global Obesity Observatory on overweight and obesity . Where available, data on common and relevant obesity drivers and comorbidities are also presented.

View the latest version of this report on the Global Obesity Observatory at <https://data.worldobesity.org/country/mongolia-143/>.

Contents	Page
Obesity prevalence	3
Trend: % Adults living with obesity in Mongolia 2005-2019	6
Trend: % Adults living with overweight or obesity in Mongolia 2005-2019	8
Overweight/obesity by education	10
Overweight/obesity by age	13
Overweight/obesity by region	16
Overweight/obesity by socio-economic group	21
Overweight/obesity by ethnicity	24
Double burden of underweight & overweight	27
Insufficient physical activity	30
Average daily frequency of carbonated soft drink consumption	36
Estimated per capita fruit intake	37
Prevalence of less than daily fruit consumption	38
Prevalence of less than daily vegetable consumption	39
Average weekly frequency of fast food consumption	40
Estimated per-capita processed meat intake	41
Estimated per capita whole grains intake	42
Mental health - depression disorders	43
Mental health - anxiety disorders	49
% Infants exclusively breastfed 0-5 months	54
Percent of population who cannot afford a healthy diet	55
Oesophageal cancer	56
Breast cancer	58
Colorectal cancer	59
Pancreatic cancer	61
Gallbladder cancer	63
Kidney cancer	65
Cancer of the uterus	67
Raised blood pressure	68
Raised cholesterol	71
Raised fasting blood glucose	74
Diabetes prevalence	76
Ovarian Cancer	77
Leukemia	78
Liver and intrahepatic bile duct Cancer	80
Multiple Myeloma	82
Non Hodgkin Lymphoma	84
Thyroid Cancer	86


Obesity prevalence

Adults, 2019




Survey type:	Measured
Age:	15-69
Sample size:	6654
Area covered:	National
References:	Fourth national STEPS Survey on the Prevalence of Noncommunicable Disease and Injury Risk Factors 2019. Brief Summary. Available at https://cdn.who.int/media/docs/default-source/ncds/ncd-surveillance/data-reporting/mongolia/mongolia-steps-survey---2019-brief-summary-english.pdf?sfvrsn=5ba7a1d3_1&download=true (last accessed 16.09.21)
Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m ² , obesity refers to a BMI greater than 30kg/m ² .	

Children, 2023



Survey type:	Measured
Age:	13-17
Sample size:	3311
Area covered:	National
References:	Global School-Based Student Health Survey, Mongolia 2023: Youth's Health Behavior Among Secondary School Children, GSHS-2023. Ulaanbaatar; 2024. Available at: https://www.who.int/publications/m/item/2021-gshs-country-report-mongolia (Accessed: 13.06.25)
Cutoffs:	WHO


0-5 years, 2023



Survey type:	Measured
Age:	0-5
Sample size:	5117
Area covered:	National
References:	UNICEF (2023) MONGOLIA Social Indicator Sample Survey-2023 KEY INDICATORS REPORT. Available at (https://mics.unicef.org/sites/mics/files/2024-07/Mongolia%202023%20SISS-MICS%20Key%20Indicators%20Report_English.pdf last accessed 11.06.25)
Cutoffs:	>2SD <3 SD, =>3SD

% Adults living with obesity in Mongolia 2005-2019

Men



Survey type:

Measured


References:

2005: Mongolian STEPS Survey on the prevalence of noncommunicable disease risk factors 2006
2009: Mongolian STEPS Survey on the Prevalence of Non-communicable Disease and Injury Risk Factors - 2009
2013: Chimeddamba O, Gearon E, Stevenson C et al. Trends in Adult Overweight and Obesity Prevalence in Mongolica 2005-2013. Obesity 2016;24(10):2194-2201
2019: Fourth national STEPS Survey on the Prevalence of Noncommunicable Disease and Injury Risk Factors 2019. Brief Summary. Available at https://cdn.who.int/media/docs/default-source/ncds/ncd-surveillance/data-reporting/mongolia/mongolia-steps-survey---2019_brief-summary_english.pdf?sfvrsn=5ba7a1d3_1&download=true (last accessed 16.09.21)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women



Survey type:

Measured

References:


2005: Mongolian STEPS Survey on the prevalence of noncommunicable disease risk factors 2006
2009: Mongolian STEPS Survey on the Prevalence of Non-communicable Disease and Injury Risk Factors - 2009
2013: Chimeddamba O, Gearon E, Stevenson C et al. Trends in Adult Overweight and Obesity Prevalence in Mongolica 2005-2013. Obesity 2016;24(10):2194-2201
2019: Fourth national STEPS Survey on the Prevalence of Noncommunicable Disease and Injury Risk Factors 2019. Brief Summary. Available at https://cdn.who.int/media/docs/default-source/ncds/ncd-surveillance/data-reporting/mongolia/mongolia-steps-survey---2019_brief-summary_english.pdf?sfvrsn=5ba7a1d3_1&download=true (last accessed 16.09.21)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with overweight or obesity in Mongolia 2005-2019

Men



Survey type:

Measured


References:

2005: Mongolian STEPS Survey on the prevalence of noncommunicable disease risk factors 2006
 2009: Mongolian STEPS Survey on the Prevalence of Non-communicable Disease and Injury Risk Factors - 2009
 2013: Chimeddamba O, Gearon E, Stevenson C et al. Trends in Adult Overweight and Obesity Prevalence in Mongolica 2005-2013. Obesity 2016;24(10):2194-2201
 2019: Fourth national STEPS Survey on the Prevalence of Noncommunicable Disease and Injury Risk Factors 2019. Brief Summary. Available at https://cdn.who.int/media/docs/default-source/ncds/ncd-surveillance/data-reporting/mongolia/mongolia-steps-survey---2019_brief-summary_english.pdf?sfvrsn=5ba7a1d3_1&download=true (last accessed 16.09.21)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women



Survey type:

Measured

References:


2005: Mongolian STEPS Survey on the prevalence of noncommunicable disease risk factors 2006
2009: Mongolian STEPS Survey on the Prevalence of Non-communicable Disease and Injury Risk Factors - 2009
2013: Chimeddamba O, Gearon E, Stevenson C et al. Trends in Adult Overweight and Obesity Prevalence in Mongolica 2005-2013. Obesity 2016;24(10):2194-2201
2019: Fourth national STEPS Survey on the Prevalence of Noncommunicable Disease and Injury Risk Factors 2019. Brief Summary. Available at https://cdn.who.int/media/docs/default-source/ncds/ncd-surveillance/data-reporting/mongolia/mongolia-steps-survey---2019_brief-summary_english.pdf?sfvrsn=5ba7a1d3_1&download=true (last accessed 16.09.21)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.


Overweight/obesity by education

Men, 2017



Survey type:	Measured
Age:	15-49
Sample size:	3314
Area covered:	National
References:	National Centre for Public Health (2017). Nutrition status of the population of Mongolia. Fifth National Nutrition Survey. Ulaanbaatar, Mongolia. Available at https://www.unicef.org/mongolia/reports/fifth-national-nutrition-survey-report (accessed 20.04.2020)
Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m ² , obesity refers to a BMI greater than 30kg/m ² .	

Women, 2017



Survey type: Measured

Age: 15-49


Sample size: 3314

Area covered: National

References: National Centre for Public Health (2017). Nutrition status of the population of Mongolia. Fifth National Nutrition Survey. Ulaanbaatar, Mongolia. Available at <https://www.unicef.org/mongolia/reports/fifth-national-nutrition-survey-report> (accessed 20.04.2020)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

0-5 years, 2018



Survey type: Measured

Sample size: 5,882

Area covered: National

References: NSO. 2019. Social Indicator Sample Survey-2018, Survey Findings Report. Ulaanbaatar, Mongolia: National Statistical Office of Mongolia. <https://washdata.org/sites/default/files/documents/reports/2019-10/Mongolia-2018-MICS-report.pdf>


Notes: Mother's education

Definitions: Weight for height. Overweight = Between +2SD and +3SD. Obesity = >+3SD

Cutoffs: Other

Overweight/obesity by age

Adults, 2017



Survey type: Measured


Sample size: 2,578

Area covered: National

References: Nutrition Status of the Population of Mongolia: Fifth National Nutrition Survey Report 2017. Available at: https://www.unicef.org/mongolia/media/1116/file/NNS_V_undsen_tailan_EN.pdf. Accessed 02.11.21.


Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Children, 2017-2018



Survey type:	Measured
Sample size:	1750
Area covered:	National
Definitions:	Nutrition Status of the Population of Mongolia: Fifth National Nutrition Survey Report 2017. Available at: https://www.unicef.org/mongolia/media/1116/file/NNS_V_undsen_tailan_EN.pdf . Accessed 02.11.21.
Cutoffs:	WHO


0-5 years, 2013



Survey type:	Measured
Sample size:	3404
Area covered:	National
References:	Mongolia MICS 2013 Survey - data analysed by R Jackson Leach, F Montague. World Obesity Federation
Notes:	IOTF monthly cut offs
Cutoffs:	IOTF

Overweight/obesity by region

Adults, 2019




Survey type:	Measured
Age:	15-69
Sample size:	6654
Area covered:	National

References: Fourth national STEPS Survey on the Prevalence of Noncommunicable Disease and Injury Risk Factors 2019. Brief Summary. Available at https://cdn.who.int/media/docs/default-source/ncds/ncd-surveillance/data-reporting/mongolia/mongolia-steps-survey---2019-brief-summary_english.pdf?sfvrsn=5ba7a1d3_1&download=true (last accessed 16.09.21)


Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Children, 2023




Survey type:	Measured
Age:	13-17
Sample size:	3311
Area covered:	National
References:	Global School-Based Student Health Survey, Mongolia 2023: Youth's Health Behavior Among Secondary School Children, GSHS-2023. Ulaanbaatar; 2024. Available at: https://www.who.int/publications/m/item/2021-gshs-country-report-mongolia (Accessed: 13.06.25)
Cutoffs:	WHO 2007

Boys, 2023




Survey type:	Measured
Age:	13-17
Sample size:	3311
Area covered:	National
References:	Global School-Based Student Health Survey, Mongolia 2023: Youth's Health Behavior Among Secondary School Children, GSHS-2023. Ulaanbaatar; 2024. Available at: https://www.who.int/publications/m/item/2021-gshs-country-report-mongolia (Accessed: 13.06.25)
Cutoffs:	WHO 2007

Girls, 2023



Survey type:	Measured
Age:	13-17
Sample size:	3311
Area covered:	National
References:	Global School-Based Student Health Survey, Mongolia 2023: Youth's Health Behavior Among Secondary School Children, GSHS-2023. Ulaanbaatar; 2024. Available at: https://www.who.int/publications/m/item/2021-gshs-country-report-mongolia (Accessed: 13.06.25)
Cutoffs:	WHO 2007


0-5 years, 2023



Survey type:	Measured
Sample size:	5117
Area covered:	National
References:	UNICEF (2023) MONGOLIA Social Indicator Sample Survey-2023 KEY INDICATORS REPORT. Available at (https://mics.unicef.org/sites/mics/files/2024-07/Mongolia%202023%20SISS-MICS%20Key%20Indicators%20Report_English.pdf last accessed 11.06.25)
Cutoffs:	>2SD <3 SD, =>3SD

Overweight/obesity by socio-economic group

Men, 2017



Survey type: Measured

Age: 15-49


Sample size: 3314

Area covered: National

References: National Centre for Public Health (2017). Nutrition status of the population of Mongolia. Fifth National Nutrition Survey. Ulaanbaatar, Mongolia. Available at <https://www.unicef.org/mongolia/reports/fifth-national-nutrition-survey-report> (accessed 20.04.2020)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2017



Survey type: Measured

Age: 15-49


Sample size: 3314

Area covered: National

References: National Centre for Public Health (2017). Nutrition status of the population of Mongolia. Fifth National Nutrition Survey. Ulaanbaatar, Mongolia. Available at <https://www.unicef.org/mongolia/reports/fifth-national-nutrition-survey-report> (accessed 20.04.2020)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

0-5 years, 2018




Survey type:	Measured
Sample size:	5882
Area covered:	National
References:	NSO. 2019. Social Indicator Sample Survey-2018, Survey Findings Report. Ulaanbaatar, Mongolia: National Statistical Office of Mongolia. https://washdata.org/sites/default/files/documents/reports/2019-10/Mongolia-2018-MICS-report.pdf (Accessed 24.06.21)
Notes:	Infants.
Definitions:	Weight for height. Overweight = Between +2SD and +3SD. Obesity = >+3SD
Cutoffs:	Other

Overweight/obesity by ethnicity

Ethnic groups are as defined by publication of origin and are not as defined by WOF. In some instances ethnicity is conflated with nationality and/or race.


Men, 2017



Survey type:	Measured
Age:	15-49
Sample size:	2,578
Area covered:	National
References:	Nutrition Status of the Population of Mongolia: Fifth National Nutrition Survey Report 2017. Available at: https://www.unicef.org/mongolia/media/1116/file/NNS_V_undsen_tailan_EN.pdf . Accessed 02.11.21.

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Women, 2017



Survey type:	Measured
Age:	15-49
Sample size:	2,578
Area covered:	National
References:	Nutrition Status of the Population of Mongolia: Fifth National Nutrition Survey Report 2017. Available at: https://www.unicef.org/mongolia/media/1116/file/NNS_V_undsen_tailan_EN.pdf . Accessed 02.11.21.

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


0-5 years, 2018



Survey type:	Measured
Sample size:	5,882
Area covered:	National
References:	NSO. 2019. Social Indicator Sample Survey-2018, Survey Findings Report. Ulaanbaatar, Mongolia: National Statistical Office of Mongolia. https://washdata.org/sites/default/files/documents/reports/2019-10/Mongolia-2018-MICS-report.pdf
Notes:	Ethnicity of household head
Definitions:	Weight for height. Overweight = Between +2SD and +3SD. Obesity = >+3SD
Cutoffs:	Other

Double burden of underweight & overweight

Adults, 2022



Survey type: Measured


Age: 20+

References: NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population representative studies with 222 million children, adolescents, and adults. Lancet 2024; published online Feb 29. [https://doi.org/10.1016/S0140-6736\(23\)02750-2](https://doi.org/10.1016/S0140-6736(23)02750-2).

Notes: Age Standardised estimates


Definitions: Combined prevalence of BMI<18.5 kg/m² and BMI≥30 kg/m² (double burden of underweight and obesity)

Children, 2022



Survey type:	Measured
Age:	5-19
References:	NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population representative studies with 222 million children, adolescents, and adults. Lancet 2024; published online Feb 29. https://doi.org/10.1016/S0140-6736(23)02750-2 .
Notes:	Age standardised estimates
Definitions:	Combined prevalence of BMI < -2SD and BMI > 2SD (double burden of thinness and obesity)
Cutoffs:	BMI < -2SD and BMI > 2SD


0-5 years, 2018-2018



Survey type:	Measured
Age:	0-5
References:	UNICEF data available at https://data.unicef.org/topic/nutrition/malnutrition/ (last accessed 12.03.25)
Notes:	Children under 5 falling below -2 standard deviations from the median height for age and falling at or above +2 standard deviations from the median weight-for-height of the reference population

Insufficient physical activity

Adults, 2022



Survey type: Self-reported


Age: 18+

Area covered: National

References: WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-\(age-standardized-estimate\)-\(-\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-))

Definitions: Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.

Men, 2022



Survey type: Self-reported


Age: 18+

Area covered: National

References: WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-\(age-standardized-estimate\)-\(-\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-))

Definitions: Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.

Women, 2022



Survey type: Self-reported


Age: 18+

Area covered: National

References: WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-\(age-standardized-estimate\)-\(-\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-))

Definitions: Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.

Children, 2016



Survey type: Self-reported


Age: 11-17

References: Global Health Observatory data repository, World Health Organisation, <https://apps.who.int/gho/data/node.main.A893ADO?lang=en> (last accessed 16.03.21)

Notes: % of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.

Definitions: % Adolescents insufficiently active (age standardised estimate)

Boys, 2016



Survey type: Self-reported


Age: 11-17

References: Global Health Observatory data repository, World Health Organisation, <https://apps.who.int/gho/data/node.main.A893ADO?lang=en> (last accessed 16.03.21)

Notes: % of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.

Definitions: % Adolescents insufficiently active (age standardised estimate)

Girls, 2016



Survey type: Self-reported

Age: 11-17


References: Global Health Observatory data repository, World Health Organisation, <https://apps.who.int/gho/data/node.main.A893ADO?lang=en> (last accessed 16.03.21)

Notes: % of school going adolescents not meeting WHO recommendations on Physical Activity for Health, i.e. doing less than 60 minutes of moderate- to vigorous-intensity physical activity daily.

Definitions: % Adolescents insufficiently active (age standardised estimate)

Average daily frequency of carbonated soft drink consumption

Children, 2010-2015




Survey type: Measured

Age: 12-17

References: Beal et al. (2019). Global Patterns of Adolescent Fruit, Vegetable, Carbonated Soft Drink, and Fast-food consumption: A meta-analysis of global school-based student health surveys. Food and Nutrition Bulletin. <https://doi.org/10.1177/0379572119848287> sourced from Food Systems Dashboard <http://www.foodsystemsdashboard.org/food-system>

Estimated per capita fruit intake

Adults, 2017



Survey type: Measured


Age: 25+

References: Global Burden of Disease, the Institute for Health Metrics and Evaluation <http://ghdx.healthdata.org/>

Definitions: Estimated per-capita fruit intake (g/day)

Prevalence of less than daily fruit consumption

Children, 2010-2015



Survey type: Measured


Age: 12-17

References: Global School-based Student Health Surveys. Beal et al (2019). Global Patterns of Adolescent Fruit, Vegetable, Carbonated Soft Drink, and Fast-food consumption: A meta-analysis of global school-based student health surveys. Food and Nutrition Bulletin. <https://doi.org/10.1177/0379572119848287>. Sourced from Food Systems Dashboard <http://www.foodsystemsdashboard.org/food-system>

Definitions: Prevalence of less-than-daily fruit consumption (% less-than-daily fruit consumption)

Prevalence of less than daily vegetable consumption

Children, 2010-2015



Survey type: Measured


Age: 12-17

References: Beal et al. (2019). Global Patterns of Adolescent Fruit, Vegetable, Carbonated Soft Drink, and Fast-food consumption: A meta-analysis of global school-based student health surveys. Food and Nutrition Bulletin. <https://doi.org/10.1177/0379572119848287> sourced from Food Systems Dashboard <http://www.foodsystemsdashboard.org/food-system>

Definitions: Prevalence of less-than-daily vegetable consumption (% less-than-daily vegetable consumption)

Average weekly frequency of fast food consumption

Children, 2010-2015




Age: 12-17

References: Beal et al. (2019). Global Patterns of Adolescent Fruit, Vegetable, Carbonated Soft Drink, and Fast-food consumption: A meta-analysis of global school-based student health surveys. Food and Nutrition Bulletin. <https://doi.org/10.1177/0379572119848287> sourced from Food Systems Dashboard <http://www.foodsystemsdashboard.org/food-system>

Estimated per-capita processed meat intake

Adults, 2017



Survey type: Measured


Age: 25+

References: Global Burden of Disease, the Institute for Health Metrics and Evaluation <http://ghdx.healthdata.org/>

Definitions: Estimated per-capita processed meat intake (g per day)

Estimated per capita whole grains intake

Adults, 2017



Survey type: Measured


Age: 25+

References: Global Burden of Disease, the Institute for Health Metrics and Evaluation <http://ghdx.healthdata.org/>

Definitions: Estimated per-capita whole grains intake (g/day)


Mental health - depression disorders

Adults, 2021




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from http://vizhub.healthdata.org/gbd-compare . (Last accessed 23.04.25)
Definitions:	Number living with depression per 100,000 population (adults 20+ years)

Men, 2021




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from http://vizhub.healthdata.org/gbd-compare . (Last accessed 23.04.25)
Definitions:	Number living with depression per 100,000 population (adults 20+ years)

Women, 2021



Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from http://vizhub.healthdata.org/gbd-compare . (Last accessed 23.04.25)
Definitions:	Number living with depression per 100,000 population (adults 20+ years)

Children, 2021



Area covered:

National


References:

Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from <http://vizhub.healthdata.org/gbd-compare>. (Last accessed 23.04.25)

Definitions:

Number living with depressive disorder per 100,000 population (Under 20 years of age)

Boys, 2021



Area covered:

National


References:

Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from <http://vizhub.healthdata.org/gbd-compare>. (Last accessed 23.04.25)

Definitions:

Number living with depressive disorder per 100,000 population (Under 20 years of age)

Girls, 2021



Area covered:

National

References:


Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from <http://vizhub.healthdata.org/gbd-compare>. (Last accessed 23.04.25)

Definitions:

Number living with depressive disorder per 100,000 population (Under 20 years of age)

Mental health - anxiety disorders

Adults, 2021




Age: 20+

References: Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from <http://vizhub.healthdata.org/gbd-compare>. (Last accessed 23.04.25).

Definitions: Number living with anxiety per 100,000 population

Men, 2021




Age: 20+

References: Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from <http://vizhub.healthdata.org/gbd-compare>. (Last accessed 23.04.25).

Definitions: Number living with anxiety per 100,000 population

Women, 2021




Age: 20+

References: Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from <http://vizhub.healthdata.org/gbd-compare>. (Last accessed 23.04.25).

Definitions: Number living with anxiety per 100,000 population


Children, 2021



References:

Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from <http://vizhub.healthdata.org/gbd-compare>. (Last accessed 23.04.25)


Boys, 2021



References:

Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from <http://vizhub.healthdata.org/gbd-compare>. (Last accessed 23.04.25)

Girls, 2021




References:

Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2021. Seattle, WA: IHME, University of Washington, 2023. Available from <http://vizhub.healthdata.org/gbd-compare>. (Last accessed 23.04.25)

% Infants exclusively breastfed 0-5 months

0-5 years, 2007-2023



References:


Mongolia Multiple Indicator Cluster Survey 2023

Notes:

Full details are available. Original citation: United Nations Children's Fund, Division of Data, Analysis, Planning and Monitoring (2024).
Global UNICEF Global Databases: Infant and Young Child Feeding: Exclusive breastfeeding, New York, December 2024.

Percent of population who cannot afford a healthy diet

Adults, 2022



Area covered:


National

References:

The Food Systems Dashboard. The Global Alliance for Improved Nutrition (GAIN), The Columbia Climate School, and Cornell University College of Agriculture and Life Sciences. 2024. Geneva, Switzerland. <https://www.foodsystemsdashboard.org>. DOI: <https://doi.org/10.36072/db>.


Oesophageal cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Indicence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Indicence per 100,000

Breast cancer

Women, 2022



Area covered:

National

References:


Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions:

Incidence per 100,000


Colorectal cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Incidence per 100,000


Pancreatic cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Incidence per 100,000


Gallbladder cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Indicence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Indicence per 100,000


Kidney cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today , accessed [16.07.24]
Definitions:	Incidence per 100,000

Cancer of the uterus

Women, 2022




Age: 20+

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Raised blood pressure


Adults, 2015



References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A875?lang=en>

Definitions: Age Standardised estimated % Raised blood pressure 2015 (SBP $\geq$ 140 OR DBP $\geq$ 90).


Men, 2015



References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A875?lang=en>

Definitions: Age Standardised estimated % Raised blood pressure 2015 (SBP $\geq$ 140 OR DBP $\geq$ 90).

Women, 2015




References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A875?lang=en>

Definitions: Age Standardised estimated % Raised blood pressure 2015 (SBP $\geq$ 140 OR DBP $\geq$ 90).

Raised cholesterol


Adults, 2008



References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A885>

Definitions: % Raised total cholesterol (≥ 5.0 mmol/L) (age-standardized estimate).


Men, 2008



References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A885>

Definitions: % Raised total cholesterol (≥ 5.0 mmol/L) (age-standardized estimate).

Women, 2008




References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A885>

Definitions: % Raised total cholesterol (≥ 5.0 mmol/L) (age-standardized estimate).

Raised fasting blood glucose


Men, 2014



References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A869?lang=en>

Definitions: Age Standardised % raised fasting blood glucose (≥ 7.0 mmol/L or on medication).

Women, 2014




References: Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A869?lang=en>

Definitions: Age Standardised % raised fasting blood glucose (≥ 7.0 mmol/L or on medication).

Diabetes prevalence

Adults, 2024




References:

Reproduced with kind permission International Diabetes Federation. IDF Diabetes Atlas, 11th edn. Brussels, Belgium:International Diabetes Federation, 2025. <http://www.diabetesatlas.org>

Ovarian Cancer

Women, 2022




Age: 20+

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Leukemia

Men, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022



Age: 20+

Area covered: National


References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Liver and intrahepatic bile duct Cancer

Men, 2022




Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Women, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Multiple Myeloma

Men, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022



Age: 20+

Area covered: National


References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Non Hodgkin Lymphoma

Men, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022



Age: 20+

Area covered: National


References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Thyroid Cancer

Men, 2022




Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Women, 2022



Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

PDF created on July 7, 2025