
Oman



Country report card - adults


This report card contains the latest data available on the Global Obesity Observatory on overweight and obesity for adults. Where available, data on common and relevant obesity drivers and comorbidities are also presented.

View the latest version of this report on the Global Obesity Observatory at https://data.worldobesity.org/country/oman-165/.

Contents	Page
Obesity prevalence	3
Trend: % Adults living with obesity in Oman 1991-2025	6
Trend: % Adults living with overweight or obesity in Oman 1991-2025	11
Trend: % Adults living with obesity in selected countries in the EMRO Region 1984-2017, selected countries	16
Overweight/obesity by age	21
Overweight/obesity by socio-economic group	24
Overweight/obesity by ethnicity	27
Double burden of underweight & obesity	34
Insufficient physical activity	37
Mental health - depression disorders	44
Mental health - anxiety disorders	48
Oesophageal cancer	52
Breast cancer	55
Colorectal cancer	57
Pancreatic cancer	60
Gallbladder cancer	63
Kidney cancer	66
Cancer of the uterus	69
Raised blood pressure	71
Raised cholesterol	75
Raised fasting blood glucose	78
Diabetes prevalence	80
Ovarian Cancer	81
Leukemia	83
Liver and intrahepatic bile duct Cancer	88
Multiple Myeloma	91
Non Hodgkin Lymphoma	96
Thyroid Cancer	101

Obesity prevalence

Adults, 2025




Survey type:	Measured
Age:	18+
Sample size:	10,174
Area covered:	National
References:	Oman STEPS Survey 2025 Fact Sheet (National ncd survey 2025 - fact sheet (total)). Available at: https://moh.gov.om/en/statistics/studies-and-research/ [Accessed 07.04.26]

Notes:	Total population (Omani and Non-Omani)
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

% Adults living with obesity in Oman 1991-2025

Men




Survey type:	Measured
References:	1991: Al-Lawati JA, Jousilahti PJ. Prevalence and 10year secular trend of obesity in Oman. Saudi Med J 2004;25:346-351 2000: Al-Lawati JA and Jousilahti PJ. (2004). Prevalence and 10 year secular trend of obesity in Oman. Saudi Medical Journal 25: 346 - 351. 2008: Riyami AA, Elaty MAA, Morsi M et al Oman World Health Survey: Part 1 - Methodology, Sociodemographic Profile and Epidemiology of Non-Communicable Diseases in Oman. Oman Medical Journal 2012; 27(5):425-443 2017: Executive Summary. Oman National Non-Communicable Disease & their Risk Factors Survey 2017 https://mohcsr.gov.om/wp-content/uploads/2019/01/Executive-Summary_NCDsurvey2017_En.pdf (last accessed 17.10.19) 2025: Oman STEPS Survey 2025 Fact Sheet (National ncd survey 2025 - fact sheet (total)). Available at: https://moh.gov.om/en/statistics/studies-and-research/ [Accessed 07.04.26]

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women




Survey type:	Measured
References:	1991: Al-Lawati JA, Jousilahti PJ. Prevalence and 10year secular trend of obesity in Oman. Saudi Med J 2004;25:346-351 2000: Al-Lawati JA and Jousilahti PJ. (2004). Prevalence and 10 year secular trend of obesity in Oman. Saudi Medical Journal 25: 346 - 351. 2008: Riyami AA, Elaty MAA, Morsi M et al Oman World Health Survey: Part 1 - Methodology, Sociodemographic Profile and Epidemiology of Non-Communicable Diseases in Oman. Oman Medical Journal 2012; 27(5):425-443 2017: Executive Summary. Oman National Non-Communicable Disease & their Risk Factors Survey 2017 https://mohcsr.gov.om/wp-content/uploads/2019/01/Executive-Summary_NCDsurvey2017_En.pdf (last accessed 17.10.19) 2025: Oman STEPS Survey 2025 Fact Sheet (National ncd survey 2025 - fact sheet (total)). Available at: https://moh.gov.om/en/statistics/studies-and-research/ [Accessed 07.04.26]

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with overweight or obesity in Oman 1991-2025

Men




Survey type:	Measured
References:	1991: Al-Lawati JA, Jousilahti PJ. Prevalence and 10year secular trend of obesity in Oman. Saudi Med J 2004;25:346-351 2000: Al-Lawati JA and Jousilahti PJ. (2004). Prevalence and 10 year secular trend of obesity in Oman. Saudi Medical Journal 25: 346 - 351. 2008: Riyami AA, Elaty MAA, Morsi M et al Oman World Health Survey: Part 1 - Methodology, Sociodemographic Profile and Epidemiology of Non-Communicable Diseases in Oman. Oman Medical Journal 2012; 27(5):425-443 2017: Executive Summary. Oman National Non-Communicable Disease & their Risk Factors Survey 2017 https://mohcsr.gov.om/wp-content/uploads/2019/01/Executive-Summary_NCDsurvey2017_En.pdf (last accessed 17.10.19) 2025: Oman STEPS Survey 2025 Fact Sheet (National ncd survey 2025 - fact sheet (total)). Available at: https://moh.gov.om/en/statistics/studies-and-research/ [Accessed 07.04.26]

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women




Survey type:	Measured
References:	1991: Al-Lawati JA, Jousilahti PJ. Prevalence and 10year secular trend of obesity in Oman. Saudi Med J 2004;25:346-351 2000: Al-Lawati JA and Jousilahti PJ. (2004). Prevalence and 10 year secular trend of obesity in Oman. Saudi Medical Journal 25: 346 - 351. 2008: Riyami AA, Elaty MAA, Morsi M et al Oman World Health Survey: Part 1 - Methodology, Sociodemographic Profile and Epidemiology of Non-Communicable Diseases in Oman. Oman Medical Journal 2012; 27(5):425-443 2017: Executive Summary. Oman National Non-Communicable Disease & their Risk Factors Survey 2017 https://mohcsr.gov.om/wp-content/uploads/2019/01/Executive-Summary_NCDsurvey2017_En.pdf (last accessed 17.10.19) 2025: Oman STEPS Survey 2025 Fact Sheet (National ncd survey 2025 - fact sheet (total)). Available at: https://moh.gov.om/en/statistics/studies-and-research/ [Accessed 07.04.26]

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with obesity in selected countries in the EMRO Region 1984-2017, selected countries

Women




References:

1984, 1998: Benjelloun S. Nutrition transition in Morocco. Public Health Nutrition. 2002;5(1A):135
1991: Al-Lawati JA, Jousilahti PJ. Prevalence and 10year secular trend of obesity in Oman. Saudi Med J 2004;25:346
1997: Sibai AM, Hwalla N, Adra N, Rahal B. Prevalence and Covariates of Obesity in Lebanon: Findings from the First Epidemiolgical Study. Ob
Research 203;11:1353-
2000: Al-Lawati JA and Jousilahti PJ. (2004). Prevalence and 10 year secular trend of obesity in Oman. Saudi Medical Journal 25: 346 -
2002: WHO Infobase, Macro international
2007: Department of Statistics [Jordan] and Macro International. 2008. Jordan Population and Family Health Survey 2007: Key Findings. Calve
Maryland, USA: Department of Statistics and Macro Internati
2008: Riyami AA, Elaty MAA, Morsi M et al Oman World Health Survey: Part 1 - Methodology, Sociodemographic Profile and Epidemiology of
Communicable Diseases in Oman. Oman Medical Journal 2012; 27(5):425
2009: WHO STEPS Report 2009 Leb

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data so
for methodologies

Men




References:

1984, 1998: Benjelloun S. Nutrition transition in Morocco. Public Health Nutrition. 2002;5(1A):135
1991: Al-Lawati JA, Jousilahti PJ. Prevalence and 10year secular trend of obesity in Oman. Saudi Med J 2004;25:346
1997: Sibai AM, Hwalla N, Adra N, Rahal B. Prevalence and Covariates of Obesity in Lebanon: Findings from the First Epidemiolgical Study. Ob
Research 203;11:1353-
2000: Al-Lawati JA and Jousilahti PJ. (2004). Prevalence and 10 year secular trend of obesity in Oman. Saudi Medical Journal 25: 346 -
2002: WHO Infobase, Macro international
2007: Department of Statistics [Jordan] and Macro International. 2008. Jordan Population and Family Health Survey 2007: Key Findings. Calve
Maryland, USA: Department of Statistics and Macro Internati
2008: Riyami AA, Elaty MAA, Morsi M et al Oman World Health Survey: Part 1 - Methodology, Sociodemographic Profile and Epidemiology of
Communicable Diseases in Oman. Oman Medical Journal 2012; 27(5):425
2009: WHO STEPS Report 2009 Leb

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data so
for methodologies

Overweight/obesity by age

Women, 2016-2017




Survey type:	Measured
Sample size:	4159
Area covered:	National
References:	Oman National Nutrition Survey 2016-2017 http://groundworkhealth.org/wp-content/uploads/2020/04/ONNS_Report_2017.pdf (Accessed 04.09.20)
Notes:	Omani citizens only.

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Overweight/obesity by socio-economic group

Women, 2016-2017




Survey type:	Measured
Age:	15-49
Sample size:	4159
Area covered:	National
References:	Oman National Nutrition Survey 2016-2017 http://groundworkhealth.org/wp-content/uploads/2020/04/ONNS_Report_2017.pdf (Accessed 04.09.20)

Notes:	Omani citizens only.
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Overweight/obesity by ethnicity

Ethnic groups are as defined by publication of origin and are not as defined by WOF. In some instances ethnicity is conflated with nationality and/or race.


Adults, 2025



Survey type:	Measured
Age:	18+
Sample size:	10,174
Area covered:	National
References:	Oman STEPS Survey 2025 Fact Sheets (National ncd survey 2025 - fact sheet (omani) and National ncd survey 2025 - fact sheet (non-omani)). Available at: https://moh.gov.om/en/statistics/studies-and-research/ [Accessed 07.04.26]

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Men, 2025



Survey type:	Measured
Age:	18+
Sample size:	10,174
Area covered:	National
References:	Oman STEPS Survey 2025 Fact Sheets (National ncd survey 2025 - fact sheet (omani) and National ncd survey 2025 - fact sheet (non-omani)). Available at: https://moh.gov.om/en/statistics/studies-and-research/ [Accessed 07.04.26]

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2025




Survey type:	Measured
Age:	18+
Sample size:	10,174
Area covered:	National
References:	Oman STEPS Survey 2025 Fact Sheets (National ncd survey 2025 - fact sheet (omani) and National ncd survey 2025 - fact sheet (non-omani)). Available at: https://moh.gov.om/en/statistics/studies-and-research/ [Accessed 07.04.26]

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Double burden of underweight & obesity

Adults, 2022




Survey type:	Measured
Age:	20+
References:	NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population representative studies with 222 million children, adolescents, and adults. Lancet 2024; published online Feb 29. https://doi.org/10.1016/S0140-6736(23)02750-2.
Notes:	Age Standardised estimates

Definitions:	Combined prevalence of BMI<18.5 kg/m² and BMI>=30 kg/m² (double burden of underweight and obesity)

Insufficient physical activity


Adults, 2022



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.


Men, 2022



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.

Women, 2022




Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.


Mental health - depression disorders

Adults, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)

Men, 2023



Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)


Women, 2023



Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)


Mental health - anxiety disorders

Adults, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)

Men, 2023



Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)


Women, 2023



Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)


Oesophageal cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Breast cancer


Women, 2022



Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Colorectal cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Pancreatic cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Gallbladder cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Kidney cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Cancer of the uterus


Women, 2022



Age:	20+
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Raised blood pressure

Adults, 2019




Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.

Men, 2019



Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.


Women, 2019



Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.


Raised cholesterol

Adults, 2008




References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).

Men, 2008



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).


Women, 2008



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).


Raised fasting blood glucose

Men, 2014



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A869?lang=en
Definitions:	Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication).


Women, 2014



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A869?lang=en
Definitions:	Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication).

Diabetes prevalence


Adults, 2024



References: Reproduced with kind permission International Diabetes Federation. IDF Diabetes Atlas, 11th edn. Brussels, Belgium:International Diabetes Federation, 2025. http://www.diabetesatlas.org

Ovarian Cancer


Women, 2022



Age:	20+
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Leukemia


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Liver and intrahepatic bile duct Cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Multiple Myeloma


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Non Hodgkin Lymphoma


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Thyroid Cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

PDF created on November 3, 2026