




# Peru



## Country report card - adults


*This report card contains the latest data available on the Global Obesity Observatory on overweight and obesity for adults. Where available, data on common and relevant obesity drivers and comorbidities are also presented.*

*View the latest version of this report on the Global Obesity Observatory at <https://data.worldobesity.org/country/peru-171/>*

Contents	Page
Obesity prevalence	3
Trend: % Adults (15+) living with obesity by education level	4
Trend: % Adults (15+) living with obesity, 2013-2024	5
Trend: % Adults (15+) living with overweight or obesity, 2013-2024	8
Trend: % Adults living with obesity by ethnic self-identification	11
Trend: % Adults living with obesity by region	12
Trend: % Adults living with obesity by wealth quintile	13
Trend: % Adults living with obesity in Peru 1992-2014	14
Trend: % Adults living with overweight or obesity in Peru 1992-2014	15
Trend: % Adults living with obesity in selected countries in the Americas Region 1960-2018, selected countries	16
Overweight/obesity by education	21
Overweight/obesity by age	22
Overweight/obesity by region	23
Overweight/obesity by socio-economic group	24
Overweight/obesity by ethnicity	25
Double burden of underweight & overweight	26
Insufficient physical activity	27
Mental health - depression disorders	30
Mental health - anxiety disorders	33
Percent of population who cannot afford a healthy diet	36
Oesophageal cancer	37
Breast cancer	39
Colorectal cancer	40
Pancreatic cancer	42
Gallbladder cancer	44
Kidney cancer	46
Cancer of the uterus	48
Raised blood pressure	49
Raised cholesterol	52
Raised fasting blood glucose	55
Diabetes prevalence	57
Ovarian Cancer	58
Leukemia	59
Liver and intrahepatic bile duct Cancer	61
Multiple Myeloma	63
Non Hodgkin Lymphoma	65
Thyroid Cancer	67

## Obesity prevalence

### Adults, 2024



Survey type:	Measured
Age:	15+
Sample size:	30,631
Area covered:	National


**References:** Instituto Nacional de Estadística e Informática. Perú: Enfermedades No Transmisibles y Transmisibles 2024. [https://proyectos.inei.gob.pe/endes/2024/SALUD/ENFERMEDADES\\_ENDES\\_2024.pdf](https://proyectos.inei.gob.pe/endes/2024/SALUD/ENFERMEDADES_ENDES_2024.pdf) [Accessed 01.10.25]

**Notes:** ENDES 2024. Standard overweight and obesity cutoffs used except in adolescents aged 15-17 where they are measured using WHO cutoffs, and for adults aged 60 and over, overweight refers to BMI between 28kg/m<sup>2</sup> and 31.9kg/m<sup>2</sup> and obesity refers to ≥32kg/m<sup>2</sup>. Interpret with caution when comparing cross-nationally.

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.*

## % Adults (15+) living with obesity by education level

### Men and women



Survey type: Measured


References: Instituto Nacional de Estadística e Informática. Perú: Enfermedades No Transmisibles y Transmisibles 2014-2024. <https://proyectos.inei.gob.pe/endes/>

Notes: ENDES. Standard overweight and obesity cutoffs used except in adolescents aged 15-17 where they are measured using WHO cutoffs, and for adults aged 60 and over, overweight refers to BMI between 28kg/m<sup>2</sup> and 31.9kg/m<sup>2</sup> and obesity refers to ≥32 kg/m<sup>2</sup>. Interpret with caution when comparing cross-nationally.

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

## % Adults (15+) living with obesity, 2013-2024

### Women



Survey type: Measured


References: Instituto Nacional de Estadística e Informática. Perú: Enfermedades No Transmisibles y Transmisibles 2014-2024. <https://proyectos.inei.gob.pe/endes/>

Notes: ENDES. Standard obesity cutoffs used except in adolescents aged 15-17 where they are measured using WHO cutoffs, and for adults aged 60 and over, obesity refers to  $\geq 32$  kg/m<sup>2</sup>. Interpret with caution when comparing cross-nationally.

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.*

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

## Men and women



Survey type: Measured


References: Instituto Nacional de Estadística e Informática. Perú: Enfermedades No Transmisibles y Transmisibles 2014-2024. <https://proyectos.inei.gob.pe/endes/>

Notes: ENDES. Standard obesity cutoffs used except in adolescents aged 15-17 where they are measured using WHO cutoffs, and for adults aged 60 and over, obesity refers to  $\geq 32$  kg/m<sup>2</sup>. Interpret with caution when comparing cross-nationally.

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.*

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

## Men



Survey type: Measured

References: Instituto Nacional de Estadística e Informática. Perú: Enfermedades No Transmisibles y Transmisibles 2014-2024. <https://proyectos.inei.gob.pe/endes/>


Notes: ENDES. Standard obesity cutoffs used except in adolescents aged 15-17 where they are measured using WHO cutoffs, and for adults aged 60 and over, obesity refers to  $\geq 32 \text{ kg/m}^2$ . Interpret with caution when comparing cross-nationally.

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.*

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

## % Adults (15+) living with overweight or obesity, 2013-2024

### Women



Survey type: Measured


References: Instituto Nacional de Estadística e Informática. Perú: Enfermedades No Transmisibles y Transmisibles 2014-2024. <https://proyectos.inei.gob.pe/endes/>

Notes: ENDES. Standard overweight and obesity cutoffs used except in adolescents aged 15-17 where they are measured using WHO cutoffs, and for adults aged 60 and over, overweight refers to BMI between 28kg/m<sup>2</sup> and 31.9kg/m<sup>2</sup> and obesity refers to ≥32 kg/m<sup>2</sup>. Interpret with caution when comparing cross-nationally.

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.*

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

## Men and women



Survey type: Measured


References: Instituto Nacional de Estadística e Informática. Perú: Enfermedades No Transmisibles y Transmisibles 2014-2024. <https://proyectos.inei.gob.pe/endes/>

Notes: ENDES. Standard overweight and obesity cutoffs used except in adolescents aged 15-17 where they are measured using WHO cutoffs, and for adults aged 60 and over, overweight refers to BMI between 28kg/m<sup>2</sup> and 31.9kg/m<sup>2</sup> and obesity refers to ≥32 kg/m<sup>2</sup>. Interpret with caution when comparing cross-nationally.

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.*

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

## Men



**Survey type:** Measured

**References:** Instituto Nacional de Estadística e Informática. Perú: Enfermedades No Transmisibles y Transmisibles 2014-2024. <https://proyectos.inei.gob.pe/endes/>


**Notes:** ENDES. Standard overweight and obesity cutoffs used except in adolescents aged 15-17 where they are measured using WHO cutoffs, and for adults aged 60 and over, overweight refers to BMI between 28kg/m<sup>2</sup> and 31.9kg/m<sup>2</sup> and obesity refers to ≥32 kg/m<sup>2</sup>. Interpret with caution when comparing cross-nationally.

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.*

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

## % Adults living with obesity by ethnic self-identification

### Men and women



Survey type: Measured

References: Instituto Nacional de Estadística e Informática. Perú: Enfermedades No Transmisibles y Transmisibles 2019-2024. <https://proyectos.inei.gob.pe/endes/>


Definitions: ENDES. Standard overweight and obesity cutoffs used except in adolescents aged 15-17 where they are measured using WHO cutoffs, and for adults aged 60 and over, overweight refers to BMI between 28kg/m<sup>2</sup> and 31.9kg/m<sup>2</sup> and obesity refers to ≥32 kg/m<sup>2</sup>. Interpret with caution when comparing cross-nationally.

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.*

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

## % Adults living with obesity by region

### Men and women



Survey type: Measured


References: Instituto Nacional de Estadística e Informática. Perú: Enfermedades No Transmisibles y Transmisibles 2014-2024. <https://proyectos.inei.gob.pe/endes/>

Notes: ENDES. Standard obesity cutoffs used except in adolescents aged 15-17 where they are measured using WHO cutoffs, and for adults aged 60 and over, obesity refers to BMI  $\geq 32$  kg/m<sup>2</sup>. Interpret with caution when comparing cross-nationally.

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

## % Adults living with obesity by wealth quintile

### Men and women




Survey type: Measured

References: Instituto Nacional de Estadística e Informática. Perú: Enfermedades No Transmisibles y Transmisibles 2014-2024. <https://proyectos.inei.gob.pe/endes/>

Definitions: ENDES. Standard overweight and obesity cutoffs used except in adolescents aged 15-17 where they are measured using WHO cutoffs, and for adults aged 60 and over, overweight refers to BMI between 28kg/m<sup>2</sup> and 31.9kg/m<sup>2</sup> and obesity refers to ≥32 g/m<sup>2</sup>. Interpret with caution when comparing cross-nationally.

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

## % Adults living with obesity in Peru 1992-2014



Survey type:

Measured

References:

1992: Martorell R, Khan LK, Hughes ML, Grummer Strawn LM. Obesity in women from developing countries. EJCN (2000) 54:247-252  
 2000: Instituto Nacional de Estadística e Informática - INEI/Perú and Macro International. 2001. Perú Encuesta Demográfica y de Salud Familiar 2000. Lima, Perú: INEI/Perú and Macro International.  
 2007: Instituto Nacional de Estadística e Informática - INEI/Peru and ORC Macro. 2009. Perú Encuesta Demográfica y de Salud Familiar - ENDES 2007-2008. Lima, Perú: INEI/Peru and ORC Macro.  
 2012: Instituto Nacional de Estadística e Informática - INEI/Perú. 2013. Perú Encuesta Demográfica y de Salud Familiar - ENDES 2012. Lima, Perú: INEI/Peru.  
 2014: Instituto Nacional de Estadística e Informática - INEI/Perú. 2015. Perú Encuesta Demográfica y de Salud Familiar - ENDES 2014. Lima, Perú: INEI/Perú.


Notes:

Adults aged 15 to 49

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.*

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

## % Adults living with overweight or obesity in Peru 1992-2014



**Survey type:**

Measured

**References:**

1992: Martorell R, Khan LK, Hughes ML, Grummer Strawn LM. Obesity in women from developing countries. EJCN (2000) 54:247-252  
 2000: Instituto Nacional de Estadística e Informática - INEI/Perú and Macro International. 2001. Perú Encuesta Demográfica y de Salud Familiar 2000. Lima, Perú: INEI/Perú and Macro International.  
 2007: Instituto Nacional de Estadística e Informática - INEI/Peru and ORC Macro. 2009. Perú Encuesta Demográfica y de Salud Familiar - ENDES 2007-2008. Lima, Perú: INEI/Peru and ORC Macro.  
 2012: Instituto Nacional de Estadística e Informática - INEI/Perú. 2013. Perú Encuesta Demográfica y de Salud Familiar - ENDES 2012. Lima, Perú: INEI/Peru.  
 2014: Instituto Nacional de Estadística e Informática - INEI/Perú. 2015. Perú Encuesta Demográfica y de Salud Familiar - ENDES 2014. Lima, Perú: INEI/Perú.

**Notes:**


Adults aged 15 to 49

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.*

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

**% Adults living with obesity in selected countries in the Americas Region  
1960-2018, selected countries**

## Men




### References:

- 1960, 1971, 1973, 1976, 1991: Flegal KM, Carroll MD, Kuczmarski RJ, Johnson CL. Overweight and obesity in the United States: prevalence and trends, 1960-1994. *International Journal of Obesity* (1998);22:39-47
- 1975: Monteiro CA, Conde WL, Popking BM. Is obesity replacing or adding to undernutrition? Evidence from different social classes in Brazil. 2002. *Public Health Nutrition*:51(1A), 105-112
- 1988: Berrios X, Jadue I, Zenteno J, Ross MI, Rodriguez H. Prevalencia de factores de riesgo para enfermedades cronicas. Estudio de la poblacion general de la region Metropolitana, 1986-1987. *Rev. Med. Chile*. 1990;118:597-604
- 1992, 1994, 1995: Martorell R, Khan LK, Hughes ML, Grummer Strawn LM. Obesity in women from developing countries. *EJCN* (2000) 54:247-252
- 1997: Filozof C, Gonzales C, Sereday M, Mazza C, Braguinsky J. Obesity prevalence and trends in Latin American countries. *Obesity Reviews*, 2001;2:99-196
- 1998: Instituto Nacional de Estadística - INE/Guatemala and Macro International. 1999. Guatemala Encuesta Nacional de Salud Materno Infantil 1998-1999. Calverton, Maryland, USA: Instituto Nacional de Estadística - INE/Guatemala and Macro International.
- 1999: Centres for Disease Control and Prevention. <http://www.cdc.gov/>
- 2000: Instituto Nacional de Estadística e Informática - INEI/Perú and Macro International. 2001. Perú Encuesta Demográfica y de Salud Familiar 2000. Lima, Perú: INEI/Perú and Macro International.
- 2001: Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of Overweight and Obesity in the United States, 1999-2004. *JAMA* 2006;295(13):1549-1555
- 2002: Monteiro CA, Conde WL and Popkin BA. (2007). Income-specific trends in obesity in Brazil: 1975 - 2003. *American Journal of Public Health*, 97 (10): 1808 - 1812.
- 2003: 2003 ENS Report. Final results on the National Health Survey. <http://epi.minsal.cl/epi/html/invest/ENS/informeFinalENS.pdf>.
- 2004: Instituto Colombiano de Bienestar Familiar. Encuesta Nacional de la Situación Nutricional (ENSIN), 2005. (2006). Available at: [https://www.icbf.gov.co/sites/default/files/libro\\_2005.pdf](https://www.icbf.gov.co/sites/default/files/libro_2005.pdf) [Accessed 18.07.25]
- 2005: Cayemittes, Michel, Marie Florence Placide, Soumaïla Mariko, Bernard Barrère, Blaise Sévère, and Canez Alexandre. 2007. Enquête Mortalité, Morbidité et Utilisation des Services, Haïti, 2005-2006. Calverton, Maryland, USA: Ministère de la Santé Publique et de la Population, Institut Haïtien de l'Enfance and Macro International.
- 2006: Olaiz-Fernández G, Rivera-Dommarco J, Shamah-Levy T, Rojas R, Villalpando-Hernández S, Hernández-Avila M, Sepúlveda-Amor J. Encuesta Nacional de Salud y Nutrición 2006. Cuernavaca, México: Instituto Nacional de Salud Pública, 2006. (National Health and Nutrition Survey 2006).
- 2007: Instituto Nacional de Estadística e Informática - INEI/Peru and ORC Macro. 2009. Perú Encuesta Demográfica y de Salud Familiar - ENDES 2007-2008. Lima, Perú: INEI/Peru and ORC Macro.
- 2008: Ramirez-Zea M, Kroker-Lobos MF, Close-Fernandez R, Kanter R. The double burden of malnutrition in indigenous and nonindigenous Guatemalan populations. *Am J Clin Nutr*. 2014 Dec;100(6):1644S-51S. doi: 10.3945/ajcn.114.083857
- 2009: 2nd Argentinian National Survey of Risk Factors (Encuesta Nacional de Factores de Riesgo). Results from 1-3rd survey reported in the 4th survey report: [https://www.indec.gob.ar/ftp/cuadros/publicaciones/enfr\\_2018\\_resultados\\_definitivos.pdf](https://www.indec.gob.ar/ftp/cuadros/publicaciones/enfr_2018_resultados_definitivos.pdf)

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

---

## Women



### References:


- 1960, 1971, 1973, 1976, 1991: Flegal KM, Carroll MD, Kuczmarski RJ, Johnson CL. Overweight and obesity in the United States: prevalence and trends, 1960-1994. *International Journal of Obesity* (1998);22:39-47
- 1975: Monteiro CA, Conde WL, Popking BM. Is obesity replacing or adding to undernutrition? Evidence from different social classes in Brazil. 2002. *Public Health Nutrition*:51(1A), 105-112
- 1988: Berrios X, Jadue I, Zenteno J, Ross MI, Rodriguez H. Prevalencia de factores de riesgo para enfermedades cronicas. Estudio de la poblacion general de la region Metropolitana, 1986-1987. *Rev. Med. Chile*. 1990;118:597-604
- 1992, 1994, 1995: Martorell R, Khan LK, Hughes ML, Grummer Strawn LM. Obesity in women from developing countries. *EJCN* (2000) 54:247-252
- 1997: Filozof C, Gonzales C, Sereday M, Mazza C, Braguinsky J. Obesity prevalence and trends in Latin American countries. *Obesity Reviews*, 2001;2:99-196
- 1998: Instituto Nacional de Estadística - INE/Guatemala and Macro International. 1999. Guatemala Encuesta Nacional de Salud Materno Infantil 1998-1999. Calverton, Maryland, USA: Instituto Nacional de Estadística - INE/Guatemala and Macro International.
- 1999: Centres for Disease Control and Prevention. <http://www.cdc.gov/>
- 2000: Instituto Nacional de Estadística e Informática - INEI/Perú and Macro International. 2001. Perú Encuesta Demográfica y de Salud Familiar 2000. Lima, Perú: INEI/Perú and Macro International.
- 2001: Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of Overweight and Obesity in the United States, 1999-2004. *JAMA* 2006;295(13):1549-1555
- 2002: Monteiro CA, Conde WL and Popkin BA. (2007). Income-specific trends in obesity in Brazil: 1975 - 2003. *American Journal of Public Health*, 97 (10): 1808 - 1812.
- 2003: 2003 ENS Report. Final results on the National Health Survey. <http://epi.minsal.cl/epi/html/invest/ENS/informeFinalENS.pdf>.
- 2004: Instituto Colombiano de Bienestar Familiar. Encuesta Nacional de la Situación Nutricional (ENSIN), 2005. (2006). Available at: [https://www.icbf.gov.co/sites/default/files/libro\\_2005.pdf](https://www.icbf.gov.co/sites/default/files/libro_2005.pdf) [Accessed 18.07.25]
- 2005: Cayemittes, Michel, Marie Florence Placide, Soumaïla Mariko, Bernard Barrère, Blaise Sévère, and Canez Alexandre. 2007. Enquête Mortalité, Morbidité et Utilisation des Services, Haïti, 2005-2006. Calverton, Maryland, USA: Ministère de la Santé Publique et de la Population, Institut Haïtien de l'Enfance and Macro International.
- 2006: Olaiz-Fernández G, Rivera-Dommarco J, Shamah-Levy T, Rojas R, Villalpando-Hernández S, Hernández-Avila M, Sepúlveda-Amor J. Encuesta Nacional de Salud y Nutrición 2006. Cuernavaca, México: Instituto Nacional de Salud Pública, 2006. (National Health and Nutrition Survey 2006).
- 2007: Instituto Nacional de Estadística e Informática - INEI/Peru and ORC Macro. 2009. Perú Encuesta Demográfica y de Salud Familiar - ENDES 2007-2008. Lima, Perú: INEI/Peru and ORC Macro.
- 2008: Ramirez-Zea M, Kroker-Lobos MF, Close-Fernandez R, Kanter R. The double burden of malnutrition in indigenous and nonindigenous Guatemalan populations. *Am J Clin Nutr*. 2014 Dec;100(6):1644S-51S. doi: 10.3945/ajcn.114.083857
- 2009: 2nd Argentinian National Survey of Risk Factors (Encuesta Nacional de Factores de Riesgo). Results from 1-3rd survey reported in the 4th survey report: [https://www.indec.gob.ar/ftp/cuadros/publicaciones/enfr\\_2018\\_resultados\\_definitivos.pdf](https://www.indec.gob.ar/ftp/cuadros/publicaciones/enfr_2018_resultados_definitivos.pdf)

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

---

## Overweight/obesity by education

### Women, 2024



Survey type: Measured

Age: 15-49

Sample size: 29467


Area covered: National

References: National Institute of Statistics and Informatics. Peru: Demographic and Family Health Survey, Endes 2024. <https://www.gob.pe/en/institucion/inei/informes-publicaciones/6813623-peru-demographic-and-family-health-survey-endes-2024> [Accessed 25.09.25]

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.*

## Overweight/obesity by age

### Women, 2024




Survey type:	Measured
Sample size:	29,467
Area covered:	National
References:	National Institute of Statistics and Informatics. Peru: Demographic and Family Health Survey, Endes 2024. <a href="https://www.gob.pe/en/institucion/inei/informes-publicaciones/6813623-peru-demographic-and-family-health-survey-endes-2024">https://www.gob.pe/en/institucion/inei/informes-publicaciones/6813623-peru-demographic-and-family-health-survey-endes-2024</a> [Accessed 25.09.25]

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.

## Overweight/obesity by region

### Women, 2024




Survey type:	Measured
Age:	15-49
Sample size:	29,467
Area covered:	National

References: National Institute of Statistics and Informatics. Peru: Demographic and Family Health Survey, Endes 2024. <https://www.gob.pe/en/institucion/inei/informes-publicaciones/6813623-peru-demographic-and-family-health-survey-endes-2024> [Accessed 25.09.25]

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.*

## Overweight/obesity by socio-economic group

### Women, 2024



Survey type: Measured

Age: 15-49

Sample size: 29,467

Area covered: National

References: National Institute of Statistics and Informatics. Peru: Demographic and Family Health Survey, Endes 2024. <https://www.gob.pe/en/institucion/inei/informes-publicaciones/6813623-peru-demographic-and-family-health-survey-endes-2024> [Accessed 25.09.25]


Definitions: Wealth quintile

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.

## Overweight/obesity by ethnicity

Ethnic groups are as defined by publication of origin and are not as defined by WOF. In some instances ethnicity is conflated with nationality and/or race.

### Adults, 2024



Survey type: Measured

Age: 15+

Sample size: 30,631

Area covered: National

References: Instituto Nacional de Estadística e Informática. Perú: Enfermedades No Transmisibles y Transmisibles 2024. [https://proyectos.inei.gob.pe/endes/2024/SALUD/ENFERMEDADES\\_ENDES\\_2024.pdf](https://proyectos.inei.gob.pe/endes/2024/SALUD/ENFERMEDADES_ENDES_2024.pdf) [Accessed 01.10.25]


Notes: ENDES 2024. Standard overweight and obesity cutoffs used except in adolescents aged 15-17 where they are measured using WHO cutoffs, and for adults aged 60 and over, overweight refers to BMI between 28kg/m<sup>2</sup> and 31.9kg/m<sup>2</sup> and obesity refers to ≥32kg/m<sup>2</sup>. Interpret with caution when comparing cross-nationally.

Definitions: Ethnic self-identification. Native origin: Quechua, Aymara, native of the Amazon, belonging to or part of another indigenous or native people. Negro, moreno, zambo also includes Afro-Peruvian and Afro-descendant. Other includes those who did not know.

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.

## Double burden of underweight & overweight


### Adults, 2022



<b>Survey type:</b>	Measured
<b>Age:</b>	20+
<b>References:</b>	NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population representative studies with 222 million children, adolescents, and adults. Lancet 2024; published online Feb 29. <a href="https://doi.org/10.1016/S0140-6736(23)02750-2">https://doi.org/10.1016/S0140-6736(23)02750-2</a> .
<b>Notes:</b>	Age Standardised estimates
<b>Definitions:</b>	Combined prevalence of BMI<18.5 kg/m <sup>2</sup> and BMI>=30 kg/m <sup>2</sup> (double burden of underweight and obesity)


## Insufficient physical activity

### Adults, 2022




<b>Survey type:</b>	Self-reported
<b>Age:</b>	18+
<b>Area covered:</b>	National
<b>References:</b>	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at <a href="https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)">https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)</a>
<b>Definitions:</b>	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.

## Men, 2022



<b>Survey type:</b>	Self-reported
<b>Age:</b>	18+
<b>Area covered:</b>	National
<b>References:</b>	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at <a href="https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)">https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)</a>
<b>Definitions:</b>	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.

## Women, 2022



Survey type: Self-reported

Age: 18+


Area covered: National

References: WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-\(age-standardized-estimate\)-\(-\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-))

Definitions: Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.

## Mental health - depression disorders

### Adults, 2023




Age: 20+

Area covered: National

References: Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from <https://vizhub.healthdata.org/gbd-compare/> [Accessed 17.03.26]


Definitions: Estimated number living with depressive disorders per 100,000 population (20+ years of age)

**Men, 2023**



<b>Age:</b>	20+
<b>Area covered:</b>	National
<b>References:</b>	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from <a href="https://vizhub.healthdata.org/gbd-compare/">https://vizhub.healthdata.org/gbd-compare/</a> [Accessed 17.03.26]
<b>Definitions:</b>	Estimated number living with depressive disorders per 100,000 population (20+ years of age)


### Women, 2023



<b>Age:</b>	20+
<b>Area covered:</b>	National
<b>References:</b>	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from <a href="https://vizhub.healthdata.org/gbd-compare/">https://vizhub.healthdata.org/gbd-compare/</a> [Accessed 17.03.26]
<b>Definitions:</b>	Estimated number living with depressive disorders per 100,000 population (20+ years of age)

## Mental health - anxiety disorders

### Adults, 2023




Age: 20+

Area covered: National

References: Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from <https://vizhub.healthdata.org/gbd-compare/> [Accessed 17.03.26]


Definitions: Estimated number living with anxiety disorders per 100,000 population (20+ years of age)

### Men, 2023



<b>Age:</b>	20+
<b>Area covered:</b>	National
<b>References:</b>	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from <a href="https://vizhub.healthdata.org/gbd-compare/">https://vizhub.healthdata.org/gbd-compare/</a> [Accessed 17.03.26]
<b>Definitions:</b>	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)


**Women, 2023**



<b>Age:</b>	20+
<b>Area covered:</b>	National
<b>References:</b>	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from <a href="https://vizhub.healthdata.org/gbd-compare/">https://vizhub.healthdata.org/gbd-compare/</a> [Accessed 17.03.26]
<b>Definitions:</b>	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)

## Percent of population who cannot afford a healthy diet

### Adults, 2024



Area covered:

National

References:


FAO. 2025. FAOSTAT: Cost and Affordability of a Healthy Diet (CoAHD). <https://www.fao.org/faostat/en/#data/CAHD>. Licence: CC-BY-4.0. [Last Accessed 15.10.25]

Notes:

The percent of the population whose food budget is below the cost of a healthy diet.

## Oesophageal cancer

### Men, 2022




Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000


## Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <a href="https://gco.iarc.who.int/today">https://gco.iarc.who.int/today</a> , accessed [16.07.24]
Definitions:	Indicence per 100,000

## Breast cancer

### Women, 2022



Area covered:

National

References:


Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions:

Incidence per 100,000

## Colorectal cancer

### Men, 2022




Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000


**Women, 2022**



<b>Age:</b>	20+
<b>Area covered:</b>	National
<b>References:</b>	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <a href="https://gco.iarc.who.int/today">https://gco.iarc.who.int/today</a> , accessed [16.07.24]
<b>Definitions:</b>	Incidence per 100,000

## Pancreatic cancer

### Men, 2022




Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000


## Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <a href="https://gco.iarc.who.int/today">https://gco.iarc.who.int/today</a> , accessed [16.07.24]
Definitions:	Incidence per 100,000

## Gallbladder cancer

### Men, 2022




Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000


## Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <a href="https://gco.iarc.who.int/today">https://gco.iarc.who.int/today</a> , accessed [16.07.24]
Definitions:	Indicence per 100,000

## Kidney cancer

### Men, 2022




Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000


### Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <a href="https://gco.iarc.who.int/today">https://gco.iarc.who.int/today</a> , accessed [16.07.24]
Definitions:	Incidence per 100,000

## Cancer of the uterus

### Women, 2022




Age: 20+

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

## Raised blood pressure


### Adults, 2015



**References:** Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A875?lang=en>

**Definitions:** Age Standardised estimated % Raised blood pressure 2015 (SBP $\geq$ 140 OR DBP $\geq$ 90).


## Men, 2015



**References:** Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A875?lang=en>

**Definitions:** Age Standardised estimated % Raised blood pressure 2015 (SBP $\geq$ 140 OR DBP $\geq$ 90).

## Women, 2015




**References:** Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A875?lang=en>

**Definitions:** Age Standardised estimated % Raised blood pressure 2015 (SBP $\geq$ 140 OR DBP $\geq$ 90).

## Raised cholesterol


### Adults, 2008



**References:** Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A885>

**Definitions:** % Raised total cholesterol ( $\geq 5.0$  mmol/L) (age-standardized estimate).


## Men, 2008



**References:** Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A885>

**Definitions:** % Raised total cholesterol ( $\geq 5.0$  mmol/L) (age-standardized estimate).

## Women, 2008




**References:** Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A885>

**Definitions:** % Raised total cholesterol ( $\geq 5.0$  mmol/L) (age-standardized estimate).

## Raised fasting blood glucose


### Men, 2014



**References:** Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A869?lang=en>

**Definitions:** Age Standardised % raised fasting blood glucose ( $\geq 7.0$  mmol/L or on medication).

## Women, 2014




**References:** Global Health Observatory data repository, World Health Organisation, <http://apps.who.int/gho/data/node.main.A869?lang=en>

**Definitions:** Age Standardised % raised fasting blood glucose ( $\geq 7.0$  mmol/L or on medication).

## Diabetes prevalence

### Adults, 2024




**References:**

Reproduced with kind permission International Diabetes Federation. IDF Diabetes Atlas, 11th edn. Brussels, Belgium:International Diabetes Federation, 2025. <http://www.diabetesatlas.org>

## Ovarian Cancer

### Women, 2022




Age: 20+

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

## Leukemia

### Men, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.

## Women, 2022



Age: 20+

Area covered: National


References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.

## Liver and intrahepatic bile duct Cancer

### Men, 2022




Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000


## Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <a href="https://gco.iarc.who.int/today">https://gco.iarc.who.int/today</a> , accessed [16.07.24]
Definitions:	Incidence per 100,000

## Multiple Myeloma

### Men, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.

## Women, 2022



Age: 20+

Area covered: National


References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.

## Non Hodgkin Lymphoma

### Men, 2022



Age: 20+


Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.*

## Women, 2022



Age: 20+

Area covered: National


References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m<sup>2</sup>, obesity refers to a BMI greater than 30kg/m<sup>2</sup>.

## Thyroid Cancer

### Men, 2022




Age: 20+

Area covered: National

References: Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.who.int/today>, accessed [16.07.24]

Definitions: Incidence per 100,000

**Women, 2022**



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024). Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <a href="https://gco.iarc.who.int/today">https://gco.iarc.who.int/today</a> , accessed [16.07.24]
Definitions:	Incidence per 100,000

PDF created on May 5, 2026