
Russian Federation



Country report card


This report card contains the latest data available on the Global Obesity Observatory on overweight and obesity . Where available, data on common and relevant obesity drivers and comorbidities are also presented.

View the latest version of this report on the Global Obesity Observatory at https://data.worldobesity.org/country/russian-federation-179/.

Contents	Page
Obesity prevalence	3
Trend: % Adults living with obesity in Russia 1992-2012	10
Trend: % Adults living with overweight or obesity in Russia 1992-2012	15
Overweight/obesity by education	20
Overweight/obesity by age	27
Overweight/obesity by region	32
Overweight/obesity by socio-economic group	43
Double burden of underweight & obesity	52
Insufficient physical activity	57
Prevalence of at least daily soft drink consumption	70
Prevalence of less than daily fruit consumption	73
Prevalence of less than daily vegetable consumption	76
Average weekly frequency of fast food consumption	79
Mental health - depression disorders	86
Mental health - anxiety disorders	93
Percent of population who cannot afford a healthy diet	100
Oesophageal cancer	101
Breast cancer	104
Colorectal cancer	106
Pancreatic cancer	109
Gallbladder cancer	112
Kidney cancer	115
Cancer of the uterus	118
Raised blood pressure	120
Raised cholesterol	124
Raised fasting blood glucose	127
Diabetes prevalence	129
Ovarian Cancer	130
Leukemia	132
Liver and intrahepatic bile duct Cancer	137
Multiple Myeloma	140
Non Hodgkin Lymphoma	145
Thyroid Cancer	150

Obesity prevalence


Adults, 2020-2022



Survey type:	Measured
Age:	35-74
Sample size:	28,710
Area covered:	National (15 regions)
References:	Balanova Yu.A., Drapkina, O.M. et al. Obesity in the Russian population during the COVID-19 pandemic and associated factors. Data from the ESSE-RF3 study. Cardiovascular Therapy and Prevention. 2023;22(8S):3793. (In Russ.) https://doi.org/10.15829/1728-8800-2023-3793 [Accessed 06.08.25]

Notes:	Survey conducted in 15 Russian regions covering all federal districts
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Children, 2018-2020



Survey type:	Measured
Age:	7
Sample size:	2081
Area covered:	Regional (Moscow)
References:	Report on the fifth round of data collection, 2018–2020: WHO European Childhood Obesity Surveillance Initiative (COSI). Copenhagen: WHO Regional Office for Europe; 2022. Licence: CC BY-NC-SA 3.0 IGO.

Cutoffs:	WHO 2007

0-5 years, 2005




Age:	0-5
Sample size:	343
References:	Other: Russia Longitudinal Monitoring Survey (RLMS-HSE) Round XIV 2005
Notes:	UNICEF/WHO/World Bank Joint Child Malnutrition Estimates Expanded Database: Overweight (Survey Estimates), May 2023, New York. For more information about the methodology, please consult https://data.unicef.org/resources/jme-2023-country-consultations/ Percentage of children under 5 years of age falling above 2 standard deviations (moderate and severe) from the median weight-for-height of the reference population.

Definitions:	=>+2SD

% Adults living with obesity in Russia 1992-2012

Men




Survey type:	Measured
References:	1992: Jahns L, Baturin A, Popkin BM. Obesity, diet, and poverty: trends in the Russian transition to market economy. EJCN 2003;57:1295-1302 2000: Jahns L, Baturin A, Popkin BM. Obesity, diet, and poverty: trends in the Russian transition to market economy. EJCN 2003;57:1295-1302. 2012: Kontsevaya A, Shalnova S, Deev A et al. Overweight and Obesity in the Russian Population: Prevalence in Adults and Association with Socioeconomic Parameters. and Cardiovascular Risk Factors.Obesity Facts 2019;12:103–114

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women




Survey type:	Measured
References:	1992: Jahns L, Baturin A, Popkin BM. Obesity, diet, and poverty: trends in the Russian transition to market economy. EJCN 2003;57:1295-1302 2000: Jahns L, Baturin A, Popkin BM. Obesity, diet, and poverty: trends in the Russian transition to market economy. EJCN 2003;57:1295-1302. 2012: Kontsevaya A, Shalnova S, Deev A et al. Overweight and Obesity in the Russian Population: Prevalence in Adults and Association with Socioeconomic Parameters. and Cardiovascular Risk Factors.Obesity Facts 2019;12:103–114

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

% Adults living with overweight or obesity in Russia 1992-2012

Men




Survey type:	Measured
References:	1992: Jahns L, Baturin A, Popkin BM. Obesity, diet, and poverty: trends in the Russian transition to market economy. EJCN 2003;57:1295-1302 2000: Jahns L, Baturin A, Popkin BM. Obesity, diet, and poverty: trends in the Russian transition to market economy. EJCN 2003;57:1295-1302. 2012: Kontsevaya A, Shalnova S, Deev A et al. Overweight and Obesity in the Russian Population: Prevalence in Adults and Association with Socioeconomic Parameters. and Cardiovascular Risk Factors.Obesity Facts 2019;12:103–114

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Women




Survey type:	Measured
References:	1992: Jahns L, Baturin A, Popkin BM. Obesity, diet, and poverty: trends in the Russian transition to market economy. EJCN 2003;57:1295-1302 2000: Jahns L, Baturin A, Popkin BM. Obesity, diet, and poverty: trends in the Russian transition to market economy. EJCN 2003;57:1295-1302. 2012: Kontsevaya A, Shalnova S, Deev A et al. Overweight and Obesity in the Russian Population: Prevalence in Adults and Association with Socioeconomic Parameters. and Cardiovascular Risk Factors.Obesity Facts 2019;12:103–114

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.

Overweight/obesity by education


Adults, 2020-2022



Survey type:	Measured
Age:	35-74
Sample size:	28,710
Area covered:	National (15 regions)

References:	Balanova Yu.A., Drapkina, O.M. et al. Obesity in the Russian population during the COVID-19 pandemic and associated factors. Data from the ESSE-RF3 study. Cardiovascular Therapy and Prevention. 2023;22(8S):3793. (In Russ.) https://doi.org/10.15829/1728-8800-2023-3793 [Accessed 06.08.25]
Notes:	Survey conducted in 15 Russian regions covering all federal districts
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Men, 2020-2022



Survey type:	Measured
Age:	35-74
Sample size:	28,710
Area covered:	National (15 regions)

References:	Balanova Yu.A., Drapkina, O.M. et al. Obesity in the Russian population during the COVID-19 pandemic and associated factors. Data from the ESSE-RF3 study. Cardiovascular Therapy and Prevention. 2023;22(8S):3793. (In Russ.) https://doi.org/10.15829/1728-8800-2023-3793 [Accessed 06.08.25]
Notes:	Survey conducted in 15 Russian regions covering all federal districts
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2020-2022




Survey type:	Measured
Age:	35-74
Sample size:	28,710
Area covered:	National (15 regions)

References:	Balanova Yu.A., Drapkina, O.M. et al. Obesity in the Russian population during the COVID-19 pandemic and associated factors. Data from the ESSE-RF3 study. Cardiovascular Therapy and Prevention. 2023;22(8S):3793. (In Russ.) https://doi.org/10.15829/1728-8800-2023-3793 [Accessed 06.08.25]
Notes:	Survey conducted in 15 Russian regions covering all federal districts
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Overweight/obesity by age


Adults, 2020-2022



Survey type:	Measured
Sample size:	28,710
Area covered:	National (15 regions)
References:	Balanova Yu.A., Drapkina, O.M. et al. Obesity in the Russian population during the COVID-19 pandemic and associated factors. Data from the ESSE-RF3 study. Cardiovascular Therapy and Prevention. 2023;22(8S):3793. (In Russ.) https://doi.org/10.15829/1728-8800-2023-3793 [Accessed 06.08.25]

Notes:	Survey conducted in 15 Russian regions covering all federal districts
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Children, 2005




Survey type:	Measured
Sample size:	1081
Area covered:	National
References:	World Obesity Reanalysis of RLMS 2005 Original Source: Ã?Â¬Ã?Â¬Russia Longitudinal Monitoring survey, RLMS-HSEÃ?Â¬Ã?Â®, conducted by HSE and ZAO Ã?Â¬Ã?Â¬DemoscopeÃ?Â¬Ã?Â® together with Carolina Population Center, University of North Carolina at Chapel Hill and the Institute of Sociology RAS. (RLMS-HSE sites: http://www.cpc.unc.edu/projects/rlms-hse, http://www.hse.ru/org/hse/rlms)

Cutoffs:	IOTF

Overweight/obesity by region


Adults, 2020-2022



Survey type:	Measured
Age:	35-74
Sample size:	28,710
Area covered:	National (15 regions)
References:	Balanova Yu.A., Drapkina, O.M. et al. Obesity in the Russian population during the COVID-19 pandemic and associated factors. Data from the ESSE-RF3 study. Cardiovascular Therapy and Prevention. 2023;22(8S):3793. (In Russ.) https://doi.org/10.15829/1728-8800-2023-3793 [Accessed 06.08.25]

Notes:	Survey conducted in 15 Russian regions covering all federal districts
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Men, 2020-2022



Survey type:	Measured
Age:	35-74
Sample size:	28,710
Area covered:	National (15 regions)
References:	Balanova Yu.A., Drapkina, O.M. et al. Obesity in the Russian population during the COVID-19 pandemic and associated factors. Data from the ESSE-RF3 study. Cardiovascular Therapy and Prevention. 2023;22(8S):3793. (In Russ.) https://doi.org/10.15829/1728-8800-2023-3793 [Accessed 06.08.25]

Notes:	Survey conducted in 15 Russian regions covering all federal districts
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Women, 2020-2022



Survey type:	Measured
Age:	35-74
Sample size:	28,710
Area covered:	National (15 regions)
References:	Balanova Yu.A., Drapkina, O.M. et al. Obesity in the Russian population during the COVID-19 pandemic and associated factors. Data from the ESSE-RF3 study. Cardiovascular Therapy and Prevention. 2023;22(8S):3793. (In Russ.) https://doi.org/10.15829/1728-8800-2023-3793 [Accessed 06.08.25]

Notes:	Survey conducted in 15 Russian regions covering all federal districts
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Children, 1992



Survey type:	Measured
Age:	6-9
Sample size:	Total sample size (6-18yrs old) = 6883
Area covered:	National
References:	Youfa Wang.Cross-national comparison of childhood obesity: the epidemic and the relationship between obesity and socioeconomic status. Int. J. Epidemiol. (2001) 30 (5): 1129-1136. doi: 10.1093/ije/30.5.1129

Notes:	The study used the US NCHS body mass index (BMI = wt/ht2) reference to define obesity (BMI ≥95th percentile) and overweight (85th≤BMI<95th percentile).
Cutoffs:	Other

0-5 years, 2005




Sample size:	343
References:	Other: Russia Longitudinal Monitoring Survey (RLMS-HSE) Round XIV 2005
Notes:	UNICEF/WHO/World Bank Joint Child Malnutrition Estimates Expanded Database: Overweight (Survey Estimates), May 2023, New York. For more information about the methodology, please consult https://data.unicef.org/resources/jme-2023-country-consultations/ Percentage of children under 5 years of age falling above 2 standard deviations (moderate and severe) from the median weight-for-height of the reference population.

Definitions:	=>+2SD

Overweight/obesity by socio-economic group


Adults, 2020-2022



Survey type:	Measured
Age:	35-74
Sample size:	28.708
Area covered:	National (15 regions)
References:	Balanova Yu.A., Drapkina, O.M. et al. Obesity in the Russian population during the COVID-19 pandemic and associated factors. Data from the ESSE-RF3 study. Cardiovascular Therapy and Prevention. 2023;22(8S):3793. (In Russ.) https://doi.org/10.15829/1728-8800-2023-3793 [Accessed 06.08.25]

Notes:	Survey conducted in 15 Russian regions covering all federal districts
Definitions:	Income level determined through survey questions on share of income spent on food, opinion on family's financial capabilities, and security in comparison to other families
	Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Men, 2020-2022




Survey type:	Measured
Age:	35-74
Sample size:	28.708
Area covered:	National (15 regions)

References:	Balanova Yu.A., Drapkina, O.M. et al. Obesity in the Russian population during the COVID-19 pandemic and associated factors. Data from the ESSE-RF3 study. Cardiovascular Therapy and Prevention. 2023;22(8S):3793. (In Russ.) https://doi.org/10.15829/1728-8800-2023-3793 [Accessed 06.08.25]
Notes:	Survey conducted in 15 Russian regions covering all federal districts
Definitions:	Income level determined through survey questions on share of income spent on food, opinion on family's financial capabilities, and security in comparison to other families

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2020-2022




Survey type:	Measured
Age:	35-74
Sample size:	28.708
Area covered:	National (15 regions)

References:	Balanova Yu.A., Drapkina, O.M. et al. Obesity in the Russian population during the COVID-19 pandemic and associated factors. Data from the ESSE-RF3 study. Cardiovascular Therapy and Prevention. 2023;22(8S):3793. (In Russ.) https://doi.org/10.15829/1728-8800-2023-3793 [Accessed 06.08.25]
Notes:	Survey conducted in 15 Russian regions covering all federal districts
Definitions:	Income level determined through survey questions on share of income spent on food, opinion on family's financial capabilities, and security in comparison to other families

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Children, 1992




Survey type:	Measured
Age:	6-9
Sample size:	Total sample size (6-18yrs old) = 6883
Area covered:	National
References:	Youfa Wang.Cross-national comparison of childhood obesity: the epidemic and the relationship between obesity and socioeconomic status. Int. J. Epidemiol. (2001) 30 (5): 1129-1136. doi: 10.1093/ije/30.5.1129

Notes:	The study used the US NCHS body mass index (BMI = wt/ht2) reference to define obesity (BMI ≥95th percentile) and overweight (85th≤BMI<95th percentile).
Cutoffs:	Other

Double burden of underweight & obesity


Adults, 2022



Survey type:	Measured
Age:	20+
References:	NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population representative studies with 222 million children, adolescents, and adults. Lancet 2024; published online Feb 29. https://doi.org/10.1016/S0140-6736(23)02750-2.
Notes:	Age Standardised estimates

Definitions:	Combined prevalence of BMI<18.5 kg/m² and BMI>=30 kg/m² (double burden of underweight and obesity)

Children, 2022




Survey type:	Measured
Age:	5-19
References:	NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population representative studies with 222 million children, adolescents, and adults. Lancet 2024; published online Feb 29. https://doi.org/10.1016/S0140-6736(23)02750-2.
Notes:	Age standardised estimates

Definitions:	Combined prevalence of BMI < -2SD and BMI > 2SD (double burden of thinness and obesity)
Cutoffs:	BMI < -2SD and BMI > 2SD

Insufficient physical activity


Adults, 2022



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.


Men, 2022



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.


Women, 2022



Survey type:	Self-reported
Age:	18+
Area covered:	National
References:	WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-)

Definitions:	Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent.


Children, 2025



Survey type:	Self-reported
Age:	11-17
Area covered:	National
References:	World Health Organization. (2026). Physical activity among adolescents, no active days in the past week, prevalence (%). Global Health Observatory. Available at: https://www.who.int/data/gho/data/indicators/indicator-details/GHO/gho-physical-activity-among-adolescents-no-active-days-in-the-past-week-prevalence [Accessed 06.11.26]

Notes:	An active day is a day on which an adolescent reports at least 60 minutes of moderate-to-vigorous physical activity (MVPA).
Definitions:	Percentage of adolescents aged 11–17 years who reported no active day in the past 7 days (0 active days per week).


Boys, 2025



Survey type:	Self-reported
Age:	11-17
Area covered:	National
References:	World Health Organization. (2026). Physical activity among adolescents, no active days in the past week, prevalence (%). Global Health Observatory. Available at: https://www.who.int/data/gho/data/indicators/indicator-details/GHO/gho-physical-activity-among-adolescents-no-active-days-in-the-past-week-prevalence [Accessed 06.11.26]

Notes:	An active day is a day on which an adolescent reports at least 60 minutes of moderate-to-vigorous physical activity (MVPA).
Definitions:	Percentage of adolescents aged 11–17 years who reported no active day in the past 7 days (0 active days per week).

Girls, 2025




Survey type:	Self-reported
Age:	11-17
Area covered:	National
References:	World Health Organization. (2026). Physical activity among adolescents, no active days in the past week, prevalence (%). Global Health Observatory. Available at: https://www.who.int/data/gho/data/indicators/indicator-details/GHO/gho-physical-activity-among-adolescents-no-active-days-in-the-past-week-prevalence [Accessed 06.11.26]

Notes:	An active day is a day on which an adolescent reports at least 60 minutes of moderate-to-vigorous physical activity (MVPA).
Definitions:	Percentage of adolescents aged 11–17 years who reported no active day in the past 7 days (0 active days per week).

Prevalence of at least daily soft drink consumption

Children, 2014




Survey type:	Measured
References:	World Health Organization. (2017). Adolescent obesity and related behaviours: Trends and inequalities in the who european region, 2002-2014: observations from the Health Behavior in School-aged Children (HBSC) WHO collaborative cross-national study (J. Inchley, D. Currie, J. Jewel, J. Breda, & V. Barnekow, Eds.). World Health Organization. Sourced from Food Systems Dashboard http://www.foodsystemsdashboard.org
Notes:	15-year-old adolescents

Definitions:	Prevalence of at least daily carbonated soft drink consumption (% of at least daily carbonated soft drink consumption)

Prevalence of less than daily fruit consumption

Children, 2014




Survey type:	Self-reported
Age:	11-15
Area covered:	National
References:	World Health Organization. (2017). Adolescent obesity and related behaviours: Trends and inequalities in the who european region, 2002-2014: observations from the Health Behavior in School-aged Children (HBSC) WHO collaborative cross-national study (J. Inchley, D. Currie, J. Jewel, J. Breda, & V. Barnekow, Eds.).

Definitions:	Proportion who reported eating fruit less than daily

Prevalence of less than daily vegetable consumption

Children, 2014




Survey type:	Self-reported
Age:	11-15
Area covered:	National
References:	World Health Organization. (2017). Adolescent obesity and related behaviours: Trends and inequalities in the who european region, 2002-2014: observations from the Health Behavior in School-aged Children (HBSC) WHO collaborative cross-national study (J. Inchley, D. Currie, J. Jewel, J. Breda, & V. Barnekow, Eds.).

Definitions:	Prevalence of less-than-daily vegetable consumption (% less-than-daily vegetable consumption)

Average weekly frequency of fast food consumption


Adults, 2021-2024



Survey type:	Self-reported
Age:	15+
Sample size:	2000
Area covered:	National
References:	Global Diet Quality Project. 2024. "DQQ Results Dataset 2021-2024." Harvard Dataverse. https://doi.org/doi:10.7910/DVN/KY3W8A Licence: CC BY-NC-SA 4.0. Accessed at dietquality.org on October 15, 2025.

Notes:	For more global dietary survey data, visit the Global Diet Quality Project website at https://www.dietquality.org/
Definitions:	Proportion of the population who consumed fast food during the previous day or night (self-reported)


Men, 2021-2024



Survey type:	Self-reported
Age:	15+
Sample size:	2000
Area covered:	National
References:	Global Diet Quality Project. 2024. "DQQ Results Dataset 2021-2024." Harvard Dataverse. https://doi.org/doi:10.7910/DVN/KY3W8A Licence: CC BY-NC-SA 4.0. Accessed at dietquality.org on October 15, 2025.

Notes:	For more global dietary survey data, visit the Global Diet Quality Project website at https://www.dietquality.org/
Definitions:	Proportion of the population who consumed fast food during the previous day or night (self-reported)

Women, 2021-2024




Survey type:	Self-reported
Age:	15+
Sample size:	2000
Area covered:	National
References:	Global Diet Quality Project. 2024. "DQQ Results Dataset 2021-2024." Harvard Dataverse. https://doi.org/doi:10.7910/DVN/KY3W8A Licence: CC BY-NC-SA 4.0. Accessed at dietquality.org on October 15, 2025.

Notes:	For more global dietary survey data, visit the Global Diet Quality Project website at https://www.dietquality.org/
Definitions:	Proportion of the population who consumed fast food during the previous day or night (self-reported)


Mental health - depression disorders

Adults, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)

Men, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)

Women, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (20+ years of age)

Children, 2023




Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (under 20 years of age)

Boys, 2023



Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (under 20 years of age)


Girls, 2023



Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with depressive disorders per 100,000 population (under 20 years of age)


Mental health - anxiety disorders

Adults, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)

Men, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)

Women, 2023




Age:	20+
Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (20+ years of age)

Children, 2023




Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (under 20 years of age)

Boys, 2023



Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (under 20 years of age)


Girls, 2023



Area covered:	National
References:	Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26]
Definitions:	Estimated number living with anxiety disorders per 100,000 population (under 20 years of age)

Percent of population who cannot afford a healthy diet


Adults, 2025



Area covered:	National
References:	FAO. 2026. FAOSTAT: Cost and Affordability of a Healthy Diet (CoAHD). https://www.fao.org/faostat/en/#data/CAHD. Licence: CC-BY-4.0. [Last Accessed 12.10.26]
Notes:	The percent of the population whose food budget is below the cost of a healthy diet.


Oesophageal cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Breast cancer


Women, 2022



Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Colorectal cancer

Men, 2022



Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Pancreatic cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Gallbladder cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000


Kidney cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Cancer of the uterus


Women, 2022



Age:	20+
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Raised blood pressure

Adults, 2019




Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.

Men, 2019



Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.


Women, 2019



Age:	30-79
References:	World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26]
Definitions:	Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79.


Raised cholesterol

Adults, 2008




References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).

Men, 2008



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).


Women, 2008



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885
Definitions:	% Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate).


Raised fasting blood glucose

Men, 2014



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A869?lang=en
Definitions:	Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication).


Women, 2014



References:	Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A869?lang=en
Definitions:	Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication).

Diabetes prevalence


Adults, 2024



References: Reproduced with kind permission International Diabetes Federation. IDF Diabetes Atlas, 11th edn. Brussels, Belgium:International Diabetes Federation, 2025. http://www.diabetesatlas.org

Ovarian Cancer


Women, 2022



Age:	20+
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Leukemia


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Liver and intrahepatic bile duct Cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000


Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Multiple Myeloma


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Indicence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Non Hodgkin Lymphoma


Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².

Women, 2022




Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².


Thyroid Cancer

Men, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

Women, 2022



Age:	20+
Area covered:	National
References:	Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24]
Definitions:	Incidence per 100,000

PDF created on November 20, 2026