

Report card Saudi Arabia




Contents	Page
Obesity prevalence	2
Trend: Obesity prevalence in Saudi Arabia adults, 1989-2019	5
Overweight/obesity by education	7
Overweight/obesity by age	9
Overweight/obesity by region	11
Overweight/obesity by socio-economic group	13
Insufficient physical activity	15
Estimated per capita fruit intake	18
Estimated per-capita processed meat intake	19
Estimated per capita whole grains intake	20
Mental health - depression disorders	21
Mental health - anxiety disorders	22
Oesophageal cancer	23
Breast cancer	25
Colorectal cancer	26
Pancreatic cancer	28
Gallbladder cancer	30
Kidney cancer	32
Cancer of the uterus	34
Raised blood pressure	35
Raised cholesterol	38
Raised fasting blood glucose	41
Diabetes prevalence	43
Contextual factors	44



Obesity prevalence

Adults, 2019

Obesity Overweight




Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².



Children, 2020-2022


Obesity Overweight





Infants, 2020

Overweight or obesity




Definitions:

=>+2SD



Obesity prevalence in Saudi Arabia adults, 1989-2019




Health-Indicators/Documents/World-Health-Survey-Saudi-Arabia.pdf (Accessed 16.11.2021)

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.



Women




Health-Indicators/Documents/World-Health-Survey-Saudi-Arabia.pdf (Accessed 16.11.2021)

Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.



Overweight/obesity by education


Adults, 2019



Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².



Children, 2006



growth standards.


Cutoffs: WHO



Overweight/obesity by age

Adults, 2019

Obesity Overweight




Indicators/Documents/World-Health-Survey-Saudi-Arabia.pdf (Accessed 16.11.2021)

Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².



Children, 2006



Cutoffs:


CDC



Overweight/obesity by region

Adults, 2019

Obesity Overweight




Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².



Children, 2020-2022

Obesity Overweight






Overweight/obesity by socio-economic group

Adults, 2019

Obesity Overweight




Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².



Children, 2006





Sample size:	1243	
Area covered:	Riyadh	
References:	Al Alwan İ, Al Fattani A, Longford N. The Effect of Parental Socioeconomic Class on Children's Body Mass Indices. Journal of Clinical Research in Pediatric Endocrinology. 2013;5(2):110-115. doi:10.4274/Jcrpe.898.	
Notes:	Prevalence of overweight and obesity by Income (Saudi Riyal/Month). Obesity and overweight were defined using the WHO	


es: Prevalence of overweight and obesity by income (Saudi Riyal/Month). Obesity and overweight were defined using the WHO 2007 growth standards.

Cutoffs: WHO




Insufficient physical activity

Adults, 2016



(18)30357-7


Men, 2016



ces: Guthold R, Stevens GA, Riley LM, Bull FC. Worldwide trends in insufficient physical activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1.9 million participants. Lancet 2018 <u>http://dx.doi.org/10.1016/S2214-109X</u> (18)30357-7



Women, 2016



 References:
 Guthold R, Stevens GA, Riley LM, Bull FC. Worldwide trends in insufficient physical activity from 2001 to 2016: a pooled


 analysis of 358 population-based surveys with 1.9 million participants. Lancet 2018 http://dx.doi.org/10.1016/S2214-109X

 (18)30357-7



Estimated per capita fruit intake







Estimated per-capita processed meat intake


2.5 2 1.5 g/day 1 0.5 South Sudan 0 Morocco Lebanon Bahrain Arghanistan Somalia Palestine Tunisia Iraq Iran Yemen UAE Sudan Syria Jordan Libya Saudi Arabia Pakistan Oman Kuwait Qatar Measured Survey type: Age: 25+ Global Burden of Disease, the Institute for Health Metrics and Evaluation http://ghdx.healthdata.org/ **References:** Definitions: Estimated per-capita processed meat intake (g per day)

Adults, 2017



Estimated per capita whole grains intake


Adults, 2017





Mental health - depression disorders

Adults, 2015



eferences: Prevalence data from Global Burden of Disease study 2015 (<u>http://ghdx.healthdata.org</u>) published in: Depression and Other Common Mental Disorders: Global Health Estimates. Geneva:World Health Organization; 2017. Licence: CC BY-NC-SA 3.0 IGO.


Definitions:

% of population with depression disorders



Mental health - anxiety disorders






References: Prevalence data from Global Burden of Disease study 2015 (<u>http://ghdx.healthdata.org</u>) published in: Depression and Other Common Mental Disorders: Global Health Estimates. Geneva:World Health Organization; 2017. Licence: CC BY-NC-SA 3.0 IGO.

Definitions:

% of population with anxiety disorders



Oesophageal cancer





Women, 2022






Breast cancer

Women, 2022

Age 20 Age 20+




25



Colorectal cancer

Men, 2022






Women, 2022






Pancreatic cancer

Men, 2022






Women, 2022





Gallbladder cancer

Men, 2022






Women, 2022






Kidney cancer

Men, 2022





Women, 2022





Cancer of the uterus


Women, 2022






Raised blood pressure

Adults, 2015




Men, 2015






Women, 2015






Raised cholesterol

Adults, 2008






Men, 2008






Women, 2008





Raised fasting blood glucose






Definitions:

Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication).




Women, 2014





Diabetes prevalence

Adults, 2021





Contextual factors

Disclaimer: These contextual factors should be interpreted with care. Results are updated as regularly as possible and use very specific criteria. The criteria used and full definitions are available for download at the bottom of this page.



Labelling

Is there mandatory nutrition labelling?	✓
Front-of-package labelling?	✓
Back-of-pack nutrition declaration?	✓
Color coding?	✓
Warning label?	×

Regulation and marketing	
Are there fiscal policies on unhealthy products?	\checkmark
Tax on unhealthy foods?	×
Tax on unhealthy drinks?	~
Are there fiscal policies on healthy products?	X
Subsidy on fruits?	×
Subsidy on vegetables?	×
Subsidy on other healthy products?	X
Mandatory limit or ban of trans fat (all settings)?	\checkmark
Mandatory limit of trans fats in place (all settings)?	~
Ban on trans-fats or phos in place (all settings)?	~
Are there any mandatory policies/marketing restrictions on the promotion of unhealthy food/drinks to	X
children?	
Mandatory restriction on broadcast media?	×
Mandatory restriction on non-broadcast media?	×
Voluntary policies/marketing restrictions on the promotion of unhealthy food/drinks to children?	v
Are there mandatory standards for food in schools?	X
Are there any mandatory nutrient limits in any manufactured food products?	~
Nutrition standards for public sector procurement?	X



Last updated September 13, 2022

PDF created on August 15, 2024