# Turkey



## Country report card - adults


*This report card contains the latest data available on the Global Obesity Observatory on overweight and obesity for adults. Where available, data on common and relevant obesity drivers and comorbidities are also presented.*

*View the latest version of this report on the Global Obesity Observatory at https://data.worldobesity.org/country/turkey-219/.*

| Contents | Page |
|---|---|
| Obesity prevalence | 3 |
| Trend: % Adults living with obesity in Turkey 1999-2023 | 5 |
| Trend: % Adults living with overweight or obesity in Turkey 1999-2023 | 10 |
| Overweight/obesity by age and education | 15 |
| Overweight/obesity by education | 22 |
| Overweight/obesity by age | 29 |
| Overweight/obesity by region | 31 |
| Overweight/obesity by age and socio-economic group | 34 |
| Overweight/obesity by socio-economic group | 38 |
| Overweight/obesity by age and limited activity | 45 |
| Double burden of underweight & obesity | 49 |
| Insufficient physical activity | 52 |
| Average weekly frequency of fast food consumption | 59 |
| Mental health - depression disorders | 66 |
| Mental health - anxiety disorders | 70 |
| Percent of population who cannot afford a healthy diet | 74 |
| Oesophageal cancer | 75 |
| Breast cancer | 78 |
| Colorectal cancer | 80 |
| Pancreatic cancer | 83 |
| Gallbladder cancer | 86 |
| Kidney cancer | 89 |
| Cancer of the uterus | 92 |
| Raised blood pressure | 94 |
| Raised cholesterol | 98 |
| Raised fasting blood glucose | 101 |
| Diabetes prevalence | 103 |
| Ovarian Cancer | 104 |
| Leukemia | 106 |
| Liver and intrahepatic bile duct Cancer | 111 |
| Multiple Myeloma | 114 |
| Non Hodgkin Lymphoma | 119 |
| Thyroid Cancer | 124 |

## Obesity prevalence

## Adults, 2025




| | |
|---|---|
| **Survey type:** | Self-reported |
| **Age:** | 18+ |
| **Area covered:** | National |
| **References:** | Eurostat 2025. Person distribution by body mass index, educational attainment level, sex and age. https://doi.org/10.2908/ILC_HCH10 [Accessed 20.04.26] |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

## % Adults living with obesity in Turkey 1999-2023

## Men




| | |
|---|---|
| **Survey type:** | Measured |
| **References:** | 1999: Delibasi, T., Karaaslan, Y., Ustun, I. et al. National prevalence of underweight, overweight and obesity in Turkey: cross sectional study of a representative adult population. cent.eur.j.med 2, 294–303 (2007). https://doi.org/10.2478/s11536-007-0024-4<br>2011: Chronic Diseases and Risk Factors Survey in Turkey. Republic of Turkey Ministry of Health 2013.<br>2017: National Household Health Survey – Prevalence of Noncommunicable Disease Risk Factors in Turkey 2017 (STEPS). Üner S, Balcılar M, Ergüder T editors. World Health Organization Country Office in Turkey, Ankara, 2018.<br>2023: National Household Health Survey – Prevalence of Noncommunicable Disease Risk Factors in Turkey 2023 (STEPS). Editors: Mehmet Enes Gökler, Hamit Harun BaÄ?cÄ±, Kanuni Keklik, Toker Erguder, Ä°layda. Third Ministry of Health, Ankara, 2025. |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

## Women




| | |
|---|---|
| **Survey type:** | Measured |
| **References:** | 1999: Delibasi, T., Karaaslan, Y., Ustun, I. et al. National prevalence of underweight, overweight and obesity in Turkey: cross sectional study of a representative adult population. cent.eur.j.med 2, 294–303 (2007). https://doi.org/10.2478/s11536-007-0024-4<br>2011: Chronic Diseases and Risk Factors Survey in Turkey. Republic of Turkey Ministry of Health 2013.<br>2017: National Household Health Survey – Prevalence of Noncommunicable Disease Risk Factors in Turkey 2017 (STEPS). Üner S, Balcılar M, Ergüder T editors. World Health Organization Country Office in Turkey, Ankara, 2018.<br>2023: National Household Health Survey – Prevalence of Noncommunicable Disease Risk Factors in Turkey 2023 (STEPS). Editors: Mehmet Enes Gökler, Hamit Harun BaÄ?cÄ±, Kanuni Keklik, Toker Erguder, Ä°layda. Third Ministry of Health, Ankara, 2025. |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

## % Adults living with overweight or obesity in Turkey 1999-2023

## Men




| | |
|---|---|
| **Survey type:** | Measured |
| **References:** | 1999: Delibasi, T., Karaaslan, Y., Ustun, I. et al. National prevalence of underweight, overweight and obesity in Turkey: cross sectional study of a representative adult population. cent.eur.j.med 2, 294–303 (2007). https://doi.org/10.2478/s11536-007-0024-4<br>2011: Chronic Diseases and Risk Factors Survey in Turkey. Republic of Turkey Ministry of Health 2013.<br>2017: National Household Health Survey – Prevalence of Noncommunicable Disease Risk Factors in Turkey 2017 (STEPS). Üner S, Balcılar M, Ergüder T editors. World Health Organization Country Office in Turkey, Ankara, 2018.<br>2023: National Household Health Survey – Prevalence of Noncommunicable Disease Risk Factors in Turkey 2023 (STEPS). Editors: Mehmet Enes Gökler, Hamit Harun BaÄ?cÄ±, Kanuni Keklik, Toker Erguder, Ä°layda. Third Ministry of Health, Ankara, 2025. |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

## Women




| | |
|---|---|
| **Survey type:** | Measured |
| **References:** | 1999: Delibasi, T., Karaaslan, Y., Ustun, I. et al. National prevalence of underweight, overweight and obesity in Turkey: cross sectional study of a representative adult population. cent.eur.j.med 2, 294–303 (2007). https://doi.org/10.2478/s11536-007-0024-4<br>2011: Chronic Diseases and Risk Factors Survey in Turkey. Republic of Turkey Ministry of Health 2013.<br>2017: National Household Health Survey – Prevalence of Noncommunicable Disease Risk Factors in Turkey 2017 (STEPS). Üner S, Balcılar M, Ergüder T editors. World Health Organization Country Office in Turkey, Ankara, 2018.<br>2023: National Household Health Survey – Prevalence of Noncommunicable Disease Risk Factors in Turkey 2023 (STEPS). Editors: Mehmet Enes Gökler, Hamit Harun BaÄ?cÄ±, Kanuni Keklik, Toker Erguder, Ä°layda. Third Ministry of Health, Ankara, 2025. |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

*Different methodologies may have been used to collect this data and so data from different surveys may not be strictly comparable. Please check with original data sources for methodologies used.*

## Overweight/obesity by age and education


## Adults, 2025



| | |
|---|---|
| **Survey type:** | Self-reported |
| **Area covered:** | National |
| **References:** | Eurostat 2025. Person distribution by body mass index, educational attainment level, sex and age. https://doi.org/10.2908/ILC_HCH10 [Accessed 20.04.26] |
| **Definitions:** | Level 0-2: Less than primary, primary and lower secondary education; Level 3-4: Upper secondary and post-secondary non-tertiary education; Level 5-8: Tertiary |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*


## Men, 2025



| | |
|---|---|
| **Survey type:** | Self-reported |
| **Area covered:** | National |
| **References:** | Eurostat 2025. Person distribution by body mass index, educational attainment level, sex and age. https://doi.org/10.2908/ILC_HCH10 [Accessed 20.04.26] |
| **Definitions:** | Level 0-2: Less than primary, primary and lower secondary education; Level 3-4: Upper secondary and post-secondary non-tertiary education; Level 5-8: Tertiary |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

## Women, 2025




| | |
|---|---|
| **Survey type:** | Self-reported |
| **Area covered:** | National |
| **References:** | Eurostat 2025. Person distribution by body mass index, educational attainment level, sex and age. https://doi.org/10.2908/ILC_HCH10 [Accessed 20.04.26] |
| **Definitions:** | Level 0-2: Less than primary, primary and lower secondary education; Level 3-4: Upper secondary and post-secondary non-tertiary education; Level 5-8: Tertiary |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

## Overweight/obesity by education

## Adults, 2025




| | |
|---|---|
| **Survey type:** | Self-reported |
| **Age:** | 18+ |
| **Area covered:** | National |
| **References:** | Eurostat 2025. Person distribution by body mass index, educational attainment level, sex and age. https://doi.org/10.2908/ILC_HCH10 [Accessed 20.04.26] |

| | |
|---|---|
| **Definitions:** | Level 0-2: Less than primary, primary and lower secondary education; Level 3-4: Upper secondary and post-secondary non-tertiary education; Level 5-8: Tertiary |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

## Men, 2025




| | |
|---|---|
| **Survey type:** | Self-reported |
| **Age:** | 18+ |
| **Area covered:** | National |
| **References:** | Eurostat 2025. Person distribution by body mass index, educational attainment level, sex and age. https://doi.org/10.2908/ILC_HCH10 [Accessed 20.04.26] |

| | |
|---|---|
| **Definitions:** | Level 0-2: Less than primary, primary and lower secondary education; Level 3-4: Upper secondary and post-secondary non-tertiary education; Level 5-8: Tertiary |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

## Women, 2025




| | |
|---|---|
| **Survey type:** | Self-reported |
| **Age:** | 18+ |
| **Area covered:** | National |
| **References:** | Eurostat 2025. Person distribution by body mass index, educational attainment level, sex and age. https://doi.org/10.2908/ILC_HCH10 [Accessed 20.04.26] |

| | |
|---|---|
| **Definitions:** | Level 0-2: Less than primary, primary and lower secondary education; Level 3-4: Upper secondary and post-secondary non-tertiary education; Level 5-8: Tertiary |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

## Overweight/obesity by age

## Adults, 2025




| | |
|---|---|
| **Survey type:** | Self-reported |
| **Area covered:** | National |
| **References:** | Eurostat 2025. Person distribution by body mass index, educational attainment level, sex and age. https://doi.org/10.2908/ILC_HCH10 [Accessed 20.04.26] |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

## Overweight/obesity by region

## Women, 2018-2019




| | |
|---|---|
| **Survey type:** | Measured |
| **Age:** | 15-49 |
| **Sample size:** | 6362 |
| **Area covered:** | National |
| **References:** | Hacettepe University Institute of Population Studies. (2019). 2018 Turkey Demographic and Health Survey. Hacettepe University Institute of Population Studies, T.R. Presidency of Turkey Directorate of Strategy and Budget and TÜBITAK, Ankara, Turkey. |

| | |
|---|---|
| **Notes:** | Demographic Health Survey data includes ever married women aged 15-49 years only and may include males aged 15-59. |
| | *Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².* |

## Overweight/obesity by age and socio-economic group


## Adults, 2019



| | |
|---|---|
| **Survey type:** | Self-reported |
| **Area covered:** | National |
| **References:** | Eurostat 2019. Available at https://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=hlth_ehis_bm1i&lang=en (last accessed 09.08.21). |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*


## Men, 2019



| | |
|---|---|
| **Survey type:** | Self-reported |
| **Area covered:** | National |
| **References:** | Eurostat 2019. Available at https://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=hlth_ehis_bm1i&lang=en (last accessed 09.08.21). |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

## Women, 2019




| | |
|---|---|
| **Survey type:** | Self-reported |
| **Area covered:** | National |
| **References:** | Eurostat 2019. Available at https://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=hlth_ehis_bm1i&lang=en (last accessed 09.08.21). |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

## Overweight/obesity by socio-economic group


## Adults, 2019



| | |
|---|---|
| **Survey type:** | Self-reported |
| **Age:** | 18+ |
| **Area covered:** | National |
| **References:** | Eurostat 2019. https://doi.org/10.2908/HLTH_EHIS_BM1I (Last accessed: 06.06.25) |
| **Notes:** | 1st Quintile (lowest income), 5th Quintile (highest income) |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*


## Men, 2019



| | |
|---|---|
| **Survey type:** | Self-reported |
| **Age:** | 18+ |
| **Area covered:** | National |
| **References:** | Eurostat 2019. https://doi.org/10.2908/HLTH_EHIS_BM1I (Last accessed: 06.06.25) |
| **Notes:** | 1st Quintile (lowest income), 5th Quintile (highest income) |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

## Women, 2019




| | |
|---|---|
| **Survey type:** | Self-reported |
| **Age:** | 18+ |
| **Area covered:** | National |
| **References:** | Eurostat 2019. https://doi.org/10.2908/HLTH_EHIS_BM1I (Last accessed: 06.06.25) |
| **Notes:** | 1st Quintile (lowest income), 5th Quintile (highest income) |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

## Overweight/obesity by age and limited activity


## Adults, 2014



| | |
|---|---|
| **Survey type:** | Self-reported |
| **Area covered:** | National |
| **References:** | Eurostat 2014 available at https://appsso.eurostat.ec.europa.eu/nui/submitViewTableAction.do (last accessed 06.10.21) |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*


## Men, 2014



| | |
|---|---|
| **Survey type:** | Self-reported |
| **Area covered:** | National |
| **References:** | Eurostat 2014 available at https://appsso.eurostat.ec.europa.eu/nui/submitViewTableAction.do (last accessed 06.10.21) |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

## Women, 2014




| | |
|---|---|
| **Survey type:** | Self-reported |
| **Area covered:** | National |
| **References:** | Eurostat 2014 available at https://appsso.eurostat.ec.europa.eu/nui/submitViewTableAction.do (last accessed 06.10.21) |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

## Double burden of underweight & obesity

## Adults, 2022




| | |
|---|---|
| **Survey type:** | Measured |
| **Age:** | 20+ |
| **References:** | NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population representative studies with 222 million children, adolescents, and adults. Lancet 2024; published online Feb 29. https://doi.org/10.1016/S0140-6736(23)02750-2. |
| **Notes:** | Age Standardised estimates |

| | |
|---|---|
| **Definitions:** | Combined prevalence of BMI<18.5 kg/m² and BMI>=30 kg/m² (double burden of underweight and obesity) |

## Insufficient physical activity


## Adults, 2022



| | |
|---|---|
| **Survey type:** | Self-reported |
| **Age:** | 18+ |
| **Area covered:** | National |
| **References:** | WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-) |

| | |
|---|---|
| **Definitions:** | Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent. |


## Men, 2022



| | |
|---|---|
| **Survey type:** | Self-reported |
| **Age:** | 18+ |
| **Area covered:** | National |
| **References:** | WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-) |

| | |
|---|---|
| **Definitions:** | Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent. |

## Women, 2022




| | |
|---|---|
| **Survey type:** | Self-reported |
| **Age:** | 18+ |
| **Area covered:** | National |
| **References:** | WHO (2024). Prevalence of insufficient physical activity among adults aged 18-years age-standardized estimate in 2022. Available at https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-insufficient-physical-activity-among-adults-aged-18-years-(age-standardized-estimate)-(-) |

| | |
|---|---|
| **Definitions:** | Percent of population attaining less than 150 minutes of moderate-intensity physical activity per week, or less than 75 minutes of vigorous-intensity physical activity per week, or equivalent. |

## Average weekly frequency of fast food consumption


## Adults, 2021-2024



| | |
|---|---|
| **Survey type:** | Self-reported |
| **Age:** | 15+ |
| **Sample size:** | 980 |
| **Area covered:** | National |
| **References:** | Global Diet Quality Project. 2024. "DQQ Results Dataset 2021-2024." Harvard Dataverse. https://doi.org/doi:10.7910/DVN/KY3W8A Licence: CC BY-NC-SA 4.0. Accessed at dietquality.org on October 15, 2025. |

| | |
|---|---|
| **Notes:** | For more global dietary survey data, visit the Global Diet Quality Project website at https://www.dietquality.org/ |
| **Definitions:** | Proportion of the population who consumed fast food during the previous day or night (self-reported) |


## Men, 2021-2024



| | |
|---|---|
| **Survey type:** | Self-reported |
| **Age:** | 15+ |
| **Sample size:** | 980 |
| **Area covered:** | National |
| **References:** | Global Diet Quality Project. 2024. "DQQ Results Dataset 2021-2024." Harvard Dataverse. https://doi.org/doi:10.7910/DVN/KY3W8A Licence: CC BY-NC-SA 4.0. Accessed at dietquality.org on October 15, 2025. |

| | |
|---|---|
| **Notes:** | For more global dietary survey data, visit the Global Diet Quality Project website at https://www.dietquality.org/ |
| **Definitions:** | Proportion of the population who consumed fast food during the previous day or night (self-reported) |

## Women, 2021-2024




| | |
|---|---|
| **Survey type:** | Self-reported |
| **Age:** | 15+ |
| **Sample size:** | 980 |
| **Area covered:** | National |
| **References:** | Global Diet Quality Project. 2024. "DQQ Results Dataset 2021-2024." Harvard Dataverse. https://doi.org/doi:10.7910/DVN/KY3W8A Licence: CC BY-NC-SA 4.0. Accessed at dietquality.org on October 15, 2025. |

| | |
|---|---|
| **Notes:** | For more global dietary survey data, visit the Global Diet Quality Project website at https://www.dietquality.org/ |
| **Definitions:** | Proportion of the population who consumed fast food during the previous day or night (self-reported) |


## Mental health - depression disorders

## Adults, 2023




| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26] |
| **Definitions:** | Estimated number living with depressive disorders per 100,000 population (20+ years of age) |

## Men, 2023



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26] |
| **Definitions:** | Estimated number living with depressive disorders per 100,000 population (20+ years of age) |


## Women, 2023



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26] |
| **Definitions:** | Estimated number living with depressive disorders per 100,000 population (20+ years of age) |


## Mental health - anxiety disorders

## Adults, 2023




| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26] |
| **Definitions:** | Estimated number living with anxiety disorders per 100,000 population (20+ years of age) |

## Men, 2023



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26] |
| **Definitions:** | Estimated number living with anxiety disorders per 100,000 population (20+ years of age) |


## Women, 2023



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Global Burden of Disease (GBD) Study 2023. Seattle, WA: IHME, University of Washington, 2025. Available from https://vizhub.healthdata.org/gbd-compare/ [Accessed 17.03.26] |
| **Definitions:** | Estimated number living with anxiety disorders per 100,000 population (20+ years of age) |

# Percent of population who cannot afford a healthy diet


## Adults, 2025



| | |
|---|---|
| **Area covered:** | National |
| **References:** | FAO. 2026. FAOSTAT: Cost and Affordability of a Healthy Diet (CoAHD). https://www.fao.org/faostat/en/#data/CAHD. Licence: CC-BY-4.0. [Last Accessed 12.10.26] |
| **Notes:** | The percent of the population whose food budget is below the cost of a healthy diet. |


## Oesophageal cancer

## Men, 2022



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Indicence per 100,000 |


## Women, 2022



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Indicence per 100,000 |

## Breast cancer


## Women, 2022



| | |
|---|---|
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Incidence per 100,000 |


## Colorectal cancer

## Men, 2022



| | |
|---|---|
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Incidence per 100,000 |


## Women, 2022



| | |
|---|---|
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Incidence per 100,000 |


## Pancreatic cancer

## Men, 2022



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Incidence per 100,000 |


## Women, 2022



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Incidence per 100,000 |


## Gallbladder cancer

## Men, 2022



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Indicence per 100,000 |


## Women, 2022



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Indicence per 100,000 |


## Kidney cancer

## Men, 2022



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Incidence per 100,000 |


## Women, 2022



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Incidence per 100,000 |

## Cancer of the uterus


## Women, 2022



| | |
|---|---|
| **Age:** | 20+ |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Incidence per 100,000 |


## Raised blood pressure

## Adults, 2019




| | |
|---|---|
| **Age:** | 30-79 |
| **References:** | World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26] |
| **Definitions:** | Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79. |

## Men, 2019



| | |
|---|---|
| **Age:** | 30-79 |
| **References:** | World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26] |
| **Definitions:** | Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79. |


## Women, 2019



| | |
|---|---|
| **Age:** | 30-79 |
| **References:** | World Health Organization. 2024. Hypertension: raised blood pressure (uncontrolled hypertension) among adults aged 30-79 years (%). Global Health Observatory. https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-raised-blood-pressure-among-adults-30-79-years [Accessed 16.03.26] |
| **Definitions:** | Prevalence of raised blood pressure (defined as having systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg) among adults aged 30-79. |


## Raised cholesterol

### Adults, 2008




| | |
|---|---|
| **References:** | Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885 |
| **Definitions:** | % Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate). |

## Men, 2008



| | |
|---|---|
| **References:** | Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885 |
| **Definitions:** | % Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate). |


## Women, 2008



| | |
|---|---|
| **References:** | Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A885 |
| **Definitions:** | % Raised total cholesterol (>= 5.0 mmol/L) (age-standardized estimate). |


# Raised fasting blood glucose

## Men, 2014



| | |
|---|---|
| **References:** | Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A869?lang=en |
| **Definitions:** | Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication). |


## Women, 2014



| | |
|---|---|
| **References:** | Global Health Observatory data repository, World Health Organisation, http://apps.who.int/gho/data/node.main.A869?lang=en |
| **Definitions:** | Age Standardised % raised fasting blood glucose (>= 7.0 mmol/L or on medication). |

# Diabetes prevalence


## Adults, 2024



**References:** Reproduced with kind permission International Diabetes Federation. IDF Diabetes Atlas, 11th edn. Brussels, Belgium:International Diabetes Federation, 2025. http://www.diabetesatlas.org

## Ovarian Cancer


## Women, 2022



| | |
|---|---|
| **Age:** | 20+ |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Incidence per 100,000 |

## Leukemia


## Men, 2022



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Incidence per 100,000 |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

## Women, 2022




| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Incidence per 100,000 |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*


## Liver and intrahepatic bile duct Cancer

## Men, 2022



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Incidence per 100,000 |


## Women, 2022



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Incidence per 100,000 |

## Multiple Myeloma


## Men, 2022



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Indicence per 100,000 |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

## Women, 2022




| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Indicence per 100,000 |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

## Non Hodgkin Lymphoma


## Men, 2022



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Incidence per 100,000 |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*

## Women, 2022




| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Incidence per 100,000 |

*Unless otherwise noted, overweight refers to a BMI between 25kg and 29.9kg/m², obesity refers to a BMI greater than 30kg/m².*


## Thyroid Cancer

## Men, 2022



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Incidence per 100,000 |

## Women, 2022



| | |
|---|---|
| **Age:** | 20+ |
| **Area covered:** | National |
| **References:** | Ferlay J, Ervik M, Lam F, Laversanne M, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F (2024).Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: https://gco.iarc.who.int/today, accessed [16.07.24] |
| **Definitions:** | Incidence per 100,000 |

*PDF created on November 5, 2026*