


Nigerija: Prevalencija pretilosti



Odrasle osobe, 2011-2012



Vrsta ankete:	Izmjereno
Dob:	18+
Veličina uzorka:	2983
Pokriveno područje:	Regionalno
Reference:	Chukwuonye II, Chuku A, Onyeonoro U, Ukegbu A, Anyabolu E, et al. (2015) Body Mass Index, Prevalence and Predictors of Obesity in Urban and Rural Communities in Abia State South Eastern Nigeria. J Diabetes Metab 6: 570. doi:10.4172/2155-6156.1000570 Copyright: © 2015 Chukwuonye II,
Bilješke:	Abia Region Years of survey not displayed in Chukwuonye II et al (2015), information found in Okpechi IG, Chukwuonye II, Tiffin N, Madukwe OO, Onyeonoro UU, Umeizudike TI, et al. (2013) Blood Pressure Gradients and Cardiovascular Risk Factors in Urban and Rural Populations in Abia State South Eastern Nigeria Using the WHO STEPwise Approach. PLoS ONE 8(9): e73403. https://doi.org/10.1371/journal.pone.0073403

Ako nije drukčije naznačeno, prekomjerna tjelesna težina odnosi se na BMI između 25 kg i 29,9 kg/m², a pretilost se odnosi na BMI veći od 30 kg/m².